

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

AN INVESTIGATION OF SOME DIFFERENCES BETWEEN

ASPECTS IN HILL COUNTRY

A thesis presented in partial fulfilment of the
requirements for the degree of Master of
Agricultural Science in Plant Science
at Massey University

Matthew Greg Lambert

1973

ABSTRACT

Climatic, edaphic and biotic variables were measured, over a twelve month period, at each of four aspects of a hill in the Southern Ruahine ranges. These variables were soil moisture status, soil temperature, air temperature, wind-speed, rainfall, soil nutrient status, sheep-dung deposition, and pasture botanical composition and productivity. Information on sunshine hours, maximum and minimum screen temperatures, relative humidity, and wind direction were obtained from the records of an adjacent meteorological station. Net radiation and potential evapotranspiration were calculated from meteorological data, and actual evapotranspiration from soil moisture data.

Large differences were recorded between aspects for most of the above mentioned variables. The wind during the observational period was a prevailing West/Northwesterly. Differences in net radiation between the north and south aspects were largest during the Winter and smallest during the summer months. In all cases the evapotranspiration values calculated were larger for the north than for the south aspect. Soil moisture tension differences were not detected during the winter months, but during the remainder of the year the north aspect was driest, followed by the east and west aspects, and the south aspect respectively. Differences between aspects, in terms of average monthly 4 cm. air temperature, were not apparent. However, large differences in the average monthly 4 cm. soil temperature of the various aspects were detected: during the January to August period the north aspect was warmest and the south coolest; during the October to December period the east aspect was warmest and the north and south aspects, which had similar average soil temperatures, were coolest.

The south and west aspect soils had greater nutritional limitations to plant growth than did the soils of the east and north aspects. This was probably due, at least in part, to nutrient transfer by grazing animals, and the differential action of soil-forming factors. Nitrogen mineralisation was closely associated with soil total nitrogen status, and was one of the main factors limiting pasture productivity. Soil moisture status was the other major limitation to pasture productivity. Pasture production during the observational period (346 days), for the east, south, west and north aspects respectively, was 9683, 3637, 2959 and 2771 kg./DM./ha. Some of the pasture species present were found

111

to be distributed in a definite pattern according to aspect, while for other species the pattern was indistinct. For a number of species no distribution pattern was detected. The patterns observed appeared to follow soil nutritional (especially mineral nitrogen) and soil moisture gradients.

Possible reasons for the above-mentioned differences, and some practical implications of these differences, are discussed.

ACKNOWLEDGEMENTS

Grateful acknowledgement is made to Mr E. Roberts for his supervision throughout this study, and to Mr D.A. Grant for suggesting this topic.

Thanks are also due to: Dr J.P. Kerr, Mr D.A. Grant, Mr J.S. Talbot, Dr R.H. Jackman, Mr P.R. Ball and Dr G.A. Wickham, for advice freely given, and the N.Z. Meteorological Service and the late Mr D.B. Edmond, for the loan of equipment.

This study was carried out while the author was holder of a State Services Commission Study Award.

CONTENTS

Acknowledgements

Contents

List of Figures

List of Tables

Introduction	1
Chapter I Literature Review	
1. Introduction	6
2. Microclimatic Differences between Aspects	7
3. Vegetational Differences between Aspects	13
4. Edaphic Differences between Aspects	16
5. Differences between Aspects due to Animals	17
Chapter II Methods and Materials	
(A) Introduction	18
(B) Experimental Methods	20
1. Microclimatic Factors	
1.1 Sunshine Hours, Wind Direction, Relative Humidity, Screen Minimum and Maximum Temperatures, and Rainfall.	
1.2 Wind-speed	
1.3 Rainfall	22
1.4 Soil Moisture Tension	
1.5 Soil and Air Temperatures	26
1.6 Net Radiation	31
1.7 Evapotranspiration	32
2. Soil Factors	34
2.1 Exchangeable Cations, Available Phosphorus, Total Nitrogen, Organic Carbon, Organic Matter Content, Cation Exchange Capacity, pH and Base Saturation	

2.2	Bulk Density	35
2.3	Available Nitrogen	
3.	Animal Factor	37
3.1	Amounts of Dung Deposition	
4.	Pasture Factors	
4.1	Botanical Composition and Percentage Bare Ground	
4.2	Pasture Production	40
(C)	Statistical Methods	41
1.	Microclimatic and Animal Factors	43
2.	Soil Factors	
2.1	Exchangeable Cations, Available Phosphorus, Total Nitrogen, Organic Carbon, Cation Exchange Capacity, pH and Base Saturation	
2.2	Mineral Nitrogen	
3.	Pasture Factors	44
3.1	Botanical Composition and Percentage Bare Ground	
3.2	Production	45
Chapter III Results		
1.	Microclimatic Factors	46
1.1	Wind-Speed	
1.2	Wind Direction	
1.3	Rainfall	
1.4	Net Radiation Estimates	
1.5	Evapotranspiration	
1.6	Soil Moisture Tension	51
1.7	Temperature	
2.	Soil Factors	57
2.1	Exchangeable Cations, Available Phosphorus, Total Nitrogen, Organic Carbon, Organic Matter Content, Cation Exchange Capacity, pH and Base Saturation	
2.2	Available Nitrogen	
3.	Animal Factor	62
4.	Pasture Factors	
4.1	Comparison of Point Analysis and Dry Weight Analysis	

4.2	Comparison of Botanical Composition and Percentage Bare Ground between Aspects	
4.3	Production Differences	66
Chapter IV Discussion		
1.	Microclimatic Factors	71
1.1	Wind-Speed and Wind Direction	
1.2	Rainfall	
1.3	Net Radiation and Evapotranspiration	72
1.4	Soil Moisture Status	73
1.5	Temperature	74
2.	Soil Factors and the Animal Factor	75
3.	Pasture Factors	78
3.1	Botanical Composition and Percentage Bare Ground	
3.2	Production Differences	80
4.	General Discussion and Practical Implications	82
Appendices		
		85
Bibliography		
		110

LIST OF FIGURES

I.1	Aerial View of Experimental Area and Surrounds	5
II.1	Aerial View of Experimental Area	19
II.2	South Aspect 'microclimate site'	21
II.3	East Aspect	
II.4	West Aspect	23
II.5	'F.F.F' Gypsum Blocks	
II.6	Sensor Unit for 4 cm. Air Temperature Measurements	28
II.7	Sensor Unit for 120 cm. Air Temperature Measurements	
II.8	Device used for Measuring Slope	42
III.1	Average Wind Speed: January to June	47
III.2	Average Wind Speed: July to December	48
III.3	Rainfall	50
III.4	Soil Moisture Tension: January to March	52
III.5	Soil Moisture Tension: April to September	53
III.6	Soil Moisture Tension: October to December	54
III.7	Annual Course of 4 cm. Air Temperature	55
III.8	Annual Course of 4 cm. Soil Temperature	56
III.9	Herbage Production on East Aspect, 4.4 to 28.8.72	68
III.10	Herbage Production on North Aspect, 4.4 to 28.8.72	
III.11	Herbage Production on South Aspect, 4.4 to 28.8.72	69
III.12	Herbage Production on West Aspect, 4.4 to 28.8.72	

LIST OF TABLES

I.1	Botanical Composition on Sunny and Shady Faces (Suckling, 1959)	15
I.2	Average Annual D.M. Production for the Period 1952 - 56 (Suckling, 1959)	
III.1	Wind Direction Summary for 1972	49
III.2	Rainfall Totals for Each Aspect	
III.3	Net Radiation and Evapotranspiration Estimates for North and South 'microclimate sites'	
III.4	Results of Chemical Analyses of Soil Samples taken 30.3.72	(58)
III.5	Ambient Mineral Nitrogen Levels	60
III.6	Net Nitrogen Mineralisation during Incubation	
III.7	Analysis of Variance of Net Mineralisation Means for Spring Incubation	
III.8	Net Nitrogen Mineralisation for Source Aspects during Spring Incubation	
III.9	Coefficients for Regression of Final Mineral Nitrogen Content on Final Moisture Content	61
III.10	Moisture Contents of Soil Cores used for Mineral Nitrogen Determinations	
III.11	Analysis of Variance of Final Moisture Contents of Soil Samples Incubated during Spring	
III.12	Total Nitrogen Values for Soil Cores using in Spring Ambient Level Determinations	63
III.13	Centigrade Hours above 5°C at 4 cm. in Soil, for Winter and Spring Incubations	
III.14	Dung Deposition At North, South, East and West Aspects	
III.15	Comparison of Botanical Composition as determined by Point Analysis and Dry Weight Analysis	64
III.16	Botanical Composition and Percentage Bare Ground as measured by Point Analysis	65
III.17	Productivity on Slopes of 5° to 30°	67
III.18	Comparison of Production Estimates using 25 cm ² Herbage Samples and 1m ² Herbage Samples	
III.19	Comparison of Production Estimates made for Two Slope Classes	70

INTRODUCTION

This section is intended as an introduction to the study which is to be described, and as a means of introducing certain general ideas concerning aspect differences.

"Ecology is the science concerned with living organisms, both plant and animal, in relation to their environment or habitat" (Levy, 1970). Such a definition might not find favour with the more pedantic members of the ecological discipline, but does manage to convey the basic meaning of the term 'ecology' ie. the study of organisms 'at home' (Odum, 1959). Odum (1959) states that many terrestrial ecosystems "have a particularly complex structure involving numerous species, marked stratification and variable physical environment, In local situations there is much to be gained from singling out a restricted component (for study)..... At the same time it is important that whole systems be studied simultaneously, since certain fundamental interrelationships can not be readily determined by piecemeal study." Any study of variation in a single environmental factor is purely descriptive if only that factor is measured eg. a record of pasture botanical composition differences between aspects, or between any contrasting areas, does not provide any explanation for the differences observed. A complete analysis of edaphic, climatic and biotic differences between aspects would be an extremely large and complicated undertaking. It is, however, possible to elucidate some of the interrelationships existing between the various environmental factors, through the study of selected variables. The selection of the variables which were examined in this study was based on two criteria: firstly, feasibility of measurement, and secondly, the likelihood of differences occurring between aspects, as judged by discussion and by perusal of relevant literature.

The literature reviewed in this study is of two broad categories: firstly, literature dealing with general concepts of aspect differences, and secondly, specific literature, in most cases describing or reviewing experimental studies and the results obtained. The latter category is dealt with in Chapter I and the former is reviewed briefly below.

Microclimatic⁽¹⁾ differences are known to exist between land surfaces which are inclined in different directions, ie. between surfaces of different aspect (Warming, 1909; Braun-Blanquet, 1932; Geiger, 1965; Chang, 1968). Variation in the amount of direct solar radiation

(1) see discussion of the terms 'climate' and 'microclimate' in Introduction to Chapter I.

received by sloping surfaces is one of the main reasons for the existence of these microclimatic differences (Geiger, 1965). The associated differing energy input has a large influence on the energy balances which exist at the surfaces of the various aspects encountered in the hill country situation. A simple energy balance for a bare soil surface may be written as:

$$R_N + B + L + V = 0 \text{ (Geiger, 1965)}$$

where:

R_N = net radiation.

B = sensible heat loss to the ground.

L = latent heat loss due to evaporation.

V = sensible heat loss to the air.

The interrelationships of the elements of this balance are discussed in many texts, eg. Geiger, 1965; Slatyer and McIlroy, 1961; Kunn, 1966. The magnitude of R_N is determined by the lack of balance between net incoming short-wave and net outgoing long-wave radiation. Near the ground, incoming short-wave radiation consists of a direct solar beam, and diffuse sky radiation, the latter coming from the whole hemisphere, although more intense in directions close to that of the sun itself (Slatyer and McIlroy, 1961). The proportion of incoming short-wave radiation which is diffuse varies from about 10% under clear-sky conditions, to 100% under overcast conditions. Thus, differences in short-wave radiation input between aspects will be greatest under clear-sky conditions. The proportion of the incoming short-wave radiation which is reflected from the surface it strikes is known as the albedo. The fraction not reflected is the net incoming short-wave radiation. The net outgoing long-wave radiation, mentioned above, represents the difference between terrestrial and sky long-wave radiation.

The R_N component of the above balance is expended by sensible heat transfers which directly affect the temperature of the soil and air, and as latent heat during the evaporation of water. Slatyer and McIlroy (1961) point out that of all the variables involved in the energy balance only solar radiation can be regarded as at all independent of the others. Under steady conditions the factors of the balance adjust to come to an equilibrium. Geiger (1965) notes that the situation described above is often complicated by the horizontal transfer, or advection, of heat from surrounding areas, thus introducing an additional factor to the energy balance. Although the balance written above is for a bare soil surface, the addition of a vegetative cover to the surface would little alter the concepts developed in the above discussion.

It was previously noted that variation in the amount of direct solar radiation received by sloping surfaces is one of the main reasons for the existence of microclimatic differences in hill country. Wind also

plays a role, in that air movement is involved in the determination of the magnitude of the components of the energy balance. Local winds may arise in mountainous hill country due to regions of different temperature (Geiger, 1965). However, in the New Zealand situation of prevailing winds and overall proximity to the coastline, and especially in the smaller-scale hill country of the North Island, it is probable that wind is only modified, rather than caused, by local topographic variation. Slatyer and McIlroy (1961) state that "wind is of considerable importance in microclimate, both as an element in its own right and because it is of such influence on the atmospheric structure of temperature and humidity". They conclude that the main effect of wind is to reduce extremes of variation in temperature and humidity, both in time and in space.

Local variation in rainfall may also occur in hill country. Geiger (1965) notes that the climate of slopes facing in different directions is affected to a large extent by moisture conditions, as well as radiation and wind, and that the smaller the topographic scale the more the local precipitation is determined by the wind field. In discussing small-scale precipitation differences, Geiger points out that more precipitation is found on the leeward than on the windward side of hills, especially where wind speeds are high. "By general agreement" (Geiger, 1965) precipitation is measured using horizontally disposed collecting surfaces, yet considerable controversy now exists as to the relative merits of using horizontal as opposed to tilted collecting surfaces (Geiger, 1965; Yates, 1970). Doubt also exists as to the proportions of recorded differences between aspects which are attributable to actual precipitation differences and those which are due to wind effects on rain gauge catch (Rodda, 1966; Jackson and Aldridge, 1972).

Edaphic differences⁽¹⁾ between aspects might arise for any of a number of reasons. Ross (1971), in his review of aspect as a soil forming factor, notes that aspect, acting through microclimate and its effects on organisms, plays an important role in the ecology and thus soils of hillsides. Sears et al (1948) give figures for the annual nutrient turnover, via sheep excreta, for grazed pasture, and remark that fertilizer programmes should be constructed with reference to the transfer of dung and urine from one part of a paddock to another. Hilder (1966) discusses nutrient accumulation on stock camps under a sheep grazing regime, and also equates this accumulation with a loss from other parts

(1) Soil moisture status and temperature will be considered as climatic rather than edaphic factors; see Introduction to Chapter I.

of the paddock. The potential for nutrient transfer between aspects appears to exist, and coupled with the probable influence of microclimatic and pasture differences between aspects on animal grazing behaviour, such a transfer might be expected to lead to differences in soil and pasture characteristics between aspects. Grazing animals have a number of important effects on plant communities, these effects being due to physical damage, defoliation and the deposition of excreta (Spedding, 1971). Rumball and Grant (1972) go so far as to state that "the trampling, grazing, voiding animal is one of the major determinants of pasture composition."

From basic ecological concepts, variation in pasture structure, composition and productivity would be expected to be associated with the differing environments existing at different aspects.

The detection of variation in edaphic, biotic and climatic factors due to aspect, formed the basis of the study described herein. A hill in the Southern Ruahine ranges of the North Island was selected as an experimental site. Measurements of selected climatic, edaphic and biotic factors were made at each of the north, south, east and west aspects, over a twelve month period. A description of each of the selected factors was the prime aim of the study and the collection of data was conducted with this in mind. However, a secondary aim did exist, namely to elucidate, where possible, the interrelationships existing between the environmental and pasture variables measured.

Throughout this thesis common names have been used, whenever possible, in referring to plant species. The corresponding botanical names have been noted the first time each common name is used in the text. A complete list of the common and botanical names of species encountered in this study is given in Appendix 10.




FIGURE I.1 Aerial View of Experimental area and Surrounds