

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

The interaction of multiple bodies and water waves

with the application to the motion of ice floes

A thesis presented in partial fulfillment of the requirements for the degree of

Master of Science

in

Mathematics

at Massey University, Albany,

New Zealand

Malte Andreas Peter

2002

Abstract

To understand the propagation of water waves through arrays of floating or (fully or partially) submerged bodies it is necessary to know how these bodies interact with each other under the influence of ambient waves. However, the conventional full diffraction calculation of the scattered wavefields of many interacting bodies requires a considerable computational effort.

In this thesis, a method is developed which makes it possible to quickly calculate the wave scattering of many interacting floating or (fully or partially) submerged, vertically non-overlapping bodies of arbitrary geometry in water of infinite depth. It extends Kagemoto and Yue's analysis for axisymmetric bodies in finite depth.

The idea of this method is to expand the water velocity potential into its cylindrical eigenfunctions such that the scattered potentials of the bodies are defined by a set of coefficients only. Representing the scattered wavefield of each body as an incident wave upon all other bodies, a linear system of equations for the coefficients of the scattered wavefields is derived.

Diffraction transfer matrices which relate the coefficients of the incoming wavefield upon a single body to those of its scattered wavefield play an important role in the process. The calculation of the diffraction transfer matrices for bodies of arbitrary shape requires the representation of the infinite depth free surface Green's function in the eigenfunctions of an outgoing wave. This eigenfunction expansion will be derived from the equivalent finite depth Green's function.

An important application of this interaction method is the propagation of ocean waves through fields of ice floes which can be modelled as floating flexible thin plates. Meylan's method of solution is used to calculate the motion of a single ice floe from which the solutions for multiple interacting ice floes are computed.

While the interaction theory will be derived for general floating or submerged bodies, particular examples are always given for the case of ice floes. Results are presented for ice floes of different geometries and in different arrangements and convergence tests comparing the finite and the infinite depth method are conducted with two square interacting ice floes where full diffraction calculations serve as references.

Acknowledgements

I would like to thank Dr Michael Meylan for the great support he has shown me during the completion of this thesis. He has been an excellent supervisor and I have learnt very much from his comments and suggestions.

I would also like to thank the German National Scholarship Foundation for financially supporting my research at Massey University.

Contents

1	Introduction	1
2	Mathematical formulation and solution for a single ice floe	5
2.1	The models	5
2.1.1	The water	6
2.1.2	The ice floe	8
2.1.3	Non-dimensionalisation	9
2.1.4	Transformation into an integral equation	10
2.2	Solving the problem with one ice floe	11
2.2.1	Simplification of the integral equation	11
2.2.2	Solving the plate equation	12
3	The eigenfunction expansion of the water velocity potential	15
3.1	Local cylindrical coordinates	15
3.2	Derivation of the eigenfunction expansion of the potential	16
3.2.1	The case of water of finite depth	17
3.2.2	The case of infinitely deep water	19
3.2.3	Derivation of the infinite depth solution by taking the limit of the finite depth one	21
4	Different representations of the Green's functions	25
4.1	Graf's addition theorem	25
4.2	The eigenfunction expansion in finite depth	27

4.3	The eigenfunction expansion in infinite depth	29
4.4	Numerical comparison to existing representations	33
4.5	Numerical aspects of the Green's function in a fixed coordinate system . . .	37
4.5.1	Numerical results for one single ice floe	39
5	The interaction of multiple bodies	45
5.1	Eigenfunction expansion of the scattered potential	46
5.2	The interaction in water of finite depth	48
5.3	The interaction in water of infinite depth	50
5.4	Explicit details	53
5.4.1	Calculation of the diffraction transfer matrix	53
5.4.2	The diffraction transfer matrix of rotated bodies	55
5.4.3	Eigenfunction expansion of the ambient wavefield	56
5.4.4	Solving the resulting system of equations	57
5.4.5	Calculation of the surface displacement of the ice floes	58
6	Numerical results and convergence	59
6.1	Numerical results	60
6.2	Convergence	60
6.2.1	Water of finite depth	63
6.2.2	Water of infinite depth	68
7	Summary and Discussion	77
7.1	The eigenfunction expansion of the free surface Green's function	78
7.2	Advantages of the interaction methods over the full diffraction calculation .	78
7.3	Comparison of the finite and the infinite depth methods	79
A	Nomenclature	81
B	Contents of the CD-ROM	85

List of Figures

2.1	Displacement of the ice floe under incident wave of wavelength 2	14
2.2	Displacement of the ice floe under incident wave of wavelength 1	14
3.1	Coordinate systems of the body	16
3.2	The graph of $\tanh x$	21
3.3	The graph of $x \tan x$	23
4.1	Triangle for Graf's addition theorem	26
4.2	Relation between two bodies (plan view)	26
4.3	Setting with a single point source at (s, φ, c) (plan view)	28
4.4	The graphs of the integrands in equations (4.16)	36
4.5	Decay of the product of the Bessel and Hankel function	39
4.6	Surface displacement in the vicinity of the ice floe under ambient incident wave of wavelength 2	41
4.7	Scattered wavefield in the vicinity of the ice floe under ambient incident wave of wavelength 2	42
4.8	Surface displacement in the vicinity of the ice floe under ambient incident wave of wavelength 1	43
4.9	Scattered wavefield in the vicinity of the ice floe under ambient incident wave of wavelength 1	44
6.1	Surface displacement of interacting ice floes	61
6.2	Surface displacement of interacting ice floes	62
6.3	Development of the errors as the number of propagating modes is increased	65

6.4	Development of the errors as the number of decaying angular components is increased	65
6.5	Development of the errors as the number of roots of the dispersion relation is increased	66
6.6	$ \alpha^2 - k^2 $ plotted versus the wavelength and the depth	67
6.7	Surface displacement of the ice floes and the water in their vicinity, $O_1 = (-1.4, 0)$, $O_2 = (1.4, 0)$	69
6.8	Development of the errors as the number of propagating modes is increased	70
6.9	Development of the errors as the number of decaying angular components is increased	71
6.10	Development of the errors as the number of points in the discretisation of the continuous variable is increased	72
6.11	Development of the errors as the distance between the ice floes increased . .	73
6.12	Surface displacement of the ice floes and the water in their vicinity, $O_1 = (0, -1.4)$, $O_2 = (0, 1.4)$	74
6.13	Surface displacement of the ice floes and the water in their vicinity, $O_1 = (-1.4, -0.6)$, $O_2 = (1.4, 0.6)$	75