

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**Use of turnips to reduce potassium
accumulation on areas receiving Farm
Dairy Effluent**

**A thesis presented in partial fulfilment of the
requirements for the degree of**

Master of Applied Science

in

Soil Science

at Massey University, Palmerston North, New Zealand.



Massey University

Monica Salazar

2006

Abstract

Land treatment of farm dairy effluent (FDE) on small areas of intensive of dairy farms has enriched soils with nutrients particularly K. Solving the problem solely by increasing the area allocated for land treatment requires large investment in pump, pipes and irrigator infrastructure. A less costly strategy, of sowing and grazing a summer turnip on the land treatment area in order to redistribute K to the pasture area is evaluated in this thesis. A survey (February 2006) showed that in the Manawatu region turnip crop yields (8 to 17t DM.ha⁻¹) provided profitable feed for dairy cows, were a suitable re-grassing strategy and if harvested, removed 350 to 700 kg K ha⁻¹ from the soil. In the summer of 2005/06, a turnip (*Brassica rapa* cv. Barkant) trial was established after permanent pasture on a Pallic soil (pH 6.5, Olsen P 35.2 ug. g⁻¹, exchangeable K⁺ 0.7, Ca²⁺ 6.3, Mg²⁺ 1.4 me/ 100 g soil). The following treatments pre-plant fertiliser only (38 kg N ha⁻¹, 25 kg P ha⁻¹ and 25 kg K ha⁻¹), pre-plant fertiliser plus side-dressed urea at 40 DAS (46 kg N ha⁻¹) and pre-plant fertiliser plus 5 x 10 mm FDE applications (57kgha⁻¹) all produced similar final dry matter yields (8 t DM ha⁻¹) at 100 days after sowing (DAS). Leaf was the largest component of dry matter and had higher K concentrations (4.6 and 6.8% K in the control and FDE treatments respectively) than bulb (3 and 4 %K in the control and FDE treatments respectively). The ratio of leaf to bulb dry matter however varied for each different treatment. Side-dressed urea and FDE treatments produced the largest leaf biomass and reached maximum yields earlier by 75 DAS and 64 DAS, respectively and generated more K removal at harvest (339, 428 & 537 kg K ha⁻¹ at 75 DAS and 316, 372 & 490 kg K ha⁻¹ at 100 DAS for pre-plant only, urea & FDE treatments, respectively).

The lack of yield response to N partially resulted from crop uptake of between 107 and 114 kg N ha⁻¹ from mineralisable soil N. The dynamic N crop model N-able predicted that extra side-dressed N would not increase turnip yield but in the absence of pre-plant N (38 kg N ha⁻¹) the turnips would yield 7.4 t DM ha⁻¹ at 100 DAS. The use of the N-

able model demonstrated a need for a decision support model to assist farmers in choosing appropriate N fertiliser application rates.

A simple model was created to simulate how the grazing cow can transfer K from turnip paddocks (part of a FDE treatment block) to other parts of the farm. The model simulation of 490 cows on a mixed diet of 4kg DM turnips and 12 kg DM pasture predicted that the grazing of turnips (8t DM ha⁻¹ crop) would result in the net transfer of significant quantities (>170 kg K ha⁻¹) of K from land growing turnips to other parts of the farm. To cause net transfer to occur the allocated turnip dry matter must be grazed in the shortest time possible and the cows returned to pasture after short milking times.

Acknowledgements

I would like to express my gratitude to the following people.

First of all, I am extremely thankful to my chief supervisor Dr. Mike Hedley who gave me from the beginning his guidance and devoted patience and technical support during the development of my thesis. I admire the extra committed time that he used to help me to finish on time this thesis. Also, I want to say thanks to Carolyn for being kind, noble and good person who with her family supported me during my first month living in NZ. My especial thank to Dr. Dave Horne for his collaboration, guidance, suggestions and changes during the last phase of this study.

Lots of thanks to Dr. Colin Holmes and his wife Dorothy for their friendship, fondness and wise advice. I will miss you Colin, Dorothy and Mia.

Thanks to the team of technicians and staff who was helped me all the time during my field experiment and laboratory analysis particularly: Bob Toes, Ian Furkert, Glenys Wallace, Ross Wallace, Ann West and Leighton Parker. Special gratefulness to James Hanly for his efforts in many aspects of the field related research and to Ko for being a nice Korean person.

A special thank to Mike Bretherton and Moira Hubbard for being such nice people (awesome, generous and charming), who offered their friendship and understanding at any time during my stay in New Zealand.

Many thanks to my postgraduate mates at TVL and especially to Jagrati Singh for being an extraordinary friend and such a nice girl, thanks for being with me in my up and downs moments. I have really enjoyed your friendship. Rita Bhandral, Muron Banabas, Janice Asing and Asoka Senarath with whom I shared knowledge, many beautiful moments, friendship and help in many ways.

Big thanks and love to Eduardo Amador and Julia McCormick who have been a real family for me, caring for me a lot at any time. To my friends Johanna and Hector Ballesteros who were the first friends in NZ cheering me up and encouraging me from the beginning. A special hug and greeting to Family Paine (Manuel, Geovana, Elizabeth and Gino) for their invaluable friendship, tenderness and love, which supported me since the time, I met them. Big hugs to Gaby and Marcos Fernandez for being good friends and a lovely couple and gave me their assistance at any time.

A special thanks and hug to Martin Baz for being a good and sensible person and friend who encouraged, supported and was with me and formed part of my life in the last four months of my stay in New Zealand.

Finally I gratefully acknowledge to the New Zealand Aid Scholarship and the NZ government for this great opportunity to study in this wonderful country.

Table of Contents

Abstract	i
Acknowledgements	iii
Table of Contents	v
List of Tables.....	x
List of Figures	xiii
List of Plates.....	xv
Introduction.....	1
1.1 Intensification of Dairying and the accumulation of potassium under land treatment of FDE.....	1
1.2 Solution to this problem: Strategies to reduce the accumulation of K on areas receiving FDE	3
1.2.1 Extension of effluent area	3
1.2.2 Crop nutrient removal	4
1.3 Research Objectives and Thesis Structure	7
1.4 References	7
Chapter 2.....	10
2.1 Brief review of FDE management in dairy farm systems.....	10
2.2 Environmental concern about effluent.....	10
2.3 Chemical composition of Farm Dairy Effluent.....	12
2.4 Best land-based wastewater treatment	15
2.5 Nutrient budgeting model	18
2.5.1 Area required to reduce the loads of K and N.....	18
2.5.2 Nutrient removal in crops.....	19
2.6 Review of literature on turnips, their impact on the performance of intensive dairy farms and as a crop to reduce high K concentrations on effluent paddocks	21
2.7 Introduction.....	21

2.8	Use of turnips on New Zealand dairy farms	21
2.9	Turnip yield and variety responses	22
2.10	Factors associated with yield	24
2.11	Milk solids.....	25
2.12	Effect of turnips on milk characteristics	26
2.13	Economic analysis of its use	27
2.14	Approaches to use turnips with FDE application.....	29
	2.14.1 Chemical characteristics.....	29
	2.14.2 Response to irrigation	31
	2.14.3 Responses to effluent	32
2.15	Summary	32
2.16	References	34
Chapter 3		40
3.1	A survey of seven turnip crops growing in the Palmerston North area.	40
3.2	Introduction	40
3.3	Objectives.....	41
3.4	Methods.....	42
	3.4.1 Location and Soil Description.....	42
	3.4.2 Soil types.....	42
	3.4.3 Field sampling and laboratory procedure.....	43
	3.4.5 Soil and Plant analysis	44
	3.4.5.1 Soil analysis	44
	3.4.5.2 Plant analysis.....	45
3.5	Results and discussion.....	45
	3.5.1 Turnip yields	46
	3.5.2 Factors influencing yield.....	47
	3.5.2.1 Soil nutrient status.....	47
	3.5.3 Climatic Conditions	49
	3.5.3.1 Temperature and solar radiation.....	49
	3.5.3.2 Plant available water	51
	3.5.4 Crop management	53

3.6	Turnip potential to remove K from paddocks	56
3.6.1	Plant nutrient concentrations.....	56
3.7	Conclusions	61
3.8	References	62
Chapter 4.....		65
4.1	A field experiment evaluating growth and nutrient uptake pattern of turnips (<i>Brassica rapa</i> cv. Barkant).....	65
4.2.	Introduction	65
4.3	Objectives.....	66
4.4	Materials and methods	66
4.4.1	Trial Location and soil type	66
4.4.2	Cultivation and crop establishment.....	67
4.4.3	Soil and Plant analysis	68
4.4.4	Effluent irrigation and N application	68
4.4.6	FDE nutrient concentrations	71
4.4.7	Plant sampling.....	72
4.4.8	Statistical analyses	73
4.5	Results and discussion.....	73
4.5.1	Dry matter yield and growth rate	73
4.5.1.1	The advantage of attaining maximum yield early.....	79
4.5.2	Plant Nutrient Concentration.....	80
4.5.3	Uptake and removal	83
4.5.4	Soil Nutrient Concentration	85
4.6	Conclusions	89
4.7	References	90
Chapter 5.....		93
5.1	Towards improved Nitrogen fertiliser recommendations for Turnip crops using N-able model.....	93
5.2	Introduction	93
5.3	Objective	94

5.4	Method	94
5.4.1	Model selection	94
5.5	Results & Discussion	97
5.5.1	Plant N uptake and dry matter yield.....	97
5.5.2	Soil N Mineralization rate.....	98
5.5.3	Discussion.....	101
5.6	Conclusions	103
5.7	References	103
Chapter 6.....		105
6.1	Mass balance modelling of potassium transfers resulting from mixed grazing of turnips and pasture during a summer period in the Manawatu District.....	105
6.2	Introduction	105
6.3	Objectives.....	106
6.4	Method (Calculation) assumptions	107
6.4.1	Case study Site No 4 Dairy	107
6.4.2	K loss model.....	108
6.4.3	The development of parameters and algorithms in the model	108
6.4.3.1	K ingested by cows	108
6.4.3.2	K product losses	109
6.4.3.3	K transfer losses	110
6.4.3.4	K returns in excreta	111
6.4.3.5	Net Return per cow/day	112
6.4.3.6	Net Returns per ha.....	112
6.5	Boundary conditions of variable inputs and inputs for No. Dairy	114
6.5.1	Daily intake of turnips.....	114
6.6	Results	115
6.6.1	Case study summer grazing turnips at No. 4 Dairy.....	115
6.6.2	General application of the model	117
6.6.2.1	Reduction of K transfer loss by reduced milking time.....	117

6.6.2.2 Net transfer of K from turnip paddocks depends on time allowed for grazing	118
6.6.2.3 Influence of Turnip K concentration on rate of K transfer	121
6.7 Conclusions	122
6.8 References	122
Summary	124
Appendix	127

List of Tables

Table 1.1 Nutrient content of different crops that could be grown to remove excess K from the soil	5
Table 1.2 The effect of crop type and yield on the removal of K from the soil.....	5
Table 1.3 Approximate cost per kg DM produced and the harvesting time of crops	6
Table 2.1 Problems caused by mismanagement of FDE application to the land.....	11
Table 2.2 Chemical and Physical characteristics of FDE from aerobic ponds	13
Table 2.3 Mean concentration of K in herbage (grass and white clover) on N-K plots and percent changes in Ca, Na and P due to K fertiliser application (McNought <i>et al</i> , 1973)	14
Table 2.4 Effect of dairy farm effluent application on exchangeable cations in soil (Taken from Bolan <i>et al</i> , 2004)	14
Table 2.5 The mean depth of farm dairy effluent (FDE) applied by the irrigator at each of the irrigation events, the soil moisture deficit at the commencement of irrigation, the average from all plots over three lactation seasons (2000/01 – 2002/03), and the maximum drainage from any single plot as a proportion of the applied FDE (i.e., representing the “worst case” scenario). -, no data (Taken from Houlbrooke <i>et al</i> . 2004c)	16
Table 2.5 Research and approximated turnip yield from 1994/95 to up to date.....	23
Table 2.7 Crude Protein (%) and metabolizable energy (MJME/kg DM) from New Zealand studies of turnips for dairy cows.....	29
Table 2.8 Chemical characteristics of turnips grown in Australia.....	31
Table 3.1 Location of the survey farms.	42
Table 3.2 Leaf, bulb and total dry matter production of Turnip crops surveyed in the Palmerston North area in March 2006.....	47
Table 3.3 Soil pH, Olsen P, exchangeable cations and extractable nitrogen values for Turnip crops surveyed in the Palmerston North area in March, 2006.....	48
Table 3.4 Normal ranges of soil pH, Olsen P, exchangeable cations and CEC values recommended for optimum Turnip crop growth (source: Hill Laboratories).....	49

Table 3.5 Summary of monthly climatic data obtained from Palmerston North weather station, November, 2005 – March, 2006.....	50
Table 3.6 Total climatic data obtained from Manawatu station, Palmerston North Nov, 2005 – March, 2006.....	50
Table 3.7 Sowing and harvesting dates, varieties, days after sowing, seed and plant density, grazed area and grazing time for Turnip crop surveyed in the Manawatu Region, Nov, 2005 – March, 2006.	55
Table 3.8 Fertilization, seed cultivation method, and ground preparation for Turnip crop surveyed in the Manawatu Region.	55
Table 3.9 Soil type and nutrient concentration in leaves of Turnip crop surveyed in the Manawatu Region, Nov, 2005 – March, 2006.....	57
Table 3.10 Soil type and nutrient concentration in bulbs by Turnip crop surveyed in the Manawatu Region, Nov, 2005 – March, 2006.....	57
Table 3.11 Nutrient removals by leaves on Turnip crop surveyed in the Manawatu Region, Nov, 2005 – March, 2006.	59
Table 3.12 Nutrient removals by bulbs on Turnip crop surveyed in the Manawatu Region, Nov, 2005 – March, 2006.	59
Table 3.13 Total Nutrient removals by the whole plant in Turnip crops surveyed in the Manawatu Region, Nov, 2005 – March, 2006.....	60
Table 4.1. Harvest dates, and timing of FDE irrigations during the trial.....	70
Table 4.2 Nutrient concentrations of FDE at point of application (FDE pumped from aerobic pond of a two pond system at Massey University’s Dairy Farm No 4).	71
Table 4.3 Quantity of nutrients applied to plots in FDE (kg ha ⁻¹).	72
Table 4.4. Total dry matter production in turnips Var. Barkant ¹ . Farm No 4 – Massey University.	74
Table 4.5. Leaf and bulb growth rates in turnips Var. Barkant. Farm No 4 – Massey University.	78
Table 4.6. Nutrient concentration in turnip cv. Barkant ¹ at 75 and 100 days after sowing Farm No 4 – Massey University.	81
Table 4.7 Concentration of the major nutrients in turnip leaves determined by Hill Laboratories (1998)	83

Table 4.8 Nutrient removal at 75 and 100 DAS by leaves and bulbs in turnips cv. Barkant ¹ . Farm No 4 – Massey University.....	84
Table 4.9 Total Nutrient removal at 75 and 100 DAS by the whole plant cv. Barkant ¹ . Farm No 4 – Massey University.....	85
Table 4.10. Soil pH, Olsen P, exchangeable cations and extractable mineral N in the top 0-150 mm soil depth at different times during the growth of turnips (var. Barkant) ¹ grown at Massey University’s No 4. Dairy Farm.....	86
Table 4.11 Harvest dates, soil gravimetric water content and timing of FDE irrigations during the trial.	89
Table 6.1 General information for Massey No 4 Dairy farm for the summer period 2005-2006.....	107
Table 6.2 The default K concentrations for feed and products used in the simple K balance model (Summer period 2005/06)	109
Table 6.3 Assumptions used in the simple K balance model, based on information from the Massey University No.4 dairy farm in the summer period 2005/06.....	115
Table 6.5 Daily transfers of potassium calculated by the simple model and expressed per cow and per hectare of turnips (grazed in 3.5 days by 490 cows at 4kgDM/cow/day).....	116
Table 6.5 Estimated K accumulations in the effluent blocks according to the milking time when the herd is in one mob of 490 or is divided into 2 mobs of 245 cows each.....	118
Table 6.6 Estimated K transferred from one hectare of turnips to pasture when time spent by cows in the turnip paddock (A) and time plus dry matter allocation (B) increases	119
Table 6.7 Estimated area (ha) to be grazed to transfer K to pasture paddocks per ha of turnip grazed.....	120

List of Figures

Figure 2.1 Key parameters and components for successful deferred irrigation of farm dairy effluent.....	17
Figure 3.1 Rainfall and soil water deficit obtained using N-Able and data from the Palmerston North weather station. Nov, 2005 – March, 2006.	52
Figure 3.4 Soil total mineral nitrogen at harvest and nitrogen concentrations in turnips leaves and bulbs in the Palmerston North region during the season Nov, 2005 – March, 2006.....	58
Figure 3.5 Total nutrient removal and turnip yield for K, and N in Turnip crop surveyed in the Manawatu Region, during the season Nov, 2005 – Mar, 2006....	61
Figure 4.1 Soil water balance and FDE irrigation. Farm No 4 – Massey University.....	69
Figure 4.2 The development of leaf and bulb dry matter yield for Turnips (cv. Barkant) grown in the control (a), urea (b) and FDE (c) treatments at Massey University’s No. 4 Dairy Farm – (Season 2005 – 06).....	75
Figure 4.3 The dry matter yield (a), leaves (b) shoots for Turnips (cv. Barkant) at Massey University’s No. 4 Dairy Farm – (Season 2005 – 06).....	77
Figure 4.4 Land-use schedule to illustrate the advantage of early grazing of turnips and early re-grassing.....	80
Figure 4.4 Change in growth rate of (total dry matter) turnip cv. Barkant and soil exchangeable soil K as days after sowing (DAS) increased.	87
Figure 4.5 Change in growth rate of (total dry matter) turnip cv. Barkant and soil ammonium (NH_4^+) and nitrate –N (NO_3^-) content as days after sowing (DAS) increased.	87
Figure 5.1 Flow diagram of N able model (with explanatory text from the web site)....	95
Figure 5.2 N-uptake (a) and dry matter yield (b) predicted (lines) and original data (symbols) of the Control (38 kg N ha^{-1} at planting), Urea (38 kg N ha^{-1} at planting plus 46 kg N ha^{-1} side-dressed) and FDE (38 kg N ha^{-1} at planting plus 57 kg N ha^{-1} as FDE) treatments and simulated yield without nitrogen application from the turnip trial conducted on the No.4 Dairy farm Massey University.	99

Figure 5.3 Simulated soil mineral nitrogen and average measured soil mineral nitrogen content in top 150 mm of the three treatments (control, urea and effluent) of the turnip trial.	100
Figure 6.1 The potassium cycle during mixed feeding of turnips and pasture for dairy systems	106
Figure 6.2 Estimated net K transferred from turnips paddocks to pasture paddocks per ha of turnip grazed as plant K concentration varies with fertiliser treatment (Control: 3.88% K, Urea: 4.66% K; FDE: 5.81% K).....	121

List of Plates

Plate 4.1 Farm dairy effluent (FDE) irrigations during the season 2005/2006. Farm No 4 – Massey University.	70
---	----

Introduction

1.1 Intensification of Dairying and the accumulation of potassium under land treatment of FDE

The intensification of dairy farming in New Zealand over the last 10 years has resulted in an increase in the average stocking rate of 10% i.e. from 2.48 cows ha⁻¹ to 2.75 cows ha⁻¹ (LIC, 2003/04). By 1999/00, the top 25% of farmers (herd size > 300 cows) had also increased the amount of supplementary feed purchased so that the average amount of supplement brought onto the farm was 670 kg DM cow⁻¹ or 2069 kg DM ha⁻¹ (Leslie, 2002). This increased use of supplementary feed and fertiliser has led to greater inputs of nutrients into dairy farm systems.

Increased stocking rate leads to the generation of greater volumes of farm dairy effluent (FDE). By 2003/04, approximately 63 million m³ year⁻¹ of FDE was being generated (LIC, 2003). Mismanaged irrigation of FDE to land can create a number of problems (Bolan et al., 2004; Houlbrooke et al., 2004) including:

- High application rates generating runoff and drainage of partially treated FDE, which pollute surface and ground waters.
- Excessive FDE loads leading to N and K enrichment of topsoils increasing the risk to animal health, plant quality and soil quality.
- Nutrient enrichment of winter drainage water
- Wet easily, pugged soils
- Localized areas that receive higher rates of effluent which exceed the soil's infiltration rate. Therefore, patches of saturated soil, preferential flow of FDE through the soil profile, soil structure damage, poor plant growth, etc.

Therefore, nutrient enrichment on land receiving FDE becomes a potential risk. The nutrient concentration in FDE varies widely from farm to farm (Longhurst *et al.*, 2000) but typically the N, P and K concentrations range from 181 to 400 mg N l⁻¹, 40 to 80 mg P l⁻¹ and 164 to 705 mg K l⁻¹, respectively (Longhurst *et al.*, 2000). Whilst high and poorly timed application rates of N and P in FDE can cause undesirable N and P enrichment of drainage (Houlbrooke *et al.*, 2004) and surface waters (Monaghan *et al.*, 2004), the high concentration of K in FDE can cause excessive K accumulation in soils of areas that receive FDE. Paddocks that are irrigated with FDE at the appropriate N loading rate for regional council consent (150-200 kg N ha⁻¹) accumulate and presumably leach large amounts of K (Bolan *et al.*, 2004). This results in pastures on effluent paddocks with high K contents and reduced Mg and Ca contents (Bolan *et al.*, 2004). For example, the nutrient budgeting tool 'Overseer® Nutrient Budgets 2' predicted that K was added as effluent at rates of 354 and 154 kg ha⁻¹ y⁻¹ to the effluent areas of two dairy farms in North Otago (3.7 cows ha⁻¹, Lynch, 2006) and Waikato (4.7 cows ha⁻¹, Payze, 2006), respectively. The variation in concentration depended partially on the area of the effluent blocks on each farm. Cows left for a long time to graze these paddocks may suffer from metabolic disorders (Hypocalcaemia and Hypomagnesaemia).

Problems caused by application of FDE to wet soils are being addressed by FDE storage and deferred irrigation. This system is advocated by most regional councils (Houlbrooke *et al.*, 2004). The management of deferred irrigation has been improved by the use of irrigators, such as the k-line system, that are able to apply small depths (less than 5 mm) of FDE to soils (Houlbrooke *et al.*, 2006). While such modification will minimise the risk of N and P loss to the aquatic environment, additional strategies are required to reduce the K loading and accumulation in soils.

1.2 Solution to this problem: Strategies to reduce the accumulation of K on areas receiving FDE

1.2.1 Extension of effluent area

The current criterion for applying FDE to land is based on Regional Council regulations for appropriate N loading of which $200 \text{ kg N ha}^{-1} \text{ y}^{-1}$ is the limit (Heatley, 1996). The land area recommended for FDE treatment is currently 4 ha per 100 cows (Heatley, 1996), but a new proposal suggests that the area should be based on the proportion of dung and urine that is collected from the herd. This can be estimated from the average time a cow spends off the paddock in the collecting yards and milking shed and/or feed pad (Hedley *et al.*, 2004). To prevent K accumulation in soil, the annual K loading rate should not exceed the annual K maintenance requirement of grazed pasture. This requires irrigation of FDE to a larger proportion of the farm (30-40%) than is currently used to meet the N requirements (10 -15 %). A number of case studies where areas allocated for land-treatment of FDE are described below.

Hedley *et al.*, (2004) described a farm in the Bay of Plenty with 580 cows and an original effluent area of only 19 ha. It was proposed to increase the FDE area to 43 ha (30% of the whole farm) to improve management of FDE. Nutrient application rates to the expanded area were reduced by up to 50%. Of course enlarging the effluent area requires increased expenditure on an upgraded irrigation system. In this case study, the predicted cost of installing additional irrigation on the 43 ha plus annual running costs was around \$8917 per year, which allowed a saving of \$9400 per year in fertiliser costs (Hedley *et al.*, 2004).

Two dairy farm case studies (in North Otago and Waikato) used the Overseer® nutrient budgeting model to assess the impact of nutrient management recommendations on efficiency of nutrient use in, and nutrient loss from, the effluent blocks. The two farms,

where the soil nutrients levels were already equal to or greater than optimum agronomic value for pasture, increased the area of effluent application up to the total farm area. The North Otago dairy farm reduced the application rates of nutrients by about 87%, by use of the new irrigation system. However, the predicted investment in the new irrigation system was very large (Lynch, 2006). On the other farm (Waikato) extending the application area to be equal to the whole farm area decreased the nutrient content applied to the effluent paddocks by about 53% (Payze, 2006). The difference in the sizes of the nutrient reductions in the two farms were due to the different stocking rate (3.7 and 4.7 cows ha⁻¹) and the size of the previous effluent blocks (25 and 85 ha).

1.2.2 Crop nutrient removal

Hedley *et al.*, (2002) proposed that the harvesting of silage, or hay, to remove K from the FDE area will avoid the need for excessively large land-treatment areas. For instance, Overseer® was used to simulate the ability of maize silage (spring planting) followed by barley/triticale (autumn planting) to remove nutrients for FDE paddocks on a Bay of Plenty farm (Hedley *et al.*, 2004). If these two crops yielded 27 and 5 ton DM ha⁻¹, respectively, the combined nutrient removal was predicted to be 414 kg N ha⁻¹, 61 kg P ha⁻¹ and 288 kg K ha⁻¹. Annual grass silage harvesting in spring and summer, which yields a total dry matter production of 6 t DM ha⁻¹, will remove less nutrient at 187 kg N ha⁻¹, 27 kg P ha⁻¹ and 133 kg K ha⁻¹ (Hedley *et al.*, 2004). Another problem in using grass silage as a nutrient removal strategy is that in many areas of New Zealand, wet soils in winter and spring delay the commencement of 'safe' effluent irrigation to late spring and summer. The window of opportunity to harvest hay and silage is reduced and crop sizes of 3.5 to 5 t DM ha⁻¹ are expected. Therefore, K extraction and removal can remain low relative to inputs in the FDE.

Summer crops such as maize and turnips offer the opportunity to produce larger amounts of dry matter with higher metabolisable energy content (Table 1.1) than conserved pasture (Table 1.2) and therefore are attractive as fodder crops for summer dry areas. In addition, for the same growing period, the higher yields of maize silage and turnips remove more nutrients than pasture silage (Table 1.2). Turnips, in particular,

have higher tissue K concentrations (Table 1.1) and remove markedly more K when harvested (grazed). For example, at a yield of 10 t DM ha⁻¹, turnips extract 10% less N and P but 40% more K and S than a crop of maize yielding 18 t DM ha⁻¹. If the turnips and maize had similar yields (18 t DM/ha) then the turnips would remove 60% more N and P and 150% more K and S (Table 1.2)

Table 1.1 Nutrient content of different crops that could be grown to remove excess K from the soil

	DM (%)	MJ ME/kg DM	CP (%)	Nutrients (%)					
				N	P	K	S	Ca	Mg
Pasture Silage/Baleage	35	9.5	16.0	3.5	0.29	2.2	0.22	0.70	0.18
Maize silage	33	10.3	8.0	1.3	0.23	1.2	0.13	0.25	0.18
Turnip crop	10	12.5	13.0	2.1	0.37	3.0	0.33	1.75	0.23

Source: Holmes *et al.*, 2002. Milk production from pasture. Principle and practices.

Table 1.2 The effect of crop type and yield on the removal of K from the soil

	Yield t ha ⁻¹	Nutrients (kg ha ⁻¹)			
		N	P	K	S
Pasture Silage/Baleage	2 - 4	70 - 140	6 - 12	44 - 88	4 - 8
Maize silage	18	234	41.4	216	23.4
Turnip crop	10	210	37	300	33
Turnip crop	18	378	66.6	540	59

[†] Source: The silage and baleage yields are from the Dexcel website (2006) www.dexcel.co.nz; the maize and turnip yields are from Holmes C. (Personal comm.).

Factors such as the timing of growth, available finance, and plant nutritive qualities would also affect farmers' decisions as to which crop to grow. At 0.08 – 0.12 \$/kg DM, turnips have the lowest cost per kg DM produced. Therefore, turnips could be a “best option” for this new strategy of nutrient harvest by a crop because, unlike maize, they require no specialized machinery nor added contractors costs to plant and harvest (feed to cows). However, many dairy farmers prefer to buy maize silage rather than cultivate (Holmes pers com, 2005).

Table 1.3 Approximate cost per kg DM produced and the harvesting time of crops

	Cost \$/kg DM	Harvest time (weeks)
Pasture Silage/Baleage	0.40 - 0.45	4 - 8
Maize silage	0.15 - 0.21	24 - 28
Turnip crop	0.08 - 0.12	10 - 12

Source: www.dexcel.co.nz

Forage crops such as turnips are grown for the following reasons:

- Provide a large quantity of high quality feed during summer months to increase milk production at this time (decreasing the fall in milk production that normally occurs at this time from 14 % to 7% month (McGrath *et al.*, 1998; Daniels, 1995).
- A pasture renewal strategy to introduce new productive pasture species free of endophytes, and clean the soil of pests, weeds and diseases;
- Improve soil structure and surface micro-topography by levelling or draining during the cultivation phase.
- Less stressful time for the farmer.

In addition, they could be sown on effluent areas and on/off grazed to aid the redistribution of effluent borne K around the other grazed areas of the farm.

1.3 Research Objectives and Thesis Structure

The objective of the research presented in this thesis is to provide more information on the role that a summer turnip crop could play in redistribution of K from soils that have been enriched with K by FDE application. This study involves a survey of turnip crops grown in the Palmerston North region, a field trial evaluating the K removal potential of Barkant turnips grown under FDE irrigation and two exercises in modelling the crop N requirement and K transfer by cows grazing a mixed diet of turnips and pasture.

This thesis has the following objectives:

- Review how intensive dairy farming increases the concentration of nutrients in effluent paddocks.
- Survey local turnip crops for yield and nutrient content at harvest.
- Measure the ability of turnips (*Brassica rapa* cv. Barkant) to take-up K under normal fertilization and FDE irrigation
- Use of N-able model to assess the amount of N fertiliser that is required to achieve high turnip yield.
- Build a simple model to evaluate the K transfer potential of different grazing strategies for cows on a mixed diet of turnips and pasture.

1.4 References

- Bolan, N. S., Horne, D. J., & Currie, L. D. (2004). Growth and chemical composition of legume-based pasture irrigated with dairy farm effluent. *New Zealand Journal of Agricultural Research*, 47, 85-93.
- Daniels, N. (1995). Summer forage crop survey. *Dairy Farming Annual*, 47: 32-40.

- Dexcel. (2005). Integrating turnips into your farm system. Dexcel FarmFact 5-27. Updated September, 2005. <http://www.dexcel.nz/farmfacts>.
- Dexcel. (2005). Barkant Turnips. Feeding the crop. Dexcel FarmFact 5-9 Updated August, 2005. <http://www.dexcel.nz/farmfacts>.
- Heatley, P. (1996). Dairy and the environment. In *Managing farm dairy effluent. Land application. Dairy and the Environment Committee: Palmerston North, New Zealand.* pp 2.1 - 2.8.
- Hedley, M.J., Dodd, M. and Vercoe, R. (2004). Juggling the supplement plus fertiliser nutrient balance - A responsible approach. In (I.M.Brookes ed.) *Proceedings of the 2nd Dairy3 Conference.* Vol. 2: Massey University, Palmerston North New Zealand. pp 49-60.
- Hedley, M. J., Horne, D., Hanly, J., Furkert, I and Toes, B. (2002). Deferred irrigation of dairy shed effluent to an artificially drained soil. In *Annual final report Massey University Agricultural Research-Foundation (MUARF).* pp 17-20.
- Houlbrooke D. J., Horne D. J., Hedley M. J. and Snow, V.O. (2004). A review of literature on the land treatment of farm-dairy effluent in New Zealand and its impact on water quality. *New Zealand Journal of Agricultural Research* 47: 499-511.
- Houlbrooke, D.J., Smith, M.R., and Nicolson, C. (2006). Reducing contaminant losses following application of farm dairy effluent to land using a K-line irrigation system. In *Implementing sustainable nutrient management strategies in agriculture* (Eds. L.D. Curie and J.A Hanly). Occasional Report No. 19. Fertilizer and Lime Research Centre, Massey University, Palmerston North, New Zealand. pp 290-331.
- Longhurst, R. D., Roberts, A. H. C., & O'Connor, M. B. (2000). Farm dairy effluent: a review of published data on chemical and physical characteristics in New Zealand. *New Zealand Journal of Agricultural Research*, 43, 7-14.

- Leslie, M. (2002). Efficiency for Economic Success. *Proceedings of the Farming of the Conference*, 373, 24 - 29.
- LIC. (2003). Dairy Statistics, *Livestock Improvement Corporation*. Hamilton, New Zealand.
- Lynch, B. (2006). Nutrient budget case study - North Otago dairy farm, In *implementing sustainable nutrient management strategies in agriculture* (Eds. L.D. Curie and J.A. Hanly). Occasional report No. 19. Fertiliser and Lime Research Centre, Massey University, Palmerston North, New Zealand. pp 238-242.
- McGrath, D. F., Dawson, J. E., Thomson, N. A., & Simons, H. P. (1998). More summer milk - The opportunities identified. *Proceedings of the Ruakura Dairy Farmers' Conference* : 85-95.
- Monaghan, R. M and Smith, L. C. (2004). Minimising surface water pollution resulting from farm dairy effluent application to mole-pipe drained soils. II. The contribution of preferential flow of effluent to whole-farm pollutant losses in subsurface drainage from a West Otago dairy farm. *New Zealand Journal of Agricultural Research* 47: 417-428.
- Payze, A. (2006). Nutrient budget case study - Waikato dairy farm, In *implementing sustainable nutrient management strategies in agriculture* (Eds. L.D. Curie and J.A Hanly). Occasional Report No. 19. Fertilizer and Lime Research Centre, Massey University, Palmerston North, New Zealand. pp 242-247.