

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

STUDIES ON THE IN VIVO CLEAVAGE AND  
THE IN VITRO CULTURE OF NEW ZEALAND  
ROMNEY SHEEP OVA

A thesis presented in partial fulfilment of the requirements for  
the degree of Master of Agricultural Science  
at  
Massey University

HARRY ROBIN TERVIT

1967

## A C K N O W L E D G E M E N T S

The candidate is indebted to his supervisor Dr M.F. McDonald for the guidance and assistance given during the course of this study.

Special thanks are due to Professor A.L. Rae and Dr F.R. Cockrem for advice in statistical analysis of data and to Dr R.E. Munford for advice in the analysis of data from the bioassay of gonadotrophins.

Grateful acknowledgement is made to Mr C. Muir for technical assistance; to Mr P.H. Whitehead and Mr W.R. Fairhall for the care of animals; to Miss M.G. Campbell and the library staff for promptly procuring the reference articles used in this work; to Miss D.J. Scott and Mr G.H. Burns for the photographic work; to Mr E. Ormsby and Mr B. Thatcher for laboratory assistance; to the staff of the local Freezing Works and Abattoir; to Mrs J. Fisher for taking special care in typing this thesis; and to my wife who spent many hours reading this manuscript.

The financial assistance of the Department of Agriculture and of a New Zealand Wool Board Scholarship is gratefully acknowledged.

T A B L E   O F   C O N T E N T S

<u>Chapter</u>		<u>Page</u>
	ACKNOWLEDGEMENTS	
	LIST OF TABLES	
	LIST OF FIGURES	
I	<u>INTRODUCTION</u>	1
II	<u>REVIEW OF LITERATURE</u>	3
	PRODUCTION OF OVA	3
	OVA TRANSFER	4
	OVA CULTURE AND STORAGE	7
	OVA CLEAVAGE	9
	OVA ABNORMALITIES	11
III	<u>MATERIALS AND METHODS</u>	13
	EXPERIMENTAL ANIMALS	13
	MANAGEMENT OF ANIMALS	13
	EXPERIMENTAL PLAN	13
	HORMONES AND TREATMENTS	17
	RECOVERY OF OVA	18
	CULTURE OF OVA	18
	EXAMINATION OF OVA	20
	TRANSFER OF OVA	21
	SLAUGHTER OF EWES AND EXAMINATION OF REPRODUCTIVE ORGANS	22

<u>Chapter</u>		<u>Page</u>
IV	<u>PRODUCTION OF OVA FOLLOWING HORMONAL STIMULATION OF DONOR EWES</u>	23
	INDUCTION OF SUPEROVULATION AND MATING OF DONOR EWES	23
	1. Length of oestrous cycle following P.M.S. injection during the breeding season	23
	2. Ovarian response	23
	3. Recovery of ova	24
	4. Fertilisation of recovered ova	26
	PRODUCTION OF OVA FROM HORMONALLY STIMULATED EWES IN THE EARLY AND LATE ANOESTROUS SEASONS	28
	1. Early anoestrous	28
	2. Late anoestrous	29
V	<u>CLEAVAGE OF OVA IN VIVO</u>	30
	NUMBERS OF CLEAVED, UNCLEAVED, FERTILISED OR UNFERTILISED OVA RECOVERED	30
	CLEAVAGE OF FERTILISED OVA	31
	1. Shortest and longest interval after the onset of oestrus to the recovery of each cleavage stage	31
	2. Mean interval after the onset of oestrus to the recovery of each cleavage stage	31
	3. Cleavage of ova recovered from the Fallopian tube unflushed at laparotomy 1, but flushed at laparotomy 2 (Treatment B - Figure 1)	32
	4. Cleavage of ova transferred into one or both Fallopian tubes at laparotomy 1 and recovered at laparotomy 2 (Treatments A and B - Figure 1)	34

<u>Chapter</u>		<u>Page</u>
VI	<u>CULTURE, ABNORMALITIES AND DIMENSIONS OF OVA</u>	36
	CULTURE OF OVA	36
	1. Media	36
	2. Cleavage of cultured ova	36
	3. Recovery of ova from the culture chambers	38
	ABNORMALITIES OF OVA	39
	1. Abnormalities of ova recovered at laparotomy	39
	2. Abnormalities of ova after culture	40
	DIMENSIONS OF OVA	41
VII	<u>STUDIES ON THE SURVIVAL OF TRANSFERRED OVA</u>	43
	TRANSFERS IN THE 1966 BREEDING AND 1966-67 EARLY ANOESTROUS SEASONS	43
	1. Control transfer of ova	44
	2. Transfer of ova after 24 hours culture	45
	3. Transfer of ova after 48 hours culture	46
	TRANSFERS IN THE 1966-67 LATE ANOESTROUS AND 1967 BREEDING SEASONS	46
	1. Control transfer of ova	47
	2. Transfer of ova after 24 hours culture	48
	3. Transfer of ova after 48 hours culture	48
	FACTORS AFFECTING SURVIVAL OF OVA IN CONTROL TRANSFERS	49
	1. Number of corpora lutea present in the ovaries of the recipient ewe	49
	2. State of recipient genital tract at transfer	49
	3. Number of ova transferred	50
	4. Cleavage stage of transferred ova	50

<u>Chapter</u>		<u>Page</u>
	5. Synchronisation of stage of development of transferred ova and recipient genital tract	50
	6. Period ova held <u>in vitro</u> before transfer	50
	MEASUREMENT OF EMBRYOS	51
	1. Measurement of 23-25 day old embryos	51
	2. Measurement of 44-101 day old foetuses	51
VIII	<u>DISCUSSION</u>	53
	PRODUCTION OF OVA FOLLOWING HORMONAL STIMULATION OF DONOR EWES	53
	THE IN VIVO CLEAVAGE OF FERTILISED OVA	62
	CULTURE, ABNORMALITIES AND DIMENSIONS OF OVA	65
	1. Culture of ova	65
	2. Abnormalities of ova	67
	3. Dimensions of ova	72
	SURVIVAL OF TRANSFERRED OVA	73
	1. Control transfers	75
	2. Transfer of cultured ova	78
IX	<u>SUMMARY</u>	83
	<u>REFERENCES</u>	90
	<u>APPENDIX 1 and 2</u>	95

L I S T      O F      T A B L E S

<u>Table</u>		<u>Page</u>
1	Oestrous cycle length following injection of P.M.S. on day 12 or 13 of the cycle	f 23
2	Ovarian response of ewes treated with P.M.S. in the breeding and anoestrous seasons	f 23
3	Ovarian response after transformation	f 24
4	Effect of dose of P.M.S. on the percentage of ova recovered and on the percentage of recovered ova fertilised per ewe	f 25
5	Effect of time of P.M.S. injection relative to progestagen withdrawal on interval to onset of oestrus and on ovarian response	f 28
6	Ovarian response of ewes treated in the late anoestrous season	f 29
7	Recovery and fertilisation of ova in the late anoestrous season	f 29
8	Numbers of cleaved, uncleaved, fertilised or unfertilised ova recovered	f 30
9	Shortest and longest interval after the onset of oestrus to recovery of each cleavage stage	f 31
10	Mean interval after the onset of oestrus to recovery of each cleavage stage	f 31
11	Effect of dose of P.M.S. on the mean interval from onset of oestrus to recovery of each cleavage stage	f 31
12	Effect of season on the interval to recovery of 2- cell ova	f 31
13	Cleavage rate of ova in the experimental Fallopian tube	f 32
14	Interval from the onset of oestrus to recovery of 6- to 8- cell and 8- cell ova from the experimental tubes and from tubes at laparotomy 1	f 33

f refers to following page.



<u>Table</u>	<u>Page</u>	
15	Effect of double laparotomy in treatment B, and transfer of ova at laparotomy 1 before recovery at laparotomy 2, on the percentage of ova recovered per ewe	f 34
16	Control cleavage rate and cleavage rate of ova transferred at laparotomy 1 and recovered at laparotomy 2	f 34
17	Interval from the onset of oestrus to recovery of 6- to 8- cell and 8- cell ova after transfer, and at laparotomy 1	f 35
18	Normal ova development and development of abnormalities relative to the initial cleavage stage of ova cultured for 24 hours	f 37
19	Comparison of the maximum times for cleavage of ova <u>in vitro</u> with the cleavage interval <u>in vivo</u>	f 38
20	The types and incidence of abnormal ova recovered at laparotomy	f 39
21	Effect of interval from onset of oestrus to laparotomy on the incidence of abnormal unfertilised ova	f 39
22	Effect of culture length on the incidence of ova abnormalities	f 40
23	Effect of initial cleavage stage of ova on the incidence of abnormalities after culture	f 40
24	Dimensions of ova	f 41
25	Mean blastomere volume and total volume of ova cytoplasm relative to the stage of ova cleavage	f 42
26	Success of ova transfer experiments conducted in the breeding and early anoestrous seasons	f 44
27	Synchronisation of stage of development of transferred ova and recipient genital tracts	f 44
28	Success of ova transfer experiments conducted in the late anoestrous and breeding seasons	f 47
29	Synchronisation of stage of development of transferred ova and recipient genital tracts	f 47
30	Synchronisation of onset of oestrus in donor and recipient ewes and ovarian response of recipient ewes	f 47

TablePage

31	Effect of number of recipient corpora lutea on the success of transfer experiments	f 49
32	Effect of state of recipient genital tract on the success of transfer experiments	f 49
33	Effect of number of ova transferred on the success of transfer experiments	f 49
34	Effect of cleavage stage of the transferred ova on the success of transfer experiments	f 49
35	Age and crown-rump length of each embryo	f 51

L I S T      O F      F I G U R E S

<u>Figure</u>		<u>Page</u>
1	Experimental plan for 1966 breeding season	14
2	Experimental plan for 1966-67 anoestrous season	15
3	Experimental plan for 1967 breeding season	16
4	Dismantled and assembled dialysis chamber	f 18
5	Mean percentage of ova recovered per ewe relative to the number of corpora lutea in the ovaries of that ewe	f 25
6	Mean percentage of ova recovered per ewe relative to the number of corpora lutea plus large follicles in the ovaries of that ewe	f 25
7	Mean percentage of ova recovered per ewe relative to the interval from onset of oestrus to laparotomy	f 26
8	Mean percentage of recovered ova fertilised per ewe relative to the number of corpora lutea in the ovaries of that ewe	f 26
9	Mean percentage of recovered ova fertilised per ewe relative to the number of corpora lutea plus large follicles in the ovaries of that ewe	f 27
10	Mean percentage of recovered ova fertilised per ewe relative to the interval from onset of oestrus to laparotomy	f 27
11	Times of recovery of fertilised ova at various cleavage stages in relation to the onset of oestrus	f 31
12	Incidence of the types of abnormal ova at the commencement of and during culture	f 40
13	Normal 4- cell ovum recovered after 24 hours culture. This ovum was fertilised but uncleaved when inserted into the culture chamber. (Approx. x 300)	f 41
14	Fragmenting 2- cell ovum recovered after 48 hours culture. This ovum was fertilised but uncleaved when inserted into the culture chamber. (Approx. x 400)	f 41

f refers to following page.

Figure

Page

- 15      Fragmenting 4- cell ovum recovered after 48 hours culture.  
This ovum was a normal 4- cell when inserted into the culture chamber. (Approx. x 400)      f 41
- 16      Involuted, unfertilised ovum beginning to fragment after 48 hours culture. This ovum was involuted when inserted into the culture chamber. (Approx. x 400)      f 41
- 17      Two 23 day viable embryos recovered from the recipient ewe pregnant after the transfer of cultured ova. (Approx. x 12.5)      f 48
- 18      Mean crown-rump lengths of embryos from 23 to 101 days of age      f 52