

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

STUDIES ON THE EFFECTS OF GRAZING REGIME

ON SWARD AND DAIRY COW PERFORMANCE

Massey University Library. Thesis Copyright Form

Title of thesis: Studies on the Effects of Grazing Regime
on Sward and Dairy Cow Performance

- (1) (a) I give permission for my thesis to be made available to readers in the Massey University Library under conditions determined by the Librarian.
- (b) ~~I do not wish my thesis to be made available to readers without my written consent for _____ months.~~
- (2) (a) I agree that my ~~thesis, or a copy,~~ may be sent to another institution ~~under conditions determined by the Librarian.~~
- (b) I do not wish my thesis, or a copy, to be sent to another institution without my written consent for 6 months.
- (3) (a) I agree that ~~my thesis may be copied~~ for Library use.
- (b) I do not wish my thesis to be copied for Library use for 6 months.

Signed Clyde G. Hoagwood
Date 4.12.86

The copyright of this thesis belongs to the author. Readers must sign their name in the space below to show that they recognise this. They are asked to add their permanent address.

NAME AND ADDRESS	DATE
_____	_____
_____	_____
_____	_____

ACKNOWLEDGEMENTS

The completion of this study depended on the contributions of many people.

Supervision was provided by C.W.Holmes and A.C.P.Chu. C.W.Holmes provided a remarkably unconstrained atmosphere in which the experiments could be conducted, and contributed ideas and constructive criticism throughout. A.C.P.Chu contributed ideas and encouraged lateral thinking.

Technical assistance was provided by C.A.Butts, K.Garrick, Y.F.Moore, W.Nelson, J.Rumbal and R.Vernal. R.A.McClenaghan and G.C.Tanner provided technical assistance, a practical perspective and essential humour. The figures in this thesis were prepared by F.Dickinson, and O.Harris typed the tables.

Numerous people including fellow post-graduate students and staff members of the Animal Science Department, my family, M.R.Doole and J.S.Tisdall gave encouragement and support.

Financial support was generously provided by the Helen E.Akers, Johannes A.Anderson, G.O.Anstiss, William Hudson and D.J.McGowan scholarships, by the Bank of New Zealand Post-Graduate Bursary in Agriculture and Horticulture, and by the Massey University Agricultural Research Foundation. The Canadian

government provided a student loan.

The assistance of all those who contributed to this study is acknowledged.

TABLE OF CONTENTS

	page
TITLE PAGE	i
ABSTRACT	ii
ACKNOWLEDGEMENTS	iv
LIST OF TABLES	xiii
LIST OF FIGURES	xxi
LIST OF APPENDICES	xxiv
CHAPTER 1 REVIEW OF LITERATURE AND OBJECTIVES	
1.1 INTRODUCTION	1
1.2 SWARD RESPONSE TO DEFOLIATION REGIME	2
1.2.1 THE EFFECTS OF DEFOLIATION ON NET HERBAGE ACCUMULATION	5
1.2.1.1 HERBAGE GROWTH	7
1.2.1.2 HERBAGE DEATH	16
1.2.2 THE INFLUENCE OF DEFOLIATION REGIME ON SWARD STRUCTURE AND COMPOSITION	18
1.3 FACTORS AFFECTING THE INTAKE OF GRAZING ANIMALS	22
1.3.1 HERBAGE MASS	25
1.3.2 HERBAGE HEIGHT	28
1.3.3 BULK DENSITY	29
1.3.4 HERBAGE QUALITY	30
1.3.5 SPATIAL DISTRIBUTION OF SWARD COMPONENTS	32
1.3.6 THE INFLUENCE OF HERBAGE ALLOWANCE ON INTAKE	34

1.4	THE RELATIONSHIP BETWEEN INTAKE AND MILK YIELD IN THE LACTATING COW	37
1.4.1	THE EFFECTS OF LEVEL OF HERBAGE INTAKE ON MILK YIELD AND LIVEWEIGHT	37
1.4.1.1	IMMEDIATE RESPONSE	39
1.4.1.2	SUBSEQUENT EFFECTS	40
1.4.2	THE EFFECTS OF LEVEL OF HERBAGE INTAKE ON MILK COMPOSITION	42
1.4.2.1	IMMEDIATE RESPONSE	42
1.4.2.2	SUBSEQUENT EFFECTS	44
1.4.3	THE EFFECTS OF HERBAGE QUALITY ON MILK YIELD	45
1.4.4	CHANGES IN MILK FAT FATTY ACID COMPOSITION IN RESPONSE TO CHANGES IN ENERGY STATUS OF THE ANIMAL	46
1.5	SCOPE AND OBJECTIVES OF THE PRESENT STUDY	47
CHAPTER 2 GENERAL MATERIALS AND METHODS		
2.1	ENVIRONMENT	49
2.2	WEATHER	50
2.3	SWARD AND ANIMAL MANAGEMENT	50
2.4	SWARD AND ANIMAL MEASUREMENTS	51
2.5	STATISTICAL ANALYSES	51
2.6	DATA PRESENTATION	55
2.7	TERMINOLOGY AND DEFINITIONS	56
2.8	COMMON ABBREVIATIONS	57
2.9	BOTANICAL NAMES OF PASTURE PLANTS	58

CHAPTER 3 THE INFLUENCE OF SEASON AND GRAZING
REGIME ON THE SWARD CHARACTERISTICS
OF THREE SOWN MIXTURES

3.1	INTRODUCTION AND OBJECTIVES	59
3.2	MATERIALS AND METHODS	61
3.2.1	PREPARATION PERIOD	61
3.2.2	EXPERIMENTAL PERIOD	62
3.2.3	STATISTICAL ANALYSIS	65
3.2.4	PRESENTATION OF RESULTS	66
3.3	RESULTS	67
3.3.1	NON-SIGNIFICANT INDEPENDENT VARIABLES	68
3.3.2	HERBAGE COMPOSITION	68
3.3.3	NET HERBAGE ACCUMULATION	71
3.3.4	HERBAGE QUALITY	72
3.3.5	HERBAGE BULK DENSITY	73
3.3.6	HERBAGE MASS COMPONENTS	73
3.4	DISCUSSION	76
3.4.1	SEASON AND GRAZING REGIME	76
3.4.2	MIXTURE SOWN	79
3.4.3	METHODS AND DESIGN USED	81
3.5	CONCLUSIONS	84

CHAPTER 4	THE EFFECT OF DEFOLIATION INTENSITY IN SPRING ON SWARD CHARACTERISTICS IN SPRING AND EARLY SUMMER AND ON DAIRY COW PERFORMANCE WHEN GRAZING THOSE SWARDS IN EARLY SUMMER	
4.1	INTRODUCTION AND OBJECTIVES	115
4.2	GENERAL MATERIALS AND METHODS	117
4.2.1	MATERIALS	117
4.2.1.1	EXPERIMENTAL PADDOCKS	117
4.2.1.2	EXPERIMENTAL ANIMALS	118
4.2.2	METHODS	118
4.2.2.1	EXPERIMENTAL DESIGN	119
4.2.2.2	MANAGEMENT OF EXPERIMENTAL PADDOCKS	119
4.2.2.3	MANAGEMENT OF EXPERIMENTAL ANIMALS	120
4.2.2.4	MEASUREMENTS TAKEN	121
4.2.2.5	STATISTICAL ANALYSES	124
4.3	TRIAL 1 THE EFFECT OF GRAZING INTENSITY IN (1982/83) SPRING ON SWARD CHARACTERISTICS IN EARLY SUMMER AND ON COW PERFORMANCE WHEN GRAZING THOSE SWARDS AT BOTH A GENEROUS AND RESTRICTED DRY MATTER ALLOWANCE IN EARLY SUMMER	125
4.3.1	OBJECTIVES	125
4.3.2	METHODS	125
4.3.3	RESULTS	126
4.3.3.1	EXPERIMENTAL PERIOD	126
4.3.3.2	POST EXPERIMENTAL PERIOD	131

4.4	Trial 2	THE EFFECT OF DEFOLIATION INTENSITY	144
	(1983/84)	IN SPRING ON SWARD CHARACTERISTICS	
		IN EARLY SUMMER, AND ON THE PERFORMANCE	
		OF COWS GRAZING THOSE SWARDS AT A	
		GENEROUS DRY MATTER ALLOWANCE IN EARLY	
		SUMMER	
4.4.1		OBJECTIVES	144
4.4.2		METHODS	144
4.4.3		RESULTS	145
4.4.3.1		EXPERIMENTAL PERIOD	145
4.4.3.2		POST EXPERIMENTAL PERIOD	150
4.5	Trial 3	THE EFFECT OF GRAZING INTENSITY IN	168
	(1984/85)	SPRING ON SWARD CHARACTERISTICS	
		THROUGHOUT SPRING AND EARLY SUMMER,	
		AND ON THE PERFORMANCE OF COWS GRAZING	
		THOSE SWARDS WHEN OFFERED A COMMON NOMINAL	
		GREEN LEAF ALLOWANCE IN EARLY SUMMER	
4.5.1		OBJECTIVES	168
4.5.2		METHODS	169
4.5.3		RESULTS	170
4.5.3.1		SWARD PREPARATION AND EXPERIMENTAL PERIODS	170
4.5.3.2		POST EXPERIMENTAL PERIOD	185
4.6		DISCUSSION AND CONCLUSIONS OF TRIALS 1, 2 AND 3	211
4.6.1		SWARD RESPONSE TO GRAZING REGIME IN SPRING	213

4.6.2	SWARD TYPE, DRY MATTER INTAKE, AND COW BEHAVIOUR	218
4.6.3	HERBAGE ALLOWANCE, DRY MATTER INTAKE AND MILK YIELD	222
4.6.4	IMPLICATIONS OF RESULTS TO DAIRY FARMING	229
4.6.5	CONCLUSIONS	231
CHAPTER 5 THE INFLUENCE OF SWARD REGROWTH INTERVAL IN AUTUMN/WINTER ON SWARD CHARACTERISTICS AND DAIRY COW PERFORMANCE IN EARLY SPRING		
5.1	INTRODUCTION AND OBJECTIVES	238
5.2	MATERIALS AND METHODS	239
5.2.1	PREPARATION PERIOD	241
5.2.2	EXPERIMENTAL PERIOD	242
5.2.3	POST EXPERIMENTAL PERIOD	242
5.2.4	MEASUREMENTS TAKEN DURING THE EXPERIMENTAL PERIOD	242
5.2.5	STATISTICAL ANALYSES	245
5.3	RESULTS	245
5.3.1	SWARD PREPARATION PERIOD	245
5.3.2	EXPERIMENTAL PERIOD	246
5.3.2.1	PRE-GRAZING SWARD CHARACTERISTICS	246
5.3.2.2	HERBAGE ALLOWANCE AND INTAKE	249
5.3.2.3	COW MEASUREMENTS	252
5.3.3	POST EXPERIMENTAL PERIOD	254
5.4	DISCUSSION	254

5.4.1	NET HERBAGE ACCUMULATION OVER THE AUTUMN/WINTER PERIOD	255
5.4.2	SWARD STRUCTURE AND COMPOSITION	256
5.4.3	ALLOWANCE AND INTAKE	258
5.4.4	MILK PRODUCTION	260
5.5	CONCLUSIONS	264
CHAPTER 6 GENERAL DISCUSSION AND CONCLUSIONS		
6.1	RESUME OF RESULTS AND CONCLUSIONS	279
6.2	COMPARISON OF RESULTS WITH RELEVANT PUBLISHED WORK	282
6.3	SOME PROBLEMS ASSOCIATED WITH MEASUREMENT TECHNIQUES USED	284
6.4	FUTURE WORK	288
6.5	RELEVANCE OF RESULTS TO DAIRY FARMING	290
APPENDICES		294
BIBLIOGRAPHY		338

LIST OF TABLES

3.1	Description and sowing rates of the three seed mixtures.	86
3.2	Target and actual residual herbage masses (RHM) (Kg DM/ha) to achieve the lax (L) and hard (H) grazing intensities at each season (the actual RHM's are raw data means and their standard deviations).	88
3.3	Grazing dates for the frequently (F) and infrequently (I) grazed swards.	88
3.4	The effect of sward type on pre-grazing herbage composition (% dry weight) averaged across all seasons and grazing regimes.	89
3.5	The effect of grazing regime on pre-grazing herbage composition (% dry weight) averaged across all sward types and seasons.	89
3.6	The effect of season on pre-grazing herbage composition (% dry weight) averaged across all sward types and grazing regimes.	96
3.7	The effect of sward type on pre-grazing herbage dry matter digestibility (%), nitrogen concentration (%) and bulk density (Kg DM/ha per cm sward surface height), and on net herbage accumulation (Kg DM/ha daily) averaged across all grazing regimes and seasons.	96
3.8	The effect of grazing regime on pre-grazing herbage dry matter digestibility (%), nitrogen concentration (%), and bulk density (Kg DM/ha per cm sward surface height) and on net herbage accumulation (Kg DM/ha daily) averaged across all sward types and seasons.	97
3.9	The effect of season on pre-grazing herbage dry matter digestibility (%), nitrogen concentration (%), and bulk density (Kg DM/ha per cm sward surface height), and on net herbage accumulation (Kg DM/ha daily) averaged across all sward types and grazing regimes.	97
3.10	The effect of sward type on pre-grazing herbage mass components (Kg DM/ha) averaged across all grazing regimes and seasons.	105
3.11	The effect of grazing regime on pre-grazing herbage mass components (Kg DM/ha) averaged across all sward types and seasons.	105
3.12	The effect of season on pre-grazing herbage mass components (Kg DM/ha) averaged across all sward types and grazing regimes.	114

4.3.1	The effect of grazing intensity in spring on the total pre-grazing herbage mass (Kg DM/ha) and composition (% dry weight), and on sward surface height (cm) and herbage bulk density (Kg DM/ha per cm of sward surface height) in the experimental period (early summer).	133
4.3.2	The effect of grazing intensity in spring on the pre-grazing mass of total green and leaf matter (Kg DM/ha) and digestible dry matter (DDM) (Kg DDM/ha) in the experimental period (early summer).	133
4.3.3	The effect of grazing intensity in spring on the (a) <u>in vitro</u> dry matter digestibility (DMD) (%) of the pre-grazing total herbage, grass leaf, and grass stem and (b) <u>in vivo</u> dry and organic matter digestibility (DMD and OMD respectively) (%) of the pre-grazing total herbage in the experimental period (early summer).	135
4.3.4	The effect of sward type and daily dry matter allowance (DA) on the daily allowance of green and leaf matter (GA and LA respectively) (Kg DM/cow daily), on dry matter intake (DMI) (Kg DM/cow daily), and on the composition of the diet (% dry weight) in the experimental period (early summer).	136
4.3.5	The effect of sward type and daily dry matter allowance (DA) on the residual herbage mass (Kg DM/ha), residual herbage height (cm) and <u>in vitro</u> dry matter digestibility of the residual herbage (%) in the experimental period (early summer).	137
4.3.6	The effect of sward type and daily dry matter allowance (DA) on milk, milk fat, and milk protein yield (Kg/cow daily) in week 2 of the experimental period (early summer).	137
4.3.7	The effect of sward type and daily dry matter allowance (DA) on the composition of milk fat (% weight of short and long chain fatty acids) at the end of week 2 of the experimental period (early summer).	141
4.3.8	The effect of sward type and daily dry matter allowance (DA) on unfasted liveweight (Kg/cow) and condition score at the end of the experimental period and on changes in unfasted liveweight (Kg/cow over 16 days) over the experimental period (early summer).	141
4.3.9	The effect of sward type and daily dry matter allowance (DA) in the experimental period (early summer) on milk, milk fat, and milk protein yield in the first week of the post experimental period (early summer) (Kg/cow daily).	142

- 4.3.10 The effect of grazing intensity in spring and of daily dry matter allowance (DA) in the experimental period on the sward characteristics of pre-grazing herbage in the post experimental period (February/March). 143
- 4.4.1 The effect of defoliation intensity in spring on the total pre-grazing herbage mass (Kg DM/ha) and composition (% dry weight), and on sward surface height (cm), herbage bulk density (Kg DM/ha per cm sward surface height) and tiller density (no. tillers/m) in the experimental period - Round 1 (early summer). 151
- 4.4.2 The effect of defoliation intensity in spring on the pre-grazing mass of total green and leaf matter (Kg DM/ha) and digestible dry matter (DDM) (Kg DDM/ha) in the experimental period - round 1 (early summer). 152
- 4.4.3 The effect of defoliation intensity in spring on the (a) in vitro dry and organic matter digestibility (DMD and OMD respectively) (%) of the pre-grazing total herbage, grass stem, grass leaf, clover, and senescent matter, and (b) in vivo dry and organic matter digestibility (%) of the total pre-grazing herbage in the experimental period - round 1 (early summer). 154
- 4.4.4 The effect of defoliation intensity in spring on (a) the nitrogen concentration (N%) and (b) the in vitro dry and (c) organic matter digestibility (%) of the pre-grazing total herbage in each of five vertical strata, in the experimental period - round 1 (early summer). 155
- 4.4.5 The effect of sward type and daily dry matter allowance (DA) on the residual herbage mass (Kg DM/ha), residual herbage height (cm) and in vitro dry and organic matter digestibility (%) of the residual herbage in the experimental period - round 1 (early summer). 156
- 4.4.6 The effect of defoliation intensity in spring on the total pre-grazing herbage mass (Kg DM/ha) and composition (% dry weight), and on sward surface height (cm), herbage bulk density (Kg DM/ha per cm of sward surface height) and tiller density (no. tillers/m²) in the experimental period - round 2 (early summer). 157
- 4.4.7 The effect of defoliation intensity in spring on (a) net herbage accumulation (Kg DM/ha daily) between rounds 1 and 2 in the experimental period in early summer, and (b) the pre-grazing mass of total green and leaf matter (Kg DM/ha) and digestible dry matter (DDM) (Kg DDM/ha) in the experimental period - round 2 (early summer). 158

- 4.4.8 The effect of defoliation intensity in spring on the in vitro dry and organic matter digestibility (DMD and OMD respectively) (%) of pre-grazing total herbage, grass leaf, clover, and senescent matter in the experimental period - round 2 (early summer). 158
- 4.4.9 The effect of defoliation intensity in spring on the (a) nitrogen concentration (%) and the in vitro (b) dry and (c) organic matter digestibility (%) of the pre-grazing total herbage in each of five vertical strata in the experimental period - round 2 (early summer). 159
- 4.4.10 The effect of sward type and daily dry matter allowance (DA) on the daily allowance of green (GA) and leaf (LA) matter (Kg DM/cow daily), dry matter intake (DMI) (Kg DM/cow daily), and on the composition of the diet (% dry weight) in the experimental period - round 1 (early summer). 160
- 4.4.11 The effect of sward type and daily dry matter allowance (DA) on the daily allowance of green (GA) and leaf (LA) matter (Kg DM/cow daily) and on dry matter intake (DMI) (Kg DM/cow daily) in the experimental period - round 2 (early summer). 161
- 4.4.12 The effect of sward type and daily dry matter allowance (DA) on the residual herbage mass (Kg DM/ha), residual herbage height (cm) and in vitro dry and organic matter digestibility (%) of the residual herbage in the experimental period - round 2 (early summer). 161
- 4.4.13 The effect of sward type on milk, milk fat and milk protein yield (Kg/cow daily) in (a) week 3 of round 1 and (b) week 2 of round 2 of the experimental period (early summer). 162
- 4.4.14 The effect of sward type on the composition of milk fat (% weight of short and long chain fatty acids) at (a) the end of week 3 of round 1 and (b) the end of week 2 of the experimental period (early summer). 162
- 4.4.15 The effect of sward type on unfasted liveweight (Kg/cow) and condition score at the end of the experimental period, and on changes in unfasted liveweight (Kg/cow over 35 days) and condition score over the experimental period (early summer). 166
- 4.4.16 The effect of sward type in the experimental period on milk, milk fat, and milk protein yield (Kg/cow daily) in the first week of the post experimental period (early summer). 166

4.4.17 The effect of defoliation intensity in spring on pre-grazing sward characteristics in the post experimental period (late summer).	167
4.5.1 Pre-grazing sward characteristics at the commencement of the first round of differential grazing in spring (October).	188
4.5.2 Post-grazing sward characteristics of swards that were grazed laxly (L), moderately (M), or intensely (I) during the sward preparation (Oct-Nov) and experimental (Nov-Dec) periods.	195
4.5.3 The effect of grazing intensity in spring on (a) net herbage accumulation (Kg DM/ha daily), (b) sward surface height (cm) and (c) herbage bulk density (Kg DM/ha per cm sward surface height) during the sward preparation (Oct-Nov), experimental (Nov-Dec) and post experimental (Feb-March) periods.	196
4.5.4 The effect of grazing intensity in spring on the pre-grazing masses of total (a) dry, (b) green (c) leaf and (d) digestible dry matter during the sward preparation (Oct-Nov), experimental (Nov-Dec) and post experimental (Feb-March) periods (Kg DM/ha).	197
4.5.5 The effect of grazing regime in spring on the weights (mg dry weight) of individual (a) reproductive and (b) vegetative tillers and on the (c) percentage of reproductive tillers in the sward (% by no. of total tillers) before grazing during the sward preparation (Oct-Nov) and experimental periods (Nov-Dec).	198
4.5.6 The effect of grazing intensity in spring on the number of visible inflorescences (no. per m ²) during the sward preparation (Oct-Nov) and experimental (Nov-Dec) periods.	198
4.5.7 The effect of grazing intensity in spring on the <u>in vitro</u> dry (DMD) and organic (OMD) matter digestibility (%) of the pre-grazing total herbage during the sward preparation (Oct-Nov) and experimental (Nov-Dec) periods.	199
4.5.8 The effect of grazing intensity in spring on the <u>in vitro</u> dry (DMD) and organic (OMD) matter digestibility (%) of the total herbage within each of five vertical strata in the pre-grazing herbage during (a) round 1 and (b) round 2 of the experimental period (Nov-Dec).	200
4.5.9 The effect of grazing intensity in spring on the <u>in vitro</u> dry (DMD) and organic (OMD) matter digestibility (%) of (a) grass stem, (b) grass leaf and (c) senescent matter in the pre-grazing herbage during the experimental period (Nov-Dec).	201

4.5.10	The effect of grazing intensity in spring and daily leaf allowance (LA) (Kg DM/cow daily) on the allowance of green (GA) and total dry (DA) matter) (Kg DM/cow daily) and on dry matter intake (DMI) (Kg DM/cow daily) during the experimental period (Nov-Dec) (a) round 1 and (b) round 2.	202
4.5.11	The effect of sward type on the composition of the diet (% dry weight) during the experimental period (Nov-Dec) (a) round 1 and (b) round 2.	203
4.5.12	The effect of sward type on the contribution (% dry weight) of each of five vertical strata to the diet during the experimental period (Nov-Dec) in (a) round 1 and (b) round 2.	204
4.5.13	The effect of sward type on the behaviour of cows in the experimental period (Nov-Dec) - round 1.	204
4.5.14	The effect of observation day on the behaviour of cows in the experimental period (Nov-Dec) - round 1.	205
4.5.15	The effect of sward type on milk, milk fat, and milk protein yield (Kg/cow daily) in (a) week 3 of round 1 and (b) week 2 of round 2, of the experimental period (Nov-Dec).	205
4.5.16	The effect of sward type on the composition of milk fat (% weight of short and long chain fatty acids) at (a) the end of week 3 of round 1 and (b) the end of week 2 of round 2 of the experimental period (Nov-Dec).	209
4.5.17	The effect of sward type on unfasted liveweight (Kg/cow) at the end of the experimental period, and on changes in unfasted liveweight (Kg/cow over 35 days) over the experimental period (Nov-Dec).	210
4.5.18	The effect of sward type in the experimental period on milk, milk fat, and milk protein yield (Kg/cow daily) in the first week of the post experimental period (Dec).	210
4.6.1	A comparison of ME intake calculated using diet composition, component organic matter digestibility, and dry matter intakes obtained in T1, T2 and T3, with that calculated using milk yield and unfasted liveweight data.	235
4.6.2	Results of stepwise regression of sward characteristics and herbage allowance on dry matter intake (Kg DM/cow daily) and milk yield (Kg/cow daily).	236
4.6.3	Correlation matrix for pre- and post-grazing (BG and AG respectively) total herbage mass, height, and dry matter digestibility (DMD), dry green, and leaf matter	237

- allowance (DA, GA, and LA respectively), dry matter intake (DMI) and milk, milk fat, and milk protein yield.
- 5.1 The effect of regrowth interval in autumn/winter on net herbage accumulation over May to September (Kg DM/ha daily). 266
- 5.2 The effect of regrowth interval in autumn/winter on pre-grazing herbage mass (Kg DM/ha), sward surface height (cm) and herbage bulk density (Kg DM/ha per cm sward surface height) in early spring. 266
- 5.3 The effect of regrowth interval in autumn/winter on the composition (% dry weight), in vitro (DMD) and organic (OMD) matter digestibility (%), and nitrogen concentration (N%) in early spring. 267
- 5.4 The effect of regrowth interval in autumn/winter on the pre-grazing mass of herbage components in early spring (Kg DM/ha). 268
- 5.5 The effect of regrowth interval in autumn/winter on pre-grazing herbage in vivo dry (DMD) and organic (OMD) matter digestibility (%) in early spring (all numbers are raw data means and their SD's). 268
- 5.6 The effect of regrowth interval in autumn/winter on the percentage contribution of each vertical stratum to the total pre-grazing herbage mass in early spring (% dry weight). 270
- 5.7 The effect of regrowth interval in autumn/winter on the in vitro dry (DMD) and organic (OMD) matter digestibility of the total herbage within each vertical stratum of pre-grazing herbage present in early spring (%). 271
- 5.8 The effect of sward type on the allowance (Kg DM/cow daily) of total green and leaf matter when dry matter allowance was kept constant; on the dry matter intake (Kg DM/cow daily) of cows grazing those swards; and on the residual herbage mass (Kg DM/ha). 272
- 5.9 The effect of sward type on the percentage contribution of each vertical stratum to the total diet of cows grazing those swards in early spring (%). 272
- 5.10 The effect of sward type on the composition of the diet of cows grazing those swards in early spring (% dry weight). 273
- 5.11 The effect of sward type on milk, milk fat, and milk protein yield (Kg/cow daily), and on the fatty acid composition of milk fat (% by weight) in week 3 of the 273

- experimental period, and on unfasted liveweight and changes in unfasted liveweight over the experimental period (Kg).
- 5.12 The effect of sward type on the amount of time cows spent at each activity during (a) daylight and (b) darkness (in hours). 277
- 5.13 The effect of observation day on the amount of time cows spent at each activity during (a) daylight and (b) darkness (in hours). 278
- 5.14 The effect of sward type in the experimental period on milk and milk fat yield in week 1 of the post experimental period (Kg/cow daily). 278
- 6.1 A comparison of dry matter intakes (Kg DM/cow daily) measured (by difference) in chapter 4 and 5 and predicted using the equation of MAFF (1975). 291
- 6.2 A comparison of organic matter intakes (Kg OM/cow daily) measured (by difference) in chapters 4 and 5 and predicted using the equation of Caird and Holmes (1986). 292
- 6.3 A comparison of dry matter intakes (Kg DM/cow daily) measured by difference) in chapters 4 and 5 and predicted using the equation of Stockdale (1985). 293

LIST OF FIGURES

1.1	Relationships between herbage mass (Kg DM/ha), sward height (cm) and LAI, and rates of herbage growth (G), senescence (S) and net production (NP) (all Kg DM/ha daily) in swards continuously grazed by sheep (Bircham and Hodgson, 1983).	8
1.2	Conceptual illustration of the plant-animal interface in grazed pastures: I. Illustrating the combined effects of herbage quality and character on the behavioural, physical and metabolic limits to intake. The effects of current intake (quantity and composition) on subsequent herbage quality and character are also illustrated.	24
3.1	Diagram of experimental area.	87
3.2	The effect of season and sward type on the percentage of clover in the pre-grazing herbage, averaged across all grazing regimes.	90
3.3	The effect of season and grazing regime on pre-grazing herbage composition, averaged across all sward types.	92
3.4	The effect of season and grazing regime on the percentage of prairie grass in Matua swards.	94
3.5	The effect of season and grazing regime on net herbage accumulation, averaged across all sward types.	98
3.6	The effect of season and grazing regime on the dry matter digestibility of the pre-grazing herbage, averaged across all sward types.	99
3.7	The effect of season and grazing regime on the nitrogen concentration of the pre-grazing herbage, averaged across all sward types.	101
3.8	The effect of season and grazing regime on the bulk density of the pre-grazing herbage, averaged across all sward types.	103
3.9	The effect of season and sward type on the pre-grazing masses of total dry, green and senescent matter, averaged across all grazing regimes.	106
3.10	The effect of season and sward type on the pre-grazing mass of total digestible dry matter, averaged across all grazing regimes.	108

3.11 The effect of season and grazing regime on the pre-grazing masses of total dry, green and senescent matter, averaged across all sward types.	110
3.12 The effect of season and grazing regime on the pre-grazing mass of total digestible dry matter, averaged across all sward types.	112
4.3.1 The effect of grazing intensity in spring on the pre-grazing mass of weed, clover, grass leaf, grass stem and senescent matter in early and late summer.	134
4.3.2 The effect of sward type and daily dry matter allowance on milk yield in the experimental and post experimental periods.	138
4.3.3 The effect of sward type and daily dry matter allowance on milk fat yield in the experimental and post experimental periods.	139
4.3.4 The effect of sward type and daily dry matter allowance on milk protein yield in the experimental and post experimental periods.	140
4.4.1 The effect of defoliation intensity in spring on the pre-grazing mass of weed, clover, grass leaf, grass stem and senescent matter in rounds 1 and 2 of the experimental period (early summer) and in the post experimental period (late summer).	153
4.4.2 The effect of sward type on milk yield in the experimental and post experimental periods.	163
4.4.3 The effect of sward type on milk fat yield in the experimental and post experimental periods.	164
4.4.4 The effect of sward type on milk protein yield in the experimental and post experimental periods.	165
4.5.1 The effect of grazing intensity in spring on the percentage of grass stem, grass leaf, clover stem, clover leaf and senescent matter in the total pre-grazing herbage in spring and summer.	189-190
4.5.2 The effect of grazing intensity in spring on the percentage of grass stem, grass leaf, clover stem, clover leaf and senescent matter in the total pre-grazing herbage in each of five vertical strata, and of the percentage contribution of total herbage in each stratum to that of the total herbage mass over all strata.	191-194

4.5.3	The effect of sward type on milk yield in the experimental and post experimental periods.	206
4.5.4	The effect of sward type on milk fat yield in the experimental and post experimental periods.	207
4.5.5	The effect of sward type on milk protein yield in the experimental and post experimental periods.	208
5.1	Diagram of experimental area.	265
5.2	The effect of regrowth interval in autumn/winter on the pre-grazing mass of clover, grass leaf, grass stem and senescent matter in each of five vertical strata in the sward in early spring.	269
5.3	The effect of sward type on milk yield in the experimental and post experimental periods.	274
5.4	The effect of sward type on milk fat yield in the experimental and post experimental period.	275
5.5	The effect of sward type on milk protein yield in the experimental period.	276

LIST OF APPENDICES

2.1	Meteorological data for each month from April 1982 to March 1985.	294
2.2	Measurement techniques.	295
3.1	Statistical analysis - split plot in space and time.	311
3.2	Statistical analysis - percentage and mass of prairie grass present in M swards.	314
3.3	Analysis of variance of pre-grazing herbage composition (% dry weight). All values are F-values.	316
3.4	Analysis of variance of the pre-grazing percentage and total mass of prairie grass in the M swards only. All values are F-values.	316
3.5	Analysis of variance of pre-grazing herbage bulk density and of net herbage accumulation. All values are F-values.	317
3.6	Analysis of variance of pre-grazing masses of total digestible dry matter and nitrogen, and pre-grazing herbage dry matter digestibility and nitrogen concentration. All values are F-values.	318
3.7	Analysis of variance of pre-grazing masses of total dry, green, and senescent matter. All values are F-values.	319
4.2	An illustration of the method of division of each experimental paddock in (a) Trial 1 (1982/83), (b) Trial 2 (1983/84), and (c) Trial 3 (1984/85).	320
4.3.1	Details of cow age (yrs) and milk fat yield (Kg/cow daily) and unfasted liveweight (kg) in the preparation period (trial 1).	321
4.3.2	A t-test of the hypothesis that the mean percentage of a component in the diet comes from the same population as the mean percentage of a component in the herbage on offer (SoKal and Rohlf, 1981).	322
4.4.1	Details of cow age (yrs) and milk fat yield (Kg/cow daily) and unfasted liveweight (Kg/cow) in the preparation period (trial 2).	323

- 4.4.2 A t-test of the hypothesis that the mean percentage of a component in the diet comes from the same population as the mean percentage of a component in the herbage on offer (SoKal and Rohlf, 1981). 323
- 4.5.1 Details of cow age (yrs) and milk fat yield (Kg/cow daily) and unfasted liveweight (kg/cow) in the preparation period (trials 3). 324
- 4.5.2 The effect of grazing intensity in spring on the percentage contribution of each of five vertical strata to the pre-grazing total herbage mass (% dry weight) in the experimental period (Nov-Dec). 325
- 4.5.3 The effect of grazing intensity on the composition (% dry weight) of the pre-grazing total herbage mass in the sward preparation (Oct-Nov), experimental (Nov-Dec), and post experimental (Feb-March) periods. 326
- 4.5.4 A t-test of the hypothesis that the mean proportion of a component in the diet comes from the same population as the mean proportion of that component in the herbage on offer (Sokal and Rohlf, 1981). 327
- 4.5.5 A t-test of the hypothesis that the mean percentage contribution of a stratum to the diet comes from the same population as the mean percentage contribution of that stratum to the total herbage on offer (SoKal and Rohlf, 1981). 328
- 4.5.6 Meteorological data for the days when cow behaviour was monitored. 329
- 4.6.1 Simple regression of the percentage of a component in the diet on the percentage of that component in the herbage on offer in T1, T2, and T3. 330
- 4.6.2 Simple regression of dry matter intake (DMI) (Kg/cow daily) on dry, green, and leaf allowance (DA, GA and LA respectively) (Kg DM/cow daily) in T1, T2, and T3; after adjustment for year, cow type and grazing round. 330
- 5.1 Details of cow age (years) and milk fat yield (Kg/cow daily) and unfasted liveweight (kg) in the preparation period. 331
- 5.2 Details of sward characteristics and management during the sward preparation period (May - September 1984). 332
- 5.3 The effect of regrowth interval in autumn/winter on the pre-grazing mass of herbage components in each vertical stratum in early spring (Kg DM/ha). 333

- 5.4 A t-test of the hypothesis that the mean of the percentage contribution of each stratum to the diet comes from the same population as the percentage contribution of each stratum to the total pre-grazing herbage mass (Sokal and Rohlf, 1981). 334
- 5.5 A t-test of the hypothesis that the mean of the percentage of a component in the diet comes from the same population as the percentage of a component in the pre-grazing herbage mass (Sokal and Rohlf, 1981). 335
- 5.6 Meteorological data for the days when cow behaviour was monitored. 336
- 5.7 Sample calculation of the ME concentration of herbage ingested and of total ME ingested (per cow daily). 336
- 5.8 Sample calculation of ME intake needed to sustain production levels measured in the experimental period (per cow daily). 337