

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**STUDIES ON PLANTS THAT  
HYPERACCUMULATE COPPER, COBALT  
AND NICKEL: THEIR POTENTIAL FOR  
USE IN PHYTOMINING AND  
PHYTOREMEDIATION.**

A thesis presented in partial fulfillment of the requirements for the degree of Masterate of Science in Soil Science at Massey University.

**Frances Anneli Bennett  
1998**

## ABSTRACT

This thesis reviews three lines of research on hyperaccumulators to examine their potential for phytoremediation and phytomining. The first line of research was to test the affect of nutrient addition on biomass and nickel uptake by two nickel hyperaccumulators, *Alyssum bertolonii* and *Streptanthus polygaloides*. Addition of fertiliser increased the biomass although the maximum amount added was found to be suboptimal. Nutrient addition did not affect the rate of nickel uptake. Larger plants contained a more dilute nickel content but still had an overall larger amount than smaller plants.

The second line of research was to test the affect of chelates on metal uptake by copper and cobalt flora of the Democratic Republic of Congo (formally Zaire) and a copper tolerant plant from Spain, *Erica andevalensis*. EDTA and Citric Acid increased uptake of copper in these plants but had no effect on the uptake of cobalt and nickel. EDTA increased the uptake of lead by *Alyssum bertolonii* but did not affect the uptake of zinc and cadmium.

The third line of research was to examine the reality of hyperaccumulators of copper and cobalt. Copper and cobalt hyperaccumulation does in fact exist but not to the extent reported previously. There is a good possibility that the previously reported values for copper and cobalt hyperaccumulation are in some cases erroneous due to high iron levels indicating contamination of plant samples by soil.

## ACKNOWLEDGEMENTS

I would like to thank several people who aided me in the production of my thesis. I would like to thank Mr Brynn Neilson who used his computer wizardry to scan in, resize and adjust all the photographs embedded in the text. I would like to thank my father, Mr Arthur Bennett, who initially proof read my text and made constructive suggestions on format and style. I would like to thank most sincerely Doctor Robert Stewart of Massey University who proof read my text for content and made many helpful comments. Finally I would like to express my deep indebtedness and gratitude to Emeritus Professor Robert R. Brooks of Massey University. Professor Brooks, known affectionately to his students as Brooksy, has guided me with his wisdom, experience and profound knowledge, has tolerated my weaknesses and has encouraged and motivated me to finish a task which I at times found daunting. Any student thriving in the fertile soil of his instruction must acquire mettle.

# TABLE OF CONTENTS

Abstract	ii
Acknowledgements	iii
Table of Contents	iv
List of Figures	v
List of Tables	vii
<b>1 Introduction</b>	<b>1</b>
<b>2 Literature review</b>	<b>7</b>
<b>3 The geological and Climatological Environment of the Hyperaccumulators Studied</b>	<b>11</b>
The Shaban Copper Arc	11
The Ultramafic Outcrops of Southern Europe	15
Ultramafic Outcrops of the Sierra Nevada Range, California	16
The Pyritic Orebodies of Huelva, Southwest Spain	17
<b>4 Nickel Hyperaccumulators</b>	<b>19</b>
Introduction	19
Purpose and general hypotheses of this research	20
<i>Alyssum bertolonii</i>	22
<i>Streptanthus polygaloides</i>	23
Pot Trials	24
Analytical methods	26
Results	27
General discussion	35
<b>5 Copper and Cobalt Hyperaccumulators</b>	<b>37</b>
Introduction	37
<i>Haumaniastrum robertii</i>	39
<i>Haumaniastrum katangense</i>	41
<i>Aeollanthus biformifolius</i>	43
<i>Erica andevalensis</i>	44
Pot Trials	47
Analytical methods	47
Results	51
General discussion	61
<b>6 Reality of Hyperaccumulators</b>	<b>64</b>
<b>7 Discussion</b>	<b>69</b>
<b>Bibliography</b>	<b>71</b>

## LIST OF FIGURES

Figure 1:	Histograms showing the abundance of nickel in <i>Alyssum</i> species.	2
Figure 2:	Abandoned native copper smelter fashioned from a termite hill at Luisha colonised by <i>Haumaniastrum katangense</i> (not in flower) and the fern <i>Nephrolepis undulata</i> .	5
Figure 3:	The Shaban Copper Arc and Zambian Copperbelt.	12
Figure 4:	Structural setting of the Shaban Copper Arc.	12
Figure 5:	Distribution of Ophiolites in the Apennines.	15
Figure 6:	Ultramafic outcrops in California and Oregon.	17
Figure 7:	<i>Alyssum bertolonii</i> .	19
Figure 8:	<i>Streptanthus polygaloides</i> .	23
Figure 9:	Graph showing mean biomass vs. fertiliser treatment for <i>Streptanthus polygaloides</i> .	28
Figure 10:	Graph showing mean biomass vs. fertiliser treatment for <i>Alyssum bertolonii</i> .	29
Figure 11:	Graph showing mean nickel concentration vs. fertiliser treatment for <i>Alyssum bertolonii</i> .	30
Figure 12:	Effect of fertiliser treatment on the biomass of <i>Alyssum bertolonii</i> .	32
Figure 13:	Graph showing mean nickel concentration vs. biomass for <i>Alyssum bertolonii</i> .	33
Figure 14:	Graph showing the effect of EDTA on the uptake of cadmium, lead and zinc by <i>Alyssum bertolonii</i> .	34
Figure 15:	The cobalt hyperaccumulator and indicator plant <i>Haumaniastrum robertii</i> at Mindingi, Democratic Republic of Congo (Zaire).	40
Figure 16:	Distribution of some Shaban plants endemic to copper/cobalt deposits.	41
Figure 17:	Carpet of <i>Haumaniastrum katangense</i> growing over copper poisoned soil derived from the stack of the giant smelter at Lubumbashi, shown in the background.	42

- Figure 18:** The copper hyperaccumulator and indicator plant *Aeollanthus biformifolius* at Etoile mine, Democratic Republic of Congo (Zaire). 43
- Figure 19:** Distribution of some Shaban plants endemic to copper/cobalt deposits in the Lubumbashi area. 43
- Figure 20:** *Erica andevalensis*. 44
- Figure 21:** *Erica andevalensis* in flower during pot trials at Massey University. 45
- Figure 22:** Map of the Huelva region of Southwest Spain showing the locality of the occurrence of *Erica andevalensis*. 46
- Figure 23:** Graph showing effect of EDTA on the uptake of copper by *Haumaniastrum katangense* and *Haumaniastrum robertii*. 53
- Figure 24:** Graph showing effect of EDTA on the uptake of cobalt by *Haumaniastrum katangense* and *Haumaniastrum robertii*. 53
- Figure 25:** Graph showing effect of EDTA on the uptake of nickel by *Haumaniastrum katangense* and *Haumaniastrum robertii*. 54
- Figure 26:** Graph showing effect of EDTA and Citric Acid on the uptake of copper by *Haumaniastrum katangense*. 55
- Figure 27:** Graph showing effect of EDTA and citric Acid on the uptake of copper by *Haumaniastrum robertii*. 56
- Figure 28:** Graph showing mean copper, cobalt and iron values for *Haumaniastrum katangense* grown in seven Champion peat/pumice treatments and control. 56
- Figure 29:** Graph showing mean copper uptake by *Aeollanthus biformifolius* after the addition of EDTA and Citric Acid. 58
- Figure 30:** Graph showing copper in *Erica andevalensis* grown in Champion Ore/seed mix. 60

## LIST OF TABLES

<b>Table 1:</b>	Climatological data for stations in Democratic Republic of Congo (Zaire), Zambia and Zimbabwe.	14
<b>Table 2:</b>	Sixteen different fertiliser treatments used in pot trials with <i>Alyssum bertolonii</i> and <i>Streptanthus polygaloides</i> .	25
<b>Table 3:</b>	Mean biomass and Nickel concentration of <i>Streptanthus polygaloides</i> in relation to the different fertiliser treatments used.	28
<b>Table 4:</b>	Mean biomass and Nickel concentration of <i>Alyssum bertolonii</i> in relation to the different fertiliser treatments used.	30
<b>Table 5:</b>	Mean biomass and Nickel concentration of <i>Alyssum bertolonii</i> in relation to the different nitrogen fertiliser treatments used.	31
<b>Table 6:</b>	Medians and geometric means for biomass and Nickel concentration of <i>Alyssum bertolonii</i> in relation to the different nitrogen fertiliser treatments used.	32
<b>Table 7:</b>	Statistical treatment of biomass and Nickel concentration data for <i>Alyssum bertolonii</i> .	32
<b>Table 8:</b>	Statistics for biomass, nickel concentration and total nickel content for <i>Alyssum bertolonii</i> .	34
<b>Table 9:</b>	Analyses for cadmium, lead and zinc uptake by <i>Alyssum bertolonii</i> grown in Tui tailings/bark.	34
<b>Table 10:</b>	Normal and abnormal concentrations of elements in plant leaves ( $\mu\text{g/g}$ ).	38
<b>Table 11:</b>	Numbers of known hyperaccumulators for eight heavy metals and the families in which they are most often found.	39
<b>Table 12:</b>	Rates of EDTA and Citric Acid addition to test substrates.	48
<b>Table 13:</b>	Uptake of copper and nickel by <i>Haumaniastrum katangense</i> and <i>Haumaniastrum robertii</i> growing in 500 $\mu\text{g/g}$ copper and United Ore.	53
<b>Table 14:</b>	Uptake of copper, cobalt and nickel by <i>Haumaniastrum katangense</i> growing in United Ore Substrate.	55
<b>Table 15:</b>	Uptake of copper, cobalt and nickel by <i>Haumaniastrum robertii</i> growing in United Ore Substrate.	56



<b>Table 16:</b>	Copper, cobalt and iron uptake by <i>Haumaniastrum katangense</i> grown in Champion Ore.	57
<b>Table 17:</b>	Mean copper and cobalt uptake by <i>Aeollanthus biformifolius</i> grown in United Ore after the addition of EDTA and Citric Acid.	58
<b>Table 18:</b>	Copper and iron concentrations in mineralised soil from the Andévalo, Huelva province.	59
<b>Table 19:</b>	Copper and iron concentrations in specimens of Andévalo <i>Erica andevalensis</i> .	60
<b>Table 20:</b>	Copper and iron concentrations in plants of <i>Erica andevalensis</i> grown in varying admixtures of Champion ore and seed mix.	61
<b>Table 21:</b>	Copper and iron concentrations in <i>Erica andevalensis</i> grown in admixtures of Champion ore and peat/pumice.	62
<b>Table 22:</b>	Iron, apparent copper and cobalt and corrected copper and cobalt for selected plant species.	66
<b>Table 23:</b>	Analyses of herbarium specimens collected by R.R. Brooks in Shaba, Democratic Republic of Congo (Zaire).	67
<b>Table 24:</b>	Analyses of herbarium specimens collected by R.R. Brooks in Shaba, Democratic Republic of Congo (Zaire).	68

# INTRODUCTION

## HYPERACCUMULATORS

Natural mineral deposits containing particularly large quantities of heavy metals are present in many regions of the globe. These areas often support very characteristic plant assemblages and species that thrive in these metal-enriched environments. Examples of such distinct plant communities include the serpentine flora (growing on Ni, Cr, Mn, Mg and Co rich soils), the seleniferous flora (growing on Se rich soil), the uraniferous flora (growing on U rich soils), the calamine flora (growing on Zn and Cd rich soils), and Copper/Cobalt floras. As a result of their association with specific ore deposits, many metallophyte plants are used as so-called indicator species in prospecting for mineral deposits. For example, the copper flower (*Haumaniastrum katangense*) is a classical indicator of Cu and Co ores in central Africa.

There are three basic categories of plant that grow on metalliferous soils. Metal excluders effectively prevent metal from entering their aerial parts when growing in a broad range of metal concentrations in the soil. Metal non-excluders actively accumulate metals in their above-ground tissues and can be roughly divided into two groups: indicators and hyperaccumulators. Metal levels in the tissues of indicator species generally reflect metal levels in the soil. Hyperaccumulators can concentrate metals in their above-ground tissues to levels far exceeding those present in the soil.


Baumann (1885) recorded the first hyperaccumulator of any metal when he reported over 1% zinc in the dried leaves of *Thlaspi calaminare* and *Viola calaminaria* growing over zinc deposits near Aachen, Germany. Minguzzi and Vergnano (1948) were the first to discover high nickel levels in plants. They reported over 1% nickel in leaves of *Alyssum bertolonii* growing over ultramafic rocks in Tuscany.

By far the greatest volume of work and general interest in hyperaccumulators has been centered on the genus *Alyssum*. This followed the original discovery of over 1% Ni in *Alyssum bertolonii* by Minguzzi and Vergnano in 1948. Doksoopulo (1961) reported the discovery of a second hyperaccumulator of nickel, *Alyssum murale*, in 1961. In the early

1970's, Severne and Brooks (1972) discovered a third nickel plant, *Hybanthus floribundus*, in Western Australia. Following this early work, over 200 hyperaccumulators of Ni have now been reported, largely as the result of chemical tests on herbarium specimens.

The term *hyperaccumulator* was first used by Brooks *et al.* (1977a) and was originally used to define plants containing more than 1000  $\mu\text{g/g}$  nickel in dry tissue. This threshold was based on histograms showing discrete plots of two different populations of the same plant genus with no overlap of hyperaccumulators and non-accumulators at the 1000  $\mu\text{g/g}$  Ni boundary (see Figure 1). By 1983, Brooks using mainly herbarium specimens had identified 44 species of the *Alyssum* genus as hyperaccumulators of Ni.

The threshold of hyperaccumulation will vary considerably for different trace elements. Reeves *et al.* (1995) have suggested the following concentrations (in the plant dry matter) as thresholds to define accumulation: 10000  $\mu\text{g/g}$  for Zn and Mn; 1000  $\mu\text{g/g}$  for Ni, Co, Cu, Cr and Pb; and 100  $\mu\text{g/g}$  for Cd and Se. These values represent a concentration about 100 times greater than the normal elemental concentration ranges found in plants.



**Figure 1:** Histograms showing the abundance of nickel in *Alyssum* species. All values  $> 1000$   $\mu\text{g/g}$  (0.1%) are from section *Odontarrhena* of the genus. Source: Brooks (1987).



Almost all metal-hyperaccumulating species known today were discovered on metalliferous soils, either natural or man-made, and almost all hyperaccumulators are endemic to these soils. This suggests that hyperaccumulation is an important ecophysiological adaptation to heavy-metal stress and one of the manifestations of heavy-metal resistance.

For a long time the ability of plants to accumulate metals was considered a detrimental trait. Being at the bottom of many natural food chains, metal-accumulating plants are directly or indirectly responsible for a large proportion of the dietary uptake of toxic heavy metals by humans and other animals. Although some heavy metals are required for life, their excessive accumulation in living organisms is always toxic.

There are three fields in which hyperaccumulators have made significant impacts: mineral prospecting, phytoarchaeology, and 'green remediation' of toxic soils.

#### **Use of Hyperaccumulators in Mineral Prospecting**

Hyperaccumulators can be used for geobotanical exploration whereby the type of vegetation can be used to determine and map areas of mineralisation (Nicks *et al*, 1995). Biological methods of prospecting involve mapping the distribution of plant species confined to certain types of mineralisation. Although there are over 400 hyperaccumulators of nickel, there are only about 20 species that can tolerate and accumulate copper and cobalt. Until recently, these plants were believed to be confined solely to the Shaban Copper Arc in Shaba Province, Democratic Republic of Congo (Zaire) which has the world's richest and most diverse copper flora. However, there have been reports that the Chinese plant, *Elsholtzia haichowensis*, a member of the Lamiaceae (mint family) and confined to copper mines in the Daye region of Hubei province, Central China, is also a hyperaccumulator of copper.

Two species of *Haumaniastrum*, *H. katangense* and *H. robertii* were once known as copper flowers and were used by early prospectors to delineate mineralised ground since they will not grow in soils containing less than 100 µg/g copper and/or cobalt. Brooks *et al*. (1980) have now identified 26 hyperaccumulators of cobalt and 24 of copper. A further 9 taxa were found to be able to hyperaccumulate both metals. All of these 38 plant

species can be used, or have been used, to prospect for copper and/or cobalt in Democratic Republic of Congo (Zaire).

Many hyperaccumulators of nickel have been, or could be used for mineral prospecting. For example, all of the 48 *Alyssum* species with this characteristic indicate the presence of ultramafic substrates and by inference the presence of Ni, Cr and Co that are associated with this rock type. Most hyperaccumulators of nickel are endemic to nickel-rich ultramafic substrates and as such can serve as indicators of this rock type.

The association of the hyperaccumulators *Viola calaminaria* and *Thlaspi calaminare* with calamine zinc deposits in Western Europe has been known for over 100 years. As in the case of nickel, virtually all of the "zinc plants" are geobotanical indicators of zinc and/or other sulphide deposits.

There is only one clearly documented case of a lead hyperaccumulator. *Thlaspi rotundifolium* subsp. *cepaefolium* from Raibl on the Austrian/Italian border has been found to contain 7000 µg/g of this element in dried tissue (Reeves *et al.*, 1983). This species has not yet been used for mineral exploration but does have this potential.

Cannon (1959, 1960, 1964) has reported the use of species of *Astragalus* for geobotanical prospecting for uranium in the western United States. Some *Astragalus* species such as *A. pattersoni* and *A. preussi* are confined to selenium-rich soils and indirectly indicate the presence of uranium in the substrate (selenium and uranium are associated geochemically in the mineral carnotite).

### **Phytoarchaeology and Hyperaccumulators**

The hyperaccumulator *Alyssum corsicum* was first discovered growing over a few hectares of ultramafic rock in the suburbs of Bastia in Corsica. Later however it was found to have its origin in Anatolia (Turkey) on the eastern shores of the Mediterranean. It was concluded that the seed had been brought to Bastia by the Venetians as weed seed in shipments of corn and had colonised this small part of Bastia since it would only grow over ultramafic soils. This small colony of hyperaccumulating *Alyssum* traces an ancient trade route.



The Zaïrean copper flower *Haumaniastrum katangense* can also be used to indicate the presence of archaeological remains beneath the soil. Artisans of the 14<sup>th</sup> Century Kabambian culture of what is now Shaba Province, Democratic Republic of Congo (Zaire) used to smelt copper to produce copper crosses that were used for currency. The smelters were fashioned from abandoned termite mounds and copper ores were brought to the site from elsewhere. After several years, the termite hills weathered to ground level and the residual copper ore formed a toxic ring around the original smelters. This toxic soil became colonised by *H. katangense* and the sites were later recognisable as bare open areas in the savanna. Archaeologists digging below the carpet of this hyperaccumulator of copper have been able to uncover numerous artifacts from the Kabambian and later cultures (Figure 2).



**Figure 2:** Abandoned native copper smelter fashioned from a termite hill at Luisha colonised by *Haumaniastrum katangense* (not in flower) and the fern *Nephrolepis undulata*. Source: Brooks *et al.* (1992).

## Hyperaccumulators and Green Remediation of Soils

Only recently has the value of metal-accumulating plants for environmental remediation been fully realised. At present, remediation technologies consist primarily of removal and replacement, or simply isolation of contaminated sites. Water treatment facilities also do a relatively poor job of removing toxic metals from residential and industrial effluents. Hyperaccumulators can also be used to colonise polluted areas, such as old mine dumps and further to remove pollutants from the soil (Baker *et al.*, 1995). This is termed phytoremediation (removal of pollutants from the soil by use of plants) and is based on the principle that a crop of hyperaccumulating plants may be grown on the heavy metal enriched soil and harvesting of these plants would lower the heavy metal concentrations. Phytomining is a process where these plants may be used to extract heavy metals from low grade ores that would otherwise not be economic (Chaney, 1983; Robinson *et al.*, 1997). Burning the crop produces a 'bio-ore' which can then be sold to small smelting companies as a sulphur-free, non-polluting raw material (Nicks and Chambers, 1994; 1995).

The application of plants in environmental cleanup is an emerging technology that has been called phytoremediation (I Raskin, Grant Proposal #R81869, 1991). Three subsets of this technology are being developed. First, phytoextraction (Salt *et al.*, 1995), in which metal-accumulating plants are used to transport and concentrate metals from the soil into the harvestable parts of roots and aboveground shoots. Second, rhizofiltration (Dushenkov *et al.*, 1995), which uses plant roots to absorb, concentrate and precipitate toxic metal from polluted effluents. Finally, phytostabilisation, which involves the use of plants to eliminate the bioavailability of toxic metals in soils.