

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**Effect of slaughter age between 5 and 14 months on the quality
of sheep meat with a specific focus on collagen concentration
and solubility**

**A thesis presented in partial fulfilment of the requirements for the degree
of**

Masters of Science

in

Animal Science

at Massey University, Manawatu,

New Zealand

Grace Arbogast Mashele

2017

Abstract

In New Zealand, pasture-raised lamb is considered the highest quality sheep meat product while hogget is a downgraded product considered to be of poorer quality. The objectives of this study were to investigate the difference in meat quality between sheep slaughtered at 5, 8 and 14 months-of-age. This study also aimed to develop an assay to measure collagen concentration and solubility and to compare shear force to collagen concentration and solubility for meat from sheep slaughtered at 5, 8 and 14 months. Sixty Romney crossbred rams born in the spring of 2015 were allocated to one of three slaughter age groups: 5-group (n=20), 8-months group (n=20) and 14-months group (n=20). The ram lambs were grazed together on perennial ryegrass-based pasture. At slaughter, the loin (*Longissimus lumborum*) from the left side of the carcass was excised, chilled for 24 hours and then frozen. Shear force, pH, colour, drip loss, myofibrillar fragmentation index, sarcomere length, soluble collagen, insoluble collagen and total collagen were measured on the loin. *Longissimus lumborum* muscle from the 5-months group was darker (P=0.045) with higher pH (P<0.001) than 8 and 14-months groups. *Longissimus lumborum* muscle from the 14-months group was redder (P<0.001), yellower (P<0.001) with higher intramuscular fat (0.003), shorter sarcomere length (P<0.001), lower collagen concentration (P=0.020) and lower soluble collagen in percentage (P=0.007) and in g/100g fresh weight (P=0.008) than 5 and 8-months groups. The peak shear force was lower at 14 months, intermediate at 5 months and greater at 8 months (P<0.001). *Longissimus lumborum* muscle from the 8-months group had greater drip loss (P<0.01) than 5-months group and 14-months group. The results suggest that slaughtering sheep at 14 months could have benefits on eating quality due to an increase in intramuscular fat.

Keywords: lamb; slaughter; age; meat; quality

Acknowledgements

First and foremost, I would like to express my heartfelt gratitude to my Supervisors Dr. Nicola M. Schreurs (Institute of Veterinary, Animal and Biomedical Sciences, Massey University) and Dr. Michael E. Parker (Institute of Food Science and Technology, Massey University) for their support, guidance, patience and encouragement throughout my study. My studies would not have been possible without them.

I would like to express my sincere appreciation to Massey University's Keebles farm for sourcing the animals and the farm technical staff for looking after the animals and weighing the sheep on-farm and prior to transport for slaughter. I would like to express my special thanks to Alliance Group processing plant in Dannevirke for their services of slaughtering the sheep, boning out and packaging of the loin samples and providing VIAscan data.

My special thanks to Kelvin Stafford and Debbie Hill (Institute of Veterinary, Animal and Biomedical Sciences, Massey University) for their administrative support throughout my study. I thank Peter Wildbore (Institute of Veterinary, Animal and Biomedical Sciences, Massey University) with assistance on purchasing of reagents used in my experiments. I thank Steve Glasgow (Institute of Food Science and Technology, Massey University) for his patience, guidance and advice throughout my time in his laboratory.

I am grateful to the New Zealand AID Scholarship through the Ministry of Foreign Affairs and Trade for funding my Masters programme. My special thanks to International student support office staff: Silvia Hooker, Jamie Hooper, Logan Tate and Dianne Reilly for their encouragement, support and advice throughout my studies.

I am grateful to my family and friends for their encouragement and advice throughout my studies.

Table of Contents

Abstract	i
Acknowledgements	ii
Table of Contents	iii
List of Tables	vii
List of Figures.....	ix
Chapter 1. Literature Review.....	10
1.1 Introduction	10
1.2. The quality of meat from sheep	10
1.2.1 Effect of sheep slaughter age on colour of lean meat.....	12
1.2.2 Effect of sheep slaughter age on colour of fat.....	15
1.2.3 Effect of sheep slaughter age on meat flavour	16
1.2.4 Effect of sheep slaughter age on meat juiciness.....	17
1.2.5 Effect of sheep slaughter age on meat tenderness	18
1.2.6 Collagen in meat.....	19
1.2.6.1 Measurement of collagen concentration and solubility	20
1.2.6.2 Collagen concentration and meat tenderness	23
1.2.6.2 Collagen solubility and meat tenderness	27
1.2.6.3 Effect of collagen concentration and solubility on meat tenderness	32
1.3 Research Objectives.....	33
1.3.1 Research objectives and hypothesis	34
Chapter 2. Materials and Methods for Objective Meat Quality Test.....	35
2.1 Animal management and slaughter	35
2.2 Carcass measurement and meat sampling.....	35

2.3 Preparation and partitioning of the loin sample	36
2.5 Soluble, insoluble and total collagen determination.....	40
Chapter 3. Collagen Assay Development	41
3.1 Introduction	41
3.2. Methods in Literature	42
3.3 Outline of method to be tested and developed	43
3.3.1 Preparation of hydroxyproline standard solution and standard curve	43
3.3.2 Determination of total collagen concentration	44
3.3.3 Determination of heat soluble and insoluble collagen in lamb meat.....	44
3.3.4 Preparation of buffer solution	45
3.3.5 Preparation of oxidant solution	45
3.3.6 Preparation of colour reagent solution (Ehrlich's reagent)	45
3.3.7 List of equipment and chemical	45
3.4 Considerations in the method development.....	46
3.4.1 pH of hydrolysate	46
3.4.2 Standard curve and sample dilution	47
3.4.3 Colour stability	50
3.5 Outcomes of initial tests	51
3.5.1 Total collagen concentration of lamb meat samples – assessing the methodology on samples.....	51
3.5.2 Soluble and insoluble collagen of lamb meat sample.....	52
3.5.3 Conclusion	54
3.6 Further considerations.....	54
3.7.1 Preparation of buffer solution	55
3.7.2 Preparation of oxidant solution	55

3.7.3 Preparation of colour reagent solution (Ehrlich’s reagent).....	56
3.7.4 Preparation of standard solution	56
3.7.5 The method for determination of heat soluble, insoluble and total collagen in lamb meat	56
Chapter 4. Meat Quality of Sheep Slaughtered at 5, 8 and 14 Months of Age.....	58
4.1 Introduction	58
4.2 Materials and methods.....	58
4.3 Results.....	59
4.3.1 Sheep live weights and carcass characteristics.....	59
4.3.2 Objective meat quality characteristics.....	61
4.3.3 Total collagen, soluble collagen and insoluble collagen	61
4.3.4 Interrelationships	62
4.4 Discussion	64
4.5 Conclusion.....	67
Chapter 5. General Discussion	67
5.1 Collagen method development	67
5.2 Effect of slaughter age on meat quality.....	68
5.3 Relationship between shear force and collagen concentration.....	69
5.4 Relationship between shear force and collagen solubility.....	69
5.5. Conclusion.....	70
References	70
Appendix.....	81
Appendix 1. Soluble, insoluble and total collagen of 14 samples per treatment in two replicates for 5, 8 and 14 month old Romney rams	81

Appendix 2: Ranges, standard deviation, means and coefficient of variation of soluble and insoluble collagen of 14 samples per treatment in two replicates for 5, 8 and 14 month old Romney rams.84

Appendix 3. Hydroxyproline standard curves for each day of analysis of samples from the slaughter age experiment.....85

List of Tables

Table 1: Effect of slaughter age on colour and tenderness of meat from different muscles of sheep.	14
Table 2. Studies investigating collagen concentration and collagen solubility in the Longissimus lumborum or several muscles of sheep.....	24
Table 3. Collagen concentration and collagen solubility of ovine different muscle and different age (Cross et al., 1972).	26
Table 4. Age at slaughter, collagen concentration, collagen solubility, tenderness rating (9 = like extremely; 1 = dislike extremely) and shear force in sheep semimembranosus muscles.	32
Table 5. List of equipment and use	46
Table 6. List of chemicals and use	46
Table 7. Absorbance values for standards	47
Table 8. Absorbance of Spring 165 green sample using 10, 25 and 50 dilution factor.	48
Table 9: The absorbance value of higher hydroxyproline concentration	49
Table 10: Absorbances of Spring 165 green sample using 50 and 100 dilution factor..	49
Table 11: Average soluble, insoluble and total collagen of 6 lamb meat samples utilised to assess the collagen assay that was developed.	53
Table 12: Final on-farm live weight, actual slaughter age, carcass weight, dressing out percentage, GR soft tissue depth, leg yield, loin yield, shoulder yield, lean meat yield and intramuscular fat for 5, 8 and 14 months old Romney rams. Values are least square means \pm standard error of the mean.	60
Table 13. Meat quality attributes of <i>Longissimus lumborum</i> of Romney ram slaughtered at 5, 8 and 14 months old assessed using objective measures on twenty samples per age treatment with each sample coming from a different animal. Values are the least square mean \pm standard error of the mean	62
Table 14: Pearson correlation coefficients and P-values of peak force (kgF), initial yield (kgF), work done (kgF), intramuscular fat (%), insoluble collagen (g/100g fresh weight), soluble collagen (g/100g fresh weight) and total collagen (g/100g fresh weight) of Romney rams slaughtered at 5, 8 and 14 months old.	63

Table 15. Step wise regression using forward selection to consider the influence of soluble collagen (g/100g fresh weight), myofibrillar fragmentation index (%), insoluble collagen (g/100g fresh weight), sarcomere length (μm), express juice (cm^2/g) and intramuscular fat (% whole muscle) on peak shear force (kgF). 64

List of Figures

Figure 1. Structure of skeletal muscle showing three connective tissues associated with the muscle structure (Pearson, 2012).	20
Figure 2. Shear force as a function of collagen concentration. Data are plotted by age i.e., 0 means new born and 1,2,3 and 4 months-of- age (Young et al., 1993).....	27
Figure 3 (a). Collagen cross-link structure in young animal with collagen molecule linked head to tail to form an infinite polymer. b. Showing collagen cross-link in older animal as it passes between the fibrils. Adopted from Purchas et al. (1989).	28
Figure 4. Shear force as a function of collagen solubility. Data are plotted by age (0 means new born and 1, 2, 3 and 4 months- of- age). Lower shear force values are associated with meat that is more tender (Young et al., 1993).....	30
Figure 5: Decline in collagen solubility of sheep Longissimus muscle with age. Adopted from Berge et al. (2003).	30
Figure 6. Shear force for muscle samples cooked at 60 or 80 °C for 90 min from weaners 2–3 months, lambs 8–10 months, 2 tooth (12-19months), 4 tooth (18-24 months), 6 tooth (23-36 months) and 8 tooth (28-48 months). (Adapted from (Bouton et al., 1978) also presented in (Hopkins & Mortimer, 2014).	31
Figure 7. A schematic diagram showing the partitioning of lamb <i>Longissimus lumborum</i> muscle for meat quality analysis. MFI= Myofibrillar fragmentation index, EJ= Express juice, SL = Sarcomere length, IMF = Intramuscular fat.	37
Figure 8. Hydroxyproline standard curve for estimating total hydroxyproline in meat sample	48
Figure 9. Hydroxyproline standard curve of higher hydroxyproline concentration.	49
Figure 10. Change in absorbance with time.....	50
Figure 11. Hydroxyproline standard curve used to estimate total collagen in meat sample	51
Figure 12. Hydroxyproline standard curve to determine total collagen of Spring 165 green meat sample when the experiment was repeated.....	52
Figure 13. Hydroxyproline standard curve of the 6 lamb meat samples.....	54
Figure 14: On-farm live weight for actual slaughter age 5.1, 8.2 and 14.5 months old Romney rams. Points are least squares means with standard error bars.	60