

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

The effect of Rabbit
(*Oryctolagus cuniculus cuniculus*)
(Linnaeus, 1758)
Browsing on Seedling Survival.

A thesis presented in part fulfillment of the requirements

of the degree of

Master of Science
in
Zoology

At Massey University, Palmerston North
New Zealand.

Paul Andrew Doyle

2004

“From the extraordinary manner in which European productions have recently spread over New Zealand, and have seized on places which must have previously occupied, we may believe, if all the animals and plants from Great Britain were set free in New Zealand, that in the course of time a multitude of British forms would become thoroughly naturalized there and would exterminate many of the natives.”

Darwin, Charles. The Origin of the Species. 1859.

ABSTRACT

The European rabbit, *Oryctolagus cuniculus* (Linnaeus 1758) is a small 2–4 kg eutherian mammal introduced into New Zealand from the United Kingdom in the mid 1800's. By the 1950's the range of the rabbit had covered nearly 95% of the available land resulting in millions of dollars of damage. It was noticed by the 1970's that some populations were stable without any control measures being applied, and with increasing research and a changing attitude towards pest management away from extermination to control, it became policy that the status of this pest was downgraded to a regional pest rather than a pest of national importance. Control became the responsibility of the landowner rather than a national body except where populations occurred in high densities.

However, even rabbit numbers at low densities do exert a pressure on vegetation, and in some bush remnants rabbits are blamed for the lack of regeneration that occurs even when larger herbivores are excluded.

Four bush remnants within the Manawatu-Rangitikei region of New Zealand, two in the coastal sand country and two inland at Marton, were studied for the causes of seedling mortality over five seasons from August 1999 to December 2000.

One hundred and fifty seedlings were identified and numbered to species level and were placed in five treatments and controls at each site, except Legg Estate bush remnant where ninety seedlings were used in unbalanced numbers in each replicate

The data were analysed using logistic regression with scores for rabbit densities and deaths of seedlings per bush remnant within the treatments and controls.

The analysis suggests that there is a seasonal effect ($P < 0.0001$) and a treatment /control effect ($P = 0.0002$) on seedling survival with treatments at Legg Estate bush

remnant surviving consistently better over the five seasons.

At Himatangi Block Scientific Reserve seedlings in the treatment survived better than seedlings in the controls, except for winter 1999.

At Monkton's and Fulleton-Smith bush remnants, treatments survived better than in the controls.

Overall the treatment (protecting seedlings against rabbits by fencing) improves seedling survivability. This trial was conducted at a time of regionally low pest numbers because of the prior introduction of Rabbit Haemorrhagic Disease (RHD), and treatment results should be interpreted with this in mind.

Key Words: Rabbit, Treatment effect, Bush remnant, RHD, Exclosure, Seedling survival.

List of photographs

- Photograph I: An nine-year-old adult male rabbit Page 6
- Photograph II: Legg Estate bush remnant as seen from the south Page 23
- Photograph III: Himatangi Block Scientific Reserve showing a dune of the Motuiti phase. On the ridge crest is *Hedycarya arborea*, *Knightia excelsa* and *Beilschmiedia tawa*, with a variety of introduced weeds and grasses in the foreground..... Page 26
- Photograph IV. Monkton's bush remnant viewed from the east. Scrub in the foreground is backed by a mixed podocarp / tawa forest..... Page 29
- Photograph V: Fulleton-Smith bush remnant showing many large examples of native trees with a good understorey. Rabbit field sign was common in the pasture in the foreground Page 32
- Photograph VI: An adult male sambar deer Page 41
- Photograph VII: A typical light-well within a bush remnant. This photograph was taken in Legg Estate bush remnant Page 53
- Photograph VIII: An insect fall trap used to collect the seed rain. In the background is a seedling survival control plot. This photograph was taken in Legg Estate bush remnant Page 73

Photograph IX: Field sign of rabbits seen in Legg Estate bush remnant in November 2003. Scratchings in the sandy soil, faecal pellets and a urine squirt are clearly present. Rabbit field sign similar to this was plentiful within and around Legg Estate bush remnant at this time Page 139

Unless otherwise stated photographs were taken by Mrs Cheryl Doyle

List of maps

Map 1: Map of lower North Island showing the four bush remnants of the seedling survival trial. Legg Estate (1) and Himatangi Block Scientific Reserve (2) are collectively known as the sand country sites while Monkton's bush (3) and Fulleton-Smith bush (4) and collectively known as the inland sites Page 20

Map II: Ranges of sambar deer in the Manawatu/Rangitikei region. The outline shows the range (red outline) now thought to be occupied from discussion with a sambar deer stalker compared with its officially recognised range (hatched). The seedling survival trial sites are 1) Legg Estate bush remnant, 2) Himatangi Block Scientific Reserve, 3) Monkton's and 4) Fulleton-Smith bush remnants. Page 42

List of diagrams

Diagram I: The setup of a replicate in the irrigation experiment at Legg Estate bush remnant shows distances between each cage, placement of rain gauge and the areas irrigated and where the samples were taken. All replicates were of an east-west orientation. Page 64

Diagram II: Drawings of the seeds of native plant species used for the Chi-squared calculations, showing the morphological and their scientific species name. ... Page 75

List of tables

- Table I: Score means for rabbit field sign per season for the four seedling survival trial sites in the Manawatu-Rangitikei region, North Island, New Zealand Page 16
- Table II: Environmental data on the four seedling survival trial sites in the Manawatu-Rangitikei region, North Island, New Zealand Page 36
- Table III: Checks to determine the accuracy of the method of calculating missing values for lux readings at the treatment and controls of the seedling survival trial.
..... Page 55
- Table IV: Mean values for Lux reading at the four bush remnants involved in the seedling survival trial. T and C represent the values obtained for treatments and controls respectively..... Page 55
- Table V: Calculated seasonal and a regional means for unobstructed sunlight in units of Lux taken as part of the seedling survival trial at four bush remnants in the Manawatu-Rangitikei region..... Page 57
- Table VI: The mean and standard error for drying time of five replicates of irrigated Ashurst silt loam, (stony phase), at a 150^o Celsius for the calibration of an Atlas Neptune™ oven for the optimal drying times of OA horizons samples..... Page 61
- Table VII: Mean numbers of seedlings present from the five replicates of the irrigation experiment at Legg Estate bush remnant, Palmerston North, New Zealand
..... Page 65

Table VIII: Individual counts of seeds collected as seed rain from Legg Estate bush remnant and Himatangi Block Scientific Reserve, south of Palmerston North, in the Manawatu region, North Island, New Zealand from April 2000 to January 2001

..... Page 69

Table IX: Individual counts of seeds collected as seed rain from Monkton's and Fullerton-Smith bush remnants near Marton in the Rangitikei region, North Island, New Zealand from April 2000 to January 2001..... Page 77

Table X: Standardised residuals calculated from the seed rain for eight plant species at the four seedling survival trial sites in the Manawatu-Rangitikei region, North Island, New Zealand..... Page 78

Table XI: Plant species germinated from approximately 500 grams of OA horizon material sampled from Legg Estate bush remnant, Palmerston North, New Zealand. Native plant species are shown in bold type and account for 14.3% (1 out of 7) of the plant species present and 1.2 % (1 out of 85) of individual seedlings germinated

..... Page 82

Table XII: Plant species germinated from approximately 500 grams of OA horizon material sampled from Himatangi Block Scientific Reserve, Palmerston North, New Zealand. Native plant species are shown in bold type and account for 25 % (2 out of 8) of the plant species present and 27 % (10 out of 37) of the individual seedlings germinated..... Page 83

Table XIII: Plant species germinated from approximately 500 grams of OA horizon material sampled from Monkton's bush remnant, Marton, New Zealand. Native plant species are shown in bold type and make up 18.2 % (2 out of 11) of the plant species present and 27.7 % (23 out of 83) of the individual seedlings germinated..... Page 84

Table XIV: Plant species germinated from approximately 500 grams of OA horizon material sampled from Fulleton-Smith bush remnant, Marton, New Zealand. Native plant species are shown in bold type and account for 60 % (6 out of 10) of the plant species present and 73.3 % (66 out of 90) of the individual seedlings germinated.
..... Page 84

Table XV: Plant species distribution and total plant species count per site for the seedling survival trial sites in the Manawatu-Rangitikei area North Island New Zealand..... Page 95

Table XVI: Means growth per season, sample size and standard errors of individual native plant species within treatments and controls in the seedling survival trial at Legg Estate bush remnant, Palmerston North, New Zealand from August 1999 to December 2000 Page 97

Table XVII: Means growth per season, sample size and standard errors of individual native plant species within treatments and controls in the seedling survival trial at Himatangi Block Scientific Reserve, Palmerston North, New Zealand from August 1999 to December 2000 Page 98

Table XVIII: Means growth per season, sample size and standard errors of individual native plant species within treatments and controls in the seedling survival trial at Monkton' bush remnant, Marton, New Zealand from August 1999 to December 2000
..... Page 99

Table XIX: Means growth per season, sample size and standard errors of individual native plant species within treatments and controls in the seedling survival trial at Fulleton-Smith bush remnant, Marton, New Zealand from August 1999 to December 2000..... Page 101

List of graphs

- Graph I: Graph of rabbit numbers across the Horizons mw region. With the introduction of RHD in 1998 population reductions of 45% occurred and at the new lower levels the seedling survival trials commenced in August 1999 and concluded in December 2000 Page 18
- Graph II: Mean values per season of unobstructed sunlight in units of Lux at four bush remnants in the Manawatu-Rangitikei Page 58
- Graph III: Mean weights of five replicates of Ashurst Silty Loam (Stony phase) verses drying time in an Atlas Neptune™ at 150^o Celsius to obtain three consistent weights. This time was assumed to be optimal drying time for OA horizon material from the irrigation trial and background OA moisture levels at the four bush remnants of the seedling survival trial North Island, New Zealand..... Page 62
- Graph IV: OA horizons moisture levels of treatments (water addition) and controls (no water addition) at five replicates as part of the irrigation trial at Legg Estate bush remnant, Palmerston North from 15 October 2000 (Day 1) to 11 February 2001 (Day 119). The unobstructed rainfall is shown for this bush remnant was obtained from a climate station 700 metres from the bush remnant edge Page 66
- Graph V: OA horizon percentage moisture content at the four bush remnants of the seedling survival trial from October 2000 to January 2001. The data for Legg Estate bush remnant was independent of the irrigation trial Page 70
- Graph VI: Seasonal growth changes for *Alectryon excelsus* (top) and *Hoheria populena* (bottom) at Legg Estate bush remnant in the seedling survival trial August 1999 to December 2000 Page 104

-
- Graph VII: Seasonal growth changes for *Macropiper excelsum* (top) and *Beilschmiedia tawa* (bottom) at Legg Estate bush remnant in the seedling survival trial August 1999 to December 2000 Page 105
- Graph VIII: Seasonal growth changes for *Beilschmiedia tawa* (top) and *Macropiper excelsum* (bottom) at Himatangi Block Scientific Reserve in the seedling survival trial August 1999 to December 2000 Page 106
- Graph IX: Seasonal growth changes for *Hedycarta arborea* (top) and *Parsonia heteraophylla* (bottom) at Himatangi Block Scientific Reserve in the seedling survival trial August 1999 to December 2000 Page 107
- Graph X: Seasonal growth changes for *Knightia excelsa* (top) and *Macropiper excelsum* (bottom) at Himatangi Block Scientific Reserve in the seedling survival trial August 1999 to December 2000 Page 108
- Graph XI: Seasonal growth changes for *Hedycarya arborea* (top) and *Lophomyrtus obcordate* (bottom) at Monkton's bush remnant in the seedling survival trial August 1999 to December 2000 Page 109
- Graph XII: Seasonal growth changes for *Hoheria populnea* (top) and *Alectryon excelsus* (bottom) at Monkton's bush remnant in the seedling survival trial August 1999 to December 2000 Page 110
- Graph XIII: Seasonal growth changes for *Beilschmiedia tawa* (top) and *Coprosma grandifolia* (bottom) at Monkton's bush in the seedling survival trial August 1999 to December 2000 Page 111

-
- Graph XIV: Seasonal growth changes for *Alectryon exclesus* (top) and *Hedycarya arborea* (bottom) at Fulleton-Smith bush remnant bush in the seedling survival trial August 1999 to December 2000..... Page 112
- Graph XV: Seasonal growth changes for *Beilschmiedia tawa* at Fulleton-Smith bush remnant in the seedling survival trial August 1999 to December 2000..... Page 113
- Graph XVI: Seedling survival per season at Legg Estate bush remnant (top) and Himatangi Block Scientific Reserve (bottom) in the sand country area in the Manawatu region, North Island New Zealand from spring 1999 to spring 2000.
..... Page 117
- Graph XVII: Seedling survival per season at Monkton's bush remnant (top) and Fulleton-Smith bush remnant Reserve (bottom) at the inland sites in the Rangitikei region, North Island New Zealand from spring 1999 to spring 2000. Page 118
- Graph XVII: Seedling survival in Legg Estate bush remnant from spring 1999 to spring 2003. This trial was an extension of the seedling survival trial, which was left set up at the request of the land owners to provide ongoing data to assist in management decisions on this remnant's restoration..... Page 140

Acknowledgements

I would firstly like to thank my supervisor Professor Brian Springett for his patience and guidance through this thesis. As this work was partly funded by the Regional Council, I would also like to thank Mr Malcolm Todd of the Manawatu Wanganui regional council trading as Horizons mwtm for his input and his liaison with the land owners involved, as well as the council for the funding.

A special thank you to the landowners who allowed me access to their property on a regularly basis. Mr and Mrs Boulton and Mrs Legg of Legg Estate at Paranui, Palmerston North, Mr and Mrs Fulleton-Smith of Holtby Farm, Mount Curl, Marton and Mr and Mrs Monkton of Monkton Farm, Fern Flats, Marton.

A thank you to the people in the Wanganui and Palmerston North offices of Department of Conservation who allowed access to Himatangi Block Scientific Reserve and to Miss Vivian Nicholls of the Palmerston North office Department of Conservation who accessed a wealth of information on Himatangi Block Scientific Reserve.

To Kay Treader of National Institute of Water and Atmospheric Research (NIWA) who kindly provided background climate data for the irrigation trial at Legg Estate and for general climate data for the regions where the sites were situated, a grateful thank you.

A thank you to Manawatu Timber for the timber supplied free of cost for the posts of the exclusion plots.

tm Trade mark.

Mr Eddie and Margrat Millard of Bainesse, thank you for the rain fall data used in the irrigation trial as the values for background rainfall.

When setting up this trial I had many discussions with friends and professionals alike and wish to acknowledge the input of Mr Lex Foote, Miss Shona Brock, Mr Jarn Godfrey, Miss Geraldine Moore and Mr Mark Bell.

I wish to acknowledge the help with the initial statistical calculations of Mr Chris Hunt at agresearch, Palmerston North and Mr Duncan Heederley and Mr Alistair Nobel of Statistics Research and Consulting Centre, also Mr John Brock, Dr Mike Christian and Miss Yvonne Grey of agresearch for their help with the plant identification.

For help with data entry and proof reading, being a true and loyal Godmother to my son Jarn, a definite thank you to Miss Alison Southgate.

For help with obtaining mapping software I would like to thank Jean Flack of Polymedia[™], Nelson.

To Mr. Richard Morton, a thank you for your help with discussion on the range, activities and behavior of the sambar deer.

To my friends Peter Tunnicliff and Paul Morris and their families from Wellington who have given encouragement over the years thanks guys.

[™] Trade mark

To Ms Jody Inhang and Ms Erica Reid and technical support staff Mr. Paul Barrett of the Ecology Department at the Institute of Natural Resources, Massey University deserve a special mention as does the Graham Christensen of the Practical Teaching Complex for allowing use of the departments stock exclusion cages for the irrigation experiment at Legg Estate bush remnant.

To Mrs. Barbara Just, Senior Technician at the Ecology Department, for working wonders with the paper work for the account "rabbits" with Horizons mw. A big Thank You.

Jason Jones is thanked for his help in keeping my computer up to spec.

Overall I would like to thank my wife, Mrs Cheryl Doyle for her help and encouragement thought out this trial and who was assisting with data collection at Legg Estate while in the first stages of labour (and did not tell me) the day before the birth of our darling daughter Cassandra Ruby-Jo and two and a half years later on our beautiful son Jarn Alexander Doyle.

The support of friends and relatives, who have given me the encouragement I needed, especially when the going was tough, was greatly appreciated. To Mr Shane Hatcher, Miss Kerry McCracken, Miss Jennifer Neale, Ms Jacquin Ward, Ms Jackie (the rock) Tribe, Miss Sharon Dearlove, Dr David and Mrs Dorothy Page, Mr Bill and Mrs Olive Mills, Mr Nigel and Dr Jade Kerr and Ms Josey Randall and Mrs Joy and the now late Dr Colin Coates, Ms Siena Corcoran and Patch M^cKenna. To Bill Koko and Scott Shaw of Lambpackers tm and Bruce Cable, Dawn Mack and Greg Veale of Asure tm, thank you for your much needed encouragement to complete this document.

tm Trade mark

A remembrance of the memory of my childhood friend, teacher and hunting companion, my sheepdog Flynn (named after Barry Crumps dog of the same name in New Zealand's greatest work of fiction, *A Good Keen Man*), who taught me to rabbit, and in many ways laid the foundation work over thirty years ago, which has accumulated to date in this project.

To my friend of 25 years, Dr Christine Wilson, who upon receiving her doctorate motivated me to do this level of study, Thanks Chrisie.

To those who said this could not be done thank you for laying the ground work for this achievement.

To you all mentioned and to those through no deliberate fault of my own who I have omitted to mention, but whom definitely do deserve recognition, I would like to say

A SINCERE

THANK YOU!

Table of contents

Abstract	I
List of photographs	III
List of maps	V
List of diagrams	VI
List of tables	VII
List of graphs	XI
Acknowledgements	XIV
Tables of contents	XIII

Chapter 1 General introduction, species and site data

1:1. Introduction	Page 1
1:2:1. The rabbit, history and biology	Page 5
1:2:2. Classification.....	Page 5
1:2:3. Ideal habitats	Page 7
1:2:4. Life cycles in New Zealand	Page 8
1:2:5. Population fluctuations and its pest status within New Zealand.....	Page 8
1:2:6. The political history of the rabbit problem	Page 9
1:2:7. Conclusions	Page 13
1:3:1. Regional and site presence of rabbits.....	Page 14
1:3:2. Methods.....	Page 15
1:3:3. Rabbit numbers during the trial period	Page 16
1:3:4. Conclusions and general remarks	Page 17
1:4:1. Regional and site data for the four bush remnants.....	Page 19
1:4:2. Climate	Page 19
1:4:3. The sand country sites.....	Page 21
1:4:3:1. Stratigraphy	Page 21
1:4:3:2. Soils.....	Page 22
1:4:4:1. Legg Estate bush remnant	Page 22
1:4:4:2. Physical description	Page 22
1:4:4:3. Vegetation	Page 24
1:4:4:4. Legal status	Page 25

1:4:5:1. Himatangi Block scientific Reserve.....	Page 25
1:4:5:2. Physical Description	Page 25
1:4:5:3. Vegetation	Page 25
1:4:5:4. Legal status	Page 27
1:4:6. The inland sites	Page 28
1:4:6:1. Stratigraphy	Page 28
1:4:7:1. Monkton's bush remnant	Page 30
1:4:7:2. Physical description	Page 30
1:4:7:3. Soils.....	Page 30
1:4:7:4. Vegetation	Page 30
1:4:7:5. Legal status	Page 31
1:4:8:1. Fulleton-Smith bush remnant.....	Page 31
1:4:8:2. Physical description	Page 31
1:4:8:3. Soils.....	Page 33
1:4:8:4. Vegetation	Page 33
1:4:8:5. Legal status	Page 34
1:4:9. Impressions and conclusions.....	Page 34
1:5:1. The problem defined	Page 37
1:5:2. Objectives.....	Page 39
1:6:1. Sambar deer in the Rangitikei-Manawatu region	Page 40
1:6:2. Relevant biology and behaviour	Page 43
1:6:3. The deer hypothesis	Page 44
1:6:4. Implications for the trial.....	Page 45
Bibliography	Page 46

Chapter 2 Light levels and the irrigation experiment.

2:1:1. Light levels.....	Page 52
2:1:2. Materials and methods	Page 54
2:1:3. Results of lights levels at remnant floor.....	Page 54
2:1:4. Results.....	Page 56
2:1:5. Conclusions.....	Page 56
2:2:1. Results for canopy light levels	Page 56
2:2:2. Conclusions for canopy light level results	Page 56
2:3:1. OA horizons soil moisture	Page 59
2:3:2. Materials and methods	Page 60
2:3:3. Calibration of the neptune oven.....	Page 60

2:3:4:1. Irrigation experiment at Legg Estate bush remnant.....	Page 63
2:3:4:2. Materials and methods	Page 63
2:3:4:3. Results.....	Page 67
2:3:4:4. Conclusions.....	Page 68

Bibliography.....	Page 71
-------------------	---------

Chapter 3 Seed production and seed rain.

3:1:1. Seed production.....	Page 72
3:1:2. Materials and methods	Page 72
3:1:3. Results.....	Page 74
3:1:4. Conclusions.....	Page 78

3:2:1. OA horizons soil seed bank	Page 80
3:2:2. Materials and methods	Page 80
3:2:3. Results.....	Page 81
3:2:4. Conclusions.....	Page 88

Bibliography.....	Page 90
-------------------	---------

Chapter 4 Seedling Growth and survival.

4:1:1. Seedling growth	Page 91
4:1:2. Materials and methods	Page 91
4:1:3. Results.....	Page 102
4:1:4. Conclusions.....	Page 114

4:2:1. Seedling survival.....	Page 116
4:2:2. Results.....	Page 116
4:2:3. Conclusions.....	Page 119

Bibliography.....	Page 121
-------------------	----------

Chapter 5 Conclusions and post script.

5:1. Discussion.....	Page 122
----------------------	----------

Bibliography.....	Page 129
-------------------	----------

Appendices

Appendix I. Scientific and common names of species mentioned in this work	Page 130
Appendix II. Chi-squared distribution	Page 134
Appendix III. Rainfall (in mls) recorded at Bainesse, 700 metres from Legg Estate bush remnant, Palmerston North.....	Page 135
Appendix IV. Drying times for five replicates of an Ashurst silty loam (stony phase) at 150°C to calculate the optimal drying time for the samples of the irrigation experiment	Page 136
Appendix V. t distribution.....	Page 137
Appendix VI. Post script – Legg Estate in spring 2003 and 2004.....	Page 138
Bibliography	Page 143