

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**Effect of various whey protein supplements  
on recovery from prolonged endurance exercise in  
trained cyclists**

**A thesis presented in partial fulfilment of the requirements for the  
degree of**

**Master of Science  
in  
Nutritional Science**

**at Massey University, Palmerston North,  
New Zealand.**

**Dean Mathew Rankin**

**2005**

## ABSTRACT

### **Background:**

Protein-containing recovery beverages are proposed to support an athlete's recovery from exercise through stimulation of insulin release, promoting the restoration of muscle glycogen stores, and stimulation of protein synthesis and muscle protein restoration.

### **Objective:**

The present study aimed to determine, (1) whether there is an insulinotropic effect of whey proteins, when consumed in addition to carbohydrate, which is assumed to enhance muscle glycogen resynthesis and (2) whether a blend of hydrolysate and intact protein, when consumed in addition to carbohydrate, will enhance the athlete's recovery from exercise.

### **Design:**

Twelve trained top level cyclists repeated a protocol on four consecutive weeks, during which either a control beverage (Carb) or three beverages containing whey protein (carbohydrate and intact protein (Carb + I); carbohydrate and protein hydrolysate (Carb + H; carbohydrate and intact protein : protein hydrolysate mix (Carb + M)) were consumed during recovery from exhaustive endurance exercise. The beverages were formulated to supply 1.2 g/kg/hour carbohydrate and 0.4 g/kg/hour protein. Subjects followed a controlled diet two days before each experimental day. On the experimental day the athletes each performed a glycogen-depleting exercise programme, then received the designated dietary beverage every 30 minutes for the first two hours post-exercise. The progress of recovery was monitored via the measurement of cardiovascular recovery, and the appearance and relative concentration of metabolites in blood (15 samples over a four hour period, obtained via an indwelling cannula) and urine samples (13 samples over a seven hour period) collected sequentially during the post-exercise recovery period.

### **Results:**

Plasma albumin concentrations were significantly lower following consumption of beverages containing whey protein (Carb + H,  $p < 0.01$ ; Carb + M,  $p < 0.05$ ) compared to

that observed with the Carb beverage. Urine output was significantly higher after consumption of the Carb beverage than with any of the three-protein containing beverages (Carb + I,  $p < 0.01$ ; Carb + H,  $p < 0.05$ ; Carb + M,  $p < 0.05$ ) during the period of controlled fluid consumption. Heart rate recovery was found to be significantly greater following consumption of the three protein-containing beverages (Carb + I,  $p < 0.001$ ; Carb + M,  $p < 0.001$ , Carb + H,  $p < 0.01$ ) than following consumption of the Carb beverage. The Carb + M beverage produced increased heart rate recovery ( $p < 0.001$ ) compared to that observed following consumption of the other two protein-containing beverages (Carb + I, Carb + H). Following correction of the data for haematocrit, to account for the hydration status of the athletes, a significant difference ( $p < 0.05$ ) in the ratio of plasma insulin to plasma glucose concentrations was found following consumption of any of beverages containing whey protein (Carb + I, Carb + H, Carb + M) compared to that observed for the Carb beverage. Consumption of the Carb + I beverage resulted in significantly higher concentrations of urinary nitrogen excretion as urea ( $p < 0.05$ ) and ammonia ( $p < 0.01$ ), and significantly higher plasma concentrations of the amino acids Valine, Leucine, Isoleucine, Phenylalanine, Tryptophan, and Tyrosine ( $p < 0.05$ ).

### **Conclusions:**

The addition of whey protein to a carbohydrate-containing beverage stimulated enhanced recovery from exercise. A major factor in the improved recovery was increased rehydration following consumption of the protein-containing beverages, mainly due to the high sodium content of these beverages. This increased rehydration was shown to influence results for plasma insulin and plasma glucose concentrations where, after accounting for the hydration status of athletes, a difference between consumption of the Carb beverage and that observed for any of the three protein-containing beverages was observed. The results also allude to a potential benefit of protein hydrolysates over intact protein on protein recovery. Consuming a protein mix (Carb + M) also appears to improve heart rate recovery compared to consuming either intact (Carb + I) or hydrolysed (Carb + H) proteins individually. The results of this study highlight the importance of dietary protein on enhancing recovery from endurance exercise.

## ACKNOWLEDGEMENTS

I would like to express my sincere appreciation to my supervisors: Dr Alison Darragh (for her involvement right throughout this extensive project in all aspects of its planning, implementation, analysis and thesis writing); Dr Roger Lentle (for his involvement in the experimental trial, statistical analysis and thesis writing); and to Associate Professor Hugh Morton for his statistical help during the initial stages of the study. Their advice, guidance and encouragement throughout the study was excellent. I would also like to thank them for both their patience and their understanding of my other commitments outside of this thesis, which prolonged this project.

I would also like to thank Chris Booth for her time and efforts in the planning and implementation of this study, especially the time she put in during the experimental trial in the processing of samples. In addition, thanks also to Lena Lückel (from Germany) for volunteering her time and effort while in New Zealand to help out both prior to and during the experimental trial.

Thanks also to Fonterra Co-operative Group Ltd for the financial backing to allow this trial to occur. Thanks to all those at Fonterra who contributed their time and effort in supporting this study, especially Angela Rowan (for her continued support for the study both in gaining funding and overseeing it for Fonterra), Arron Harford (for his work with the proteins used in this study) and Roxanne Brassington (for her work in the masking of the dietary beverages).

Thanks must also go to all the subjects that participated in this trial. Their willingness and outstanding commitment to this trial to ensure its success was greatly appreciated. It was great privilege to have such highly talented and committed subjects on this trial.

I would also like to thank my current employer, Universal College of Learning (UCOL), for their assistance in allowing me to pursue post-graduate study and to the people there who, over the years, supported me during my studies.

And finally, a very special thanks to my wife, Paula, and two children, Dominic and Abby, for all of their patience, support and understanding throughout the entire duration of this study. Without your support, completion of this would not have been possible.

*Approval for this research has been obtained from both the Massey University Ethics Committee and the Manawatu-Whanganui Ethics Committee for the experiments described in this thesis.*

# TABLE OF CONTENTS

Page Number

---

<b>ABSTRACT</b>	<b>ii</b>
<b>ACKNOWLEDGEMENTS</b>	<b>iv</b>
<b>TABLE OF CONTENTS</b>	<b>vi</b>
<b>LIST OF TABLES</b>	<b>ix</b>
<b>LIST OF FIGURES</b>	<b>xi</b>
<b>LIST OF ABBREVIATIONS</b>	<b>xvi</b>

---

<b>GENERAL INTRODUCTION</b>	<b>1</b>
<b>CHAPTER 1        LITERATURE REVIEW</b>	<b>3</b>
1.1 Introduction	3
1.2 Nutritional Strategies During Exercise	4
1.2.1 Fluids and Electrolytes	4
1.2.2 Carbohydrate	8
1.2.3 Protein	13
1.3 Nutritional Strategies Post-Exercise	18
1.3.1 Fluids and Electrolytes	19
1.3.2 Carbohydrates	21
1.3.3 Protein	26
1.3.3.1 Protein and the Insulin Response	27
1.3.3.2 Protein and Protein Synthesis / Degradation	35
1.4 Summary / Hypothesis	42
<b>CHAPTER 2        METHODS</b>	<b>43</b>
2.1. Subjects	43
2.1.1. Screening of Subjects	43
2.2. Experimental Design	44
2.2.1. Pre-Experimental Period	44
2.2.1.1. Determination of Peak Oxygen Uptake and Maximum Power Output	44
2.2.1.2. Determination of Body Composition	46
2.2.1.3. Sum of Skinfolds	47

2.2.1.4. Percentage Body Fat	47
2.2.1.5. Body Mass Index (BMI)	47
2.2.1.6. Waist : Hip Ratio	48
2.2.1.7. Somatotype	48
2.2.2. Experimental Design	49
2.2.2.1. Dietary Beverage Composition	49
2.2.2.1.1. Carbohydrate Content of the Dietary Beverages	51
2.2.2.1.2. Protein Content of the Dietary Beverages	51
2.2.2.1.3. Anion : Cation Balance	
2.2.2.1.4. Double-Blinding of Treatments and Administration of the Beverages	53
2.2.2.2. Experimental Period	55
2.2.2.2.1. Day 1 (Diet Record)	59
2.2.2.2.2. Day 2 & 3 (Controlled Diet)	59
2.2.2.2.3. Day 4 (Experimental Day)	60
2.2.2.2.3.1. Glycogen Depletion Protocol	60
2.2.2.2.4. In-Dwelling Cannulae and Blood Sampling	63
2.2.2.2.4.1. In-Dwelling Cannulae	63
2.2.2.2.4.2. Blood Sampling	64
2.2.2.2.5. Urine Sampling	65
2.2.2.2.6. Heart Rate Recording	66
2.2.2.2.7. Blood Pressure Measurement	66
2.2.2.2.8. Snack	67
2.3. Chemical Analysis	68
2.4. Statistical Analysis	69
<b>CHAPTER 3</b>	<b>RESULTS</b>
	<b>70</b>
3.1 Experimental Trial Overview	70
3.1.1 Exercise Protocol	70
3.1.2 Trial Beverage Consumption	70
3.1.3 Blood & Urine Sampling	
3.2 Anthropometric and Exercise Test ( $VO_{2peak}$ and $W_{max}$ ) Results	71
3.3 Variability Associated with Chemical Analyses	74
3.4 Amino Acid Composition of Diets	76
3.5 Metabolic and Physiological Responses	77
3.5.1 Rehydration	77
3.5.1.1 Haematocrit	77
3.5.1.2 Plasma Albumin	79
3.5.1.3 Urine Production	81
3.5.1.3.1 Analysis of Cumulative Urine Output Over Initial 240 Minutes	81
3.5.1.3.2 Temporal Pattern of Urine	82

3.5.1.3.3	Temporal Pattern of Urine Output Over Initial 240 Minutes	84
3.5.1.3.4	Temporal Pattern of Urine Output Over Full 420 Minutes	84
3.5.2	Carbohydrate Metabolism	85
3.5.3	Cardiovascular Recovery	89
3.5.3.1	Heart Rate	89
3.5.3.2	Systolic Blood Pressure	93
3.5.3.3	Diastolic Blood Pressure	94
3.5.3.4	Pulse Pressure	97
3.5.4	Protein Recovery	99
3.5.4.1	Urinary 3-methylhistidine	99
3.5.4.2	Urinary Urea	101
3.5.4.3	Plasma Urea	103
3.5.4.4	Urinary Ammonia	106
3.5.4.5	Urinary Creatinine	108
3.5.4.6	Plasma Essential Amino Acids	109
3.5.4.7	Plasma Non-Essential Amino Acids	116
3.5.4.8	Plasma Amino Acids Not Supplied in Beverages	119
<b>CHAPTER 4</b>	<b>DISCUSSION</b>	<b>121</b>
4.1	Rehydration	121
4.1.1	Plasma Albumin	121
4.1.2	Haematocrit	123
4.1.3	Urine Output	124
4.1.4	Sodium and Rehydration	125
4.1.5	Role of Potassium in Supporting Rehydration	127
4.1.6	Comparison of the Protein-Containing Beverages on Rehydration	128
4.2	Cardiovascular Recovery	129
4.3	Carbohydrate Recovery	131
4.4	Protein Recovery	134
4.4.1	Plasma Amino Acids	134
4.4.2	Plasma Urea Concentration and Urinary Excretion of Urea and Ammonia	136
4.4.3	Proposed Mechanism for Difference Between Intact Protein and Protein Hydrolysate Beverages on Protein Recovery	137
4.4.4	3-Methylhistidine	138
<b>CHAPTER 5</b>	<b>CONCLUSIONS &amp; GENERAL DISCUSSIONS</b>	<b>140</b>
<b>REFERENCES</b>		<b>144</b>

# LIST OF TABLES

<b>CHAPTER 1</b>	<b>LITERATURE REVIEW</b>	<b>Page Number</b>
<b>Table 1.1.</b>	Endogenous fuel sources in man, (Lambert <i>et al.</i> , 1997).	9
<b>Table 1.2.</b>	Summary of research examining the effect of the consumption of protein in addition to carbohydrate on insulin response and muscle glycogen resynthesis.	30
<b>Table 1.3.</b>	Amino acid composition of hydrolysates of whey, pea and wheat protein, and intact casein protein (van Loon <i>et al.</i> , 2000a).	34
<b>Table 1.4.</b>	Leucine and Branched Chain Amino Acid (BCAA) content of foods1 (Layman & Baum, 2004).	40
<b>CHAPTER 2</b>	<b>METHODS</b>	<b>Page Number</b>
<b>Table 2.1.</b>	List of proposed performance enhancing substances that have the potential to influence a subject's carbohydrate and / or protein metabolism if used prior to or during a post-exercise recovery study.	44
<b>Table 2.2.</b>	Body composition assessment sites used in a post-exercise recovery study.	46
<b>Table 2.3.</b>	Ingredient composition of the four dietary beverages powders used in a post-exercise recovery study.	50
<b>Table 2.4.</b>	Expected intake rates for both of the dietary beverage powder (grams of dry matter (DM)) and water, expressed per kg of body weight, used in a post-exercise recovery study.	50
<b>Table 2.5.</b>	Typical amino acid composition of ALACEN <sup>TM</sup> 895 Whey Protein Isolate* used in a post-exercise recovery study.	52
<b>Table 2.6.</b>	Mineral composition* of the whey protein isolate and whey protein hydrolysate used in a post-exercise recovery study.	53
<b>Table 2.7.</b>	Initial and Final Dietary Electrolyte Balance (DEB) + for the three protein containing beverages tested in a post-exercise recovery study.	54
<b>Table 2.8.</b>	Randomised allocation of the four beverages tested used in a post-exercise recovery study designed so that each beverage does not routinely follow the same combination in each subject.	58
<b>Table 2.9.</b>	Diet composition analysis of the controlled diet consumed the two days prior to each experimental day in a post-exercise recovery study.	59
<b>Table 2.10.</b>	Desired water consumption throughout the experimental day in a post-exercise recovery study.	62
<b>Table 2.11.</b>	Diagnostic kits used for chemical analysis performed in a post-exercise recovery study.	68

<b>CHAPTER 3 RESULTS</b>		<b>Page Number</b>
<b>Table 3.1.</b>	Summary of the data gained from anthropometric assessment of the ten male subjects that were involved in a post-exercise recovery study.	72
<b>Table 3.2.</b>	Data from the pre-trial fitness assessment for the 11 males scheduled to be involved in a post-exercise recovery study.	73
<b>Table 3.3.</b>	Variability (CV%) associated with various chemical analyses conducted on samples collected during a post-exercise recovery study.	74
<b>Table 3.4.</b>	Variability (CV%) associated with the determination of amino acids in the beverages used in a post-exercise recovery study.	75
<b>Table 3.5.</b>	Determined amino acid composition of the four dietary beverages, expressed as g/kg BWT delivered, in a post-exercise recovery study.	76
<b>Table 3.6.</b>	The mean and standard error for haematocrit (%) of all athletes for the four beverages during the time period from cessation of exercise to 240 minutes post-exercise during a post-exercise recovery study.	78
<b>Table 3.7.</b>	The mean and standard error of plasma albumin concentration (g/L) for the response to the four beverages during the time period from cessation of exercise to 240 minutes post-exercise in a post-exercise recovery study	80
<b>Table 3.8.</b>	The mean and standard error of urine output results (g/30minutes) for the response to the four beverages during the time period from cessation of exercise to 420 minutes post-exercise in a post-exercise recovery study.	82
<b>Table 3.9.</b>	Desired and actual water consumption achieved throughout the experimental day in a post-exercise recovery study.	83
<b>Table 3.10.</b>	The mean and standard error results of plasma insulin concentration for the response to the four beverages during the time period from cessation of exercise to 240 minutes post-exercise in a post-exercise recovery study.	87
<b>Table 3.11.</b>	The mean and standard error results of plasma glucose concentration for the response to the four beverages during the time period from cessation of exercise to 240 minutes post-exercise in a post-exercise recovery study.	87
<b>Table 3.12.</b>	Linear Regression results for heart rate during the post-exercise recovery period in a post-exercise recovery study.	91
<b>Table 3.13.</b>	Analysis results of slope coefficients for post-exercise heart rate recovery for the response to four dietary beverages during a post-exercise recovery study.	92
<b>Table 3.14</b>	Statistical analysis of the effect of dietary beverages on individual mean plasma essential amino acid levels, by pairwise repeated measures ANOVA, during the time period from cessation of exercise to 240 minutes post-exercise during a post-exercise recovery study.	115
<b>Table 3.15</b>	Effect of dietary beverages on individual mean plasma non-essential amino acid levels, by pairwise repeated measures ANOVA, during the time period from cessation of exercise to 240 minutes post-exercise during a post-exercise recovery study.	118

# LIST OF FIGURES

<b>CHAPTER 1 LITERATURE REVIEW</b>		<b>Page Number</b>
<b>Figure 1.1.</b>	Estimated balance of Na <sup>+</sup> intake and Na <sup>+</sup> loss during 3 h of exercise at marathon pace. <i>Note:</i> 1 mmol of NaCl weighs 58.5 mg (Lambert, <i>et al.</i> , 1997).	7
<b>Figure 1.2.</b>	Summary of the main pathways of energy metabolism using carbohydrate, lipid, and protein as energy sources (Maughan <i>et al.</i> , 1997, p16).	11
<b>Figure 1.3.</b>	Amino acids are degraded to pyruvate, acetate, or to intermediates of the TCA cycle (Maughan <i>et al.</i> , 1997, p124).	14
<b>Figure 1.4.</b>	Amino acid metabolism in the muscle. (Wagenmakers, 2000, p120)..	14
<b>Figure 1.5.</b>	Interorgan relationship in the handling of amino acids. Dashed arrow, prolonged starvation only. (Wagenmakers, 2000, p122)..	16
<b>Figure 1.6.</b>	Muscle glycogen synthesis rates are depicted against the rate of carbohydrate ingestion. The horizontal line depicts the absolute maximum for muscle glycogen synthesis (Jentjens & Jeukendrup, 2003).	25
<b>Figure 1.7.</b>	Rates of leg protein dynamics (a) and whole body protein dynamics (b) for 10 subjects given an oral nutrient supplementation either immediately after exercise (EARLY) or 3 h postexercise (LATE). * Significantly different (p<0.05), EARLY vs LATE (Levenhagen <i>et al.</i> , 2001).	37
<b>Figure 1.8.</b>	Insulin signalling cascade. GLUT1, insulin independent glucose transporter; PKC, protein kinase C; eIF, translational initiation factors. (Layman & Baum, 2004).	40
<b>CHAPTER 2 METHODS</b>		<b>Page Number</b>
<b>Figure 2.1.</b>	Overview of a seven day experimental period highlighting division of the two phases for a post-exercise recovery study.	56
<b>Figure 2.2.</b>	Schematic overview of the Day 4 (Experimental Day) for a post-exercise recovery study.	57
<b>Figure 2.3.</b>	Glycogen depletion protocol used in a post-exercise recovery.	61

CHAPTER 3 RESULTS		Page Number
<b>Figure 3.1.</b>	Example graph of VO <sub>2</sub> output from a pre-experimental trial fitness test. The five-second interval for determining VO <sub>2peak</sub> is highlighted on the right hand side of the graph. The result for this athlete was recorded at 4.73 L/min or 71.77 ml/kg/min.	73
<b>Figure 3.2.</b>	Mean (± SEM) haematocrit (%) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 240 minute period following exercise and recovery beverage consumption.	78
<b>Figure 3.3.</b>	Mean (± SEM) plasma albumin concentrations (g/L) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 240 minute period following exercise and recovery beverage consumption.	80
<b>Figure 3.4.</b>	Mean (± SEM) total urine volumes expressed as ml/kg BWT for the first 240 minutes (fluid intake controlled during this period) in a post-exercise recovery study. <i>Similar letter designations implies non-significance, non-similar implies significance (p&lt;0.05).</i>	81
<b>Figure 3.5.</b>	Mean (± SEM) urine outputs (g/30minutes) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	83
<b>Figure 3.6.</b>	Mean (± SEM) plasma insulin concentrations (mmol/L) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 240 minute period following exercise and recovery beverage consumption.	88
<b>Figure 3.7.</b>	Mean (± SEM) plasma glucose concentrations (mmol/L) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 240 minute period following exercise and recovery beverage consumption.	88
<b>Figure 3.8.</b>	Linear regression analysis results of mean post-exercise pulse rate recovery (bpm) (n=4900) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H), intact whey protein (Carb + I), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M) in a 240 minute period following exercise and recovery beverage consumption.	89

<b>Figure 3.9.</b>	Linear regression analysis results of mean post-exercise pulse rate recovery (bpm) following consumption of each of the four dietary beverages (n=4900): (a) Carbohydrate Only; (b) Carbohydrate and Intact Protein; (c) Carbohydrate / Protein Hydrolysate; (d) Carbohydrate and 50% whey protein hydrolysate and 50% intact whey protein, in a 240 minute period following exercise and recovery beverage consumption.	90
<b>Figure 3.10.</b>	Mean ( $\pm$ SEM) systolic blood pressures (mmHg) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	96
<b>Figure 3.11.</b>	Mean ( $\pm$ SEM) diastolic blood pressures (mmHg) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	96
<b>Figure 3.12.</b>	Mean ( $\pm$ SEM) pulse pressure values (mmHg) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	98
<b>Figure 3.13.</b>	Mean ( $\pm$ SEM) urinary 3-methylhistidine excretion (mg) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	99
<b>Figure 3.14.</b>	Mean ( $\pm$ SEM) urinary urea nitrogen excretion (mg) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	105
<b>Figure 3.15.</b>	Mean ( $\pm$ SEM) plasma urea concentration (mmol/L) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	105
<b>Figure 3.16.</b>	Mean ( $\pm$ SEM) urinary ammonia nitrogen excretion (mg) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	107

<b>Figure 3.17.</b>	Mean ( $\pm$ SEM) total urinary creatinine nitrogen excretion (mg) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	108
<b>Figure 3.18.</b>	Mean ( $\pm$ SEM) plasma Valine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	111
<b>Figure 3.19.</b>	Mean ( $\pm$ SEM) plasma Methionine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	111
<b>Figure 3.20.</b>	Mean ( $\pm$ SEM) plasma Leucine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	112
<b>Figure 3.21.</b>	Mean ( $\pm$ SEM) plasma Isoleucine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	112
<b>Figure 3.22.</b>	Mean ( $\pm$ SEM) plasma Phenylalanine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	113
<b>Figure 3.23.</b>	Mean ( $\pm$ SEM) plasma Tryptophan concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	113
<b>Figure 3.24.</b>	Mean ( $\pm$ SEM) plasma Lysine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	114

<b>Figure 3.25.</b>	Mean ( $\pm$ SEM) plasma Arginine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	117
<b>Figure 3.26.</b>	Mean ( $\pm$ SEM) plasma Tyrosine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	117
<b>Figure 3.27.</b>	Mean ( $\pm$ SEM) plasma ornithine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	120
<b>Figure 3.28.</b>	Mean ( $\pm$ SEM) plasma 3-methylhistidine concentration (nmol/ml) after ingestion of a control drink containing carbohydrate only (Carb; n=10) and drinks containing carbohydrate plus either: whey protein hydrolysate (Carb + H; n=10), intact whey protein (Carb + I; n=11), or 50% whey protein hydrolysate and 50% intact whey protein (Carb + M; n=11) in a 420 minute period following exercise and recovery beverage consumption.	120

---

## LIST OF ABBREVIATIONS

Carb	Carbohydrate Only
Carb + H	Carbohydrate + Whey Protein Hydrolysate
Carb + I	Carbohydrate + Intact Whey Protein
Carb + M	Carbohydrate + 50% Whey Protein Hydrolysate + 50% Intact Whey Protein
ACSM	American College of Sports Medicine
ADA	American Dietetic Association
ANP	Atrial Natriuretic Peptide
BWT	Body Weight
DEB	Dietary Electrolyte Balance
DM	Dry Matter
DOC	Dietitians of Canada
GLUT	Glucose Transporter Carrier Proteins
FAO	Food and Agriculture Organization of the United Nations
ISAK	International Society for the Advancement of Kinanthropometry
TCA	Tricarboxylic Acid
UNU	United Nations University
VO <sub>2peak</sub>	Peak Oxygen Uptake
W <sub>max</sub>	Maximum Power Output
WHO	World Health Organisation

## GENERAL INTRODUCTION

Athletes around the world continually strive for that elusive edge that will give them victory over their fellow competitors. This search for the perfect performance is epitomised by the foundations of the modern Olympic Games and Olympism. The International Olympic Committee defines Olympism as...

*"... a philosophy of life, exalting and combining in a balanced whole the qualities of body, will and mind. Blending sport with culture and education, Olympism seeks to create a way of life based on the joy found in effort, the educational value of good example and respect for universal fundamental ethical principles." (International Olympic Committee, 2004, p.9)*

An athlete's performance is influenced by a multitude of factors, ranging from their genetic makeup and their physical fitness, through to motivational levels and the desire to succeed. Nutrition is another such factor which, while a relatively new science in terms of sports nutrition, now has a strong focus within the sport science research community. With the commercialism of sport, athletes have a lot to gain from the spoils of victory, ranging from the personal satisfaction gained years of training and effort, through to financial rewards. This has led to the development of a 'win at all costs' attitude, which has often taken athletes down the path of performance enhancing ergogenic aids, many of which are banned by the International Olympic Committee and the World Anti-Doping Agency. With nutrition, however, there are many dietary factors that have the potential to provide the athlete with an ergogenic, or performance-enhancing effect. Dietary protein definitely falls into this category, and its potential has recently been further enhanced through the development of manufacturing processes which produce protein forms that are well suited to an athlete's physiological needs. While protein intake is traditionally associated with strength-based athletes, it is now evident that the protein requirement of endurance athletes is also increased due to a combination of exercise-induced muscle damage and the use of protein as a fuel source.

Throughout history, athletes have trialled nutritional strategies, focused on dietary protein, that have resulted in both in enhancement and detriment of athletic performance. One of the most famous accounts of food consumption relates to Milos, a five-time Olympic Cretan wrestler champion (552-516 B.C.), who had a reported daily meat intake of 10kg per day (Maughan & Burke, 2000). Following the 1936 Berlin Olympics, it was stated that athletes frequently focused on the intake of meat and on average took in nearly half a kilogram of meat per day, with pre-event meals consisting of steak and eggs, supplemented with meat-juice (Schenk, 1936, cited in McArdle, Katch & Katch, 1999). This large consumption of protein by athletes, in the form of meat, has more recently been replaced with specially formulated protein supplements, which provide the athlete with a more convenient and healthier option.

Dietary protein supplementation is now common place both amongst athletes and the wider population, with a focus related to both exercise and weight loss. It also represents a major part of a world-wide, multi-million dollar supplement industry. In relation to athletes, there is a common belief that protein consumption has significant benefits on both performance and recovery from exercise. Yet uncertainty still remains with regard to the best form of dietary protein to use and when is the optimal time for its consumption (generally related to an exercise session). To date, nutritional research has been focused around both pre-exercise and during exercise nutritional strategies for maximising performance. However, due to the demands of modern sport, nutritional strategies focused on maximising recovery from exercise are of increased importance. It is during this post-exercise recovery period that the true benefits of protein consumption are becoming more evident. The potential benefit of protein to the recovering athlete relates to achieving maximal training adaptations, in addition to ensuring that exercise performance is not impaired due to any lingering effects from a previous exercise session.

There is still a lot to learn about the true extent that dietary protein intake may have on the recovering athlete. This study looks to extend the current knowledge related to the effect of dietary protein in exercise recovery, with a focus on the use of different forms of whey protein (intact and hydrolysed) on post-exercise recovery of endurance athletes.