

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

# **Development of a Mathematical Model for 'Hayward' Kiwifruit Softening in the Supply Chain**

A thesis presented in partial fulfilment of the requirements for the  
degree of Doctor of Philosophy in Food Technology at Massey  
University, New Zealand

Zhao Junyu, Matthew

2017



## **Abstract**

Fruit loss is a major concern to the kiwifruit industry as it incurs a high cost to monitor and remove over soft or rotten fruit to meet export standards. Kiwifruit is exposed to various temperature scenarios due to different packhouse cooling practices, and temperature control is difficult to maintain throughout the supply chain. Fruit pallet temperatures are wirelessly monitored in the supply chain. This time temperature data provides valuable rich information which could be used to predict kiwifruit quality.

In the laboratory, green 'Hayward' kiwifruit were exposed to industry coolchain scenarios to investigate their influence on fruit firmness in subsequent storage. Cooling rate and storage temperature were identified to affect fruit firmness and chilling injury development significantly, where accelerated softening and increased chilling injury development was observed in late storage (> 100 d) when fruit were cooled directly to 0 °C. However, when fast cooled fruit were stored at 2 °C instead of 0 °C, low incidence of chilling injury was observed. The influence of cooling rate and storage temperature on kiwifruit quality suggests that industry should focus on the management practices adopted by packhouses in order to maintain acceptable quality after long term storage. A proportion of the firmness data results were used to develop a mechanistic style mathematical model of kiwifruit softening. Kiwifruit softening was mathematically described based on the correlation with starch degradation, breakdown of cell wall structure, and a description of the incidence of chilling injury development during storage. The model inputs consist of solely commonly collected at-harvest attributes: firmness, dry matter and soluble solids content and time-temperature data. Applying at-harvest attributes as model inputs enabled a capability to predict different softening curves as influenced by fruit maturity, and grower line differences. The developed model

demonstrated promising softening prediction with mean absolute errors (MAE) between 0.8 to 2.1 N when fruit were exposed to fluctuating temperatures and cooling profiles. A logistic model was used to estimate the proportion of chilling injured fruit. Based on the given time temperature information, the logistic model was able to predict the proportion of chilling injured fruit reasonably well ( $R^2 = 0.735$ ). This chilling injury prediction was subsequently used to adjust the softening prediction during the late storage period (>100 d). Model validation was performed using the remaining data, identifying a lack of fit in both the rapid (MAE of 20.8 N) and gradual (MAE of 8.0 N) softening phase. The lack of fit in the rapid softening phase is proposed to be explained by the presence of an initial lag phase in softening which the developed model is unable to predict. The magnitude of firmness associated with starch content and cell wall integrity heavily influenced the lack of fit in the gradual softening phase. Fixing the initial amount of firmness associated to cell wall integrity to be constant for all maturities and grower lines improved the softening prediction.

Overall, this thesis contributes to the challenge of predictively modelling kiwifruit quality in the supply chain. However, there are still many opportunities for improvement including introducing the influence of: variation within the same batch; fruit maturity on chilling injury development; ethylene in the environment; pre-harvest management practices and extending the model to have more focus on high temperature conditions such as those experienced in the marketplace. Conducting studies on: the effect of curing on kiwifruit; using non-destructive techniques to provide information to help define model parameters for prediction; effect of high temperature exposure on kiwifruit softening are possible opportunities that may contribute to enable better prediction of kiwifruit quality in the supply chain in the future.

## **Acknowledgments**

I wish to thank my supervisors, Dr. Andrew East and Prof. John Bronlund for the incredible amount of time and support given to me during my PhD. I greatly appreciate their positive feedback, continuous efforts and always encouraging me through tough times.

I would like to say many thanks to Peter Jeffery for his help when I had troubles in either my laboratory work or technical issue with computer and Sue Nicholson for her help and patience in assisting me in experimental planning and during my time in the laboratory. I am also thankful to Byron, Warwick and Gary for their technical support.

Heaps of thanks should go to Zespri International Ltd for the fellowship and supplying kiwifruit for this study. I am also thankful to Massey University for providing extra funding to relief my financial issues especially during my last year of study.

I am indebted to many of my friends and colleagues for their help and support throughout my studies in New Zealand. They are Tamarath (An) Pranamornkith, Gayani, Khairul, Pilirani, Jantana, Li Mo, Abdul and Srikanth. Special thanks to Ivana, Wensheng, Grace, Eileen, Seng Guan and Jordan for being an asset of good friends during my time in New Zealand.

Special thanks to my mother and father for their encouragement and appreciate my siblings (Roy, Vyennus and Dominic) for taking care of the family when I am away for my studies. Finally, I would like to thank Jenny, who is a very special person of my life. I am grateful for her continuous encouragement and support during all tough stages in my studies and always there to listen to my problems. I appreciated all the time you spent waiting for me to return to Singapore.

## Table of contents

Abstract .....	i
Acknowledgments.....	iii
Table of contents.....	iv
List of Figures .....	x
List of Tables .....	xv
Nomenclature .....	xvii
1. Introduction.....	1
1.1. Thesis overview.....	3
2. Literature review.....	5
2.1. Kiwifruit.....	5
2.2. Kiwifruit ripening.....	5
2.2.1. Starch degradation.....	6
2.2.2. Breakdown of cell wall structure .....	7
2.2.2.1. Modification of cell wall polymeric network.....	8
2.2.2.2. Swelling of cell wall.....	10
2.2.3. Chilling injury development .....	11
2.3. Factors affecting fruit ripening.....	13
2.3.1. Temperature .....	13
2.3.2. Ethylene .....	14
2.3.3. Humidity .....	17

2.3.4.	Fruit maturity .....	18
2.3.5.	Rot development in kiwifruit .....	19
2.4.	Kiwifruit supply chain.....	20
2.4.1.	Pre-harvest treatments .....	20
2.4.2.	Postharvest treatment .....	21
2.5.	Softening models .....	25
2.5.1.	Empirical approach .....	25
2.5.2.	Mechanistic approach.....	26
2.5.2.1.	Temperature dependency.....	26
2.5.2.2.	Biological variability .....	27
2.6.	Summary .....	28
3.	Softening of ‘Hayward’ kiwifruit in different coolchain scenarios (*) .....	29
3.1.	Introduction .....	29
3.2.	Material and methods .....	31
3.2.1.	Supply chain simulation.....	31
3.2.2.	Fruit assessment .....	34
3.2.2.1.	Fruit firmness .....	34
3.2.2.2.	Soluble solids content.....	35
3.2.2.3.	Decay incidence .....	37
3.2.2.4.	Respiration rate.....	37
3.2.2.5.	Statistical analysis .....	37



3.3.	Result and discussions.....	38
3.3.1.	Cooling profiles achieved .....	38
3.3.2.	At-harvest attributes .....	41
3.3.3.	Effect of cooling rate on kiwifruit quality during storage .....	42
3.3.4.	Effect of storage temperature on kiwifruit quality during storage.....	48
3.3.5.	Effect of break in temperature control on kiwifruit quality during storage	50
3.3.6.	Effect of high temperature conditions on kiwifruit quality during storage	52
3.3.7.	Effect of storage humidity on kiwifruit quality during storage .....	54
3.4.	Conclusion.....	56
4.	Factors influencing development of chilling injury in ‘Hayward’ kiwifruit during coolstorage (*)......	57
4.1.	Introduction .....	57
4.2.	Materials and methods.....	59
4.2.1.	Fruit source .....	59
4.2.2.	Coolchain simulation .....	60
4.2.3.	Fruit assessment .....	62
4.3.	Results .....	65
4.3.1.	Cooling profiles achieved .....	65
4.3.2.	At-harvest fruit attributes .....	67
4.3.3.	Chilling injury development during storage .....	69

4.3.4.	Fruit firmness during storage .....	73
4.4.	Discussion .....	76
4.4.1.	Fruit maturity and grower line differences on kiwifruit quality .....	76
4.4.2.	Influence of chilling injury on firmness measurement .....	80
4.4.3.	Coolchain temperature effect on kiwifruit quality .....	82
4.5.	Conclusion.....	87
5.	Mathematical modelling of ‘Hayward’ kiwifruit softening.....	89
5.1.	Introduction .....	89
5.2.	Model conceptual framework.....	92
5.2.1.	Breakdown of starch content.....	95
5.2.2.	Breakdown of cell wall structure .....	97
5.2.3.	Development of chilling injury .....	98
5.3.	Model development .....	99
5.3.1.	Model constraints .....	99
5.3.2.	Mathematical formulation.....	100
5.3.3.	Estimation of initial starch content ( $S_0$ ) .....	102
5.3.4.	Model of starch breakdown effect on firmness ( $F_A$ ) .....	102
5.3.5.	Model of the breakdown of cell wall structure on firmness ( $F_B$ ).....	107
5.3.6.	Model for the development of chilling in jury effect on firmness .....	113
5.3.7.	Prediction of kiwifruit firmness ( $F_{pred}$ ) .....	116
5.4.	Model summary.....	117

5.5.	Conclusion.....	120
6.	Developed model performance.....	121
6.1.	Introduction .....	121
6.2.	Materials and methods.....	122
6.2.1.	Grower difference .....	123
6.2.2.	Cooling rate effect on fruit firmness .....	125
6.2.3.	Storage temperature effect on fruit firmness.....	127
6.2.4.	Break in temperature control effect on fruit firmness.....	129
6.2.5.	Effect of exposure to high temperature on fruit firmness .....	131
6.2.6.	Prediction on the proportion of chilling injured fruit.....	132
6.3.	Model error and sensitivity.....	133
6.3.1.	Model inputs .....	134
6.3.2.	Global model parameters .....	140
6.4.	Overall discussion .....	142
6.5.	Conclusion.....	144
7.	Validation of ‘Hayward’ kiwifruit softening model in coolstorage .....	146
7.1.	Introduction .....	146
7.2.	Material and methods .....	147
7.2.1.	Predictive modelling .....	149
7.3.	Results and discussion.....	149
7.3.1.	Fruit maturity difference .....	149

7.3.2.	Fruit grower difference .....	156
7.3.3.	Prediction of chilling injured fruit.....	163
7.4.	Overall discussion .....	165
7.5.	Conclusion.....	167
8.	Overall discussion and recommendations.....	168
8.1.	Introduction .....	168
8.2.	Establishment of model .....	169
8.3.	Model softening prediction .....	170
8.3.1.	Prediction of lag phase .....	171
8.3.2.	Prediction of gradual softening phase ( $F_{B0}$ ).....	178
8.3.3.	Prediction of incidence of chilling injury ( $CI$ ).....	181
8.4.	At-harvest attributes as model inputs .....	183
8.5.	Application to industry .....	184
8.6.	Possible future opportunities .....	185
8.6.1.	Curing of kiwifruit .....	185
8.6.2.	Pre-harvest effect on fruit storability .....	186
8.6.3.	Predict fruit firmness with biological variation .....	187
8.6.4.	High temperature exposure .....	189
8.7.	Conclusion.....	190
9.	References.....	191
	Appendices.....	224

## List of Figures

Figure 2.1: Schematic representation of key events .....	6
Figure 3.1: Example of time temperature profile of the air inside the box.....	29
Figure 3.2: Time temperature profile of direct (A) and gradual (B) cooling.....	33
Figure 3.3: Description of various temperature scenarios used for ‘Hayward’ kiwifruit from 2012 harvest .....	36
Figure 3.4: Fruit rots found at the side or stem end of ‘Hayward’ kiwifruit. Red arrow indicates the rotten area.....	37
Figure 3.5: Room and tray air time temperature for direct (A) and gradual (B) cooling to 0 or 2 °C.....	39
Figure 3.6: Real time temperature profile of the different temperature scenarios simulated in the laboratory. ....	40
Figure 3.7: Effect of cooling profile on ‘Hayward’ kiwifruit softening at constant storage conditions of 0 (A) and 2 (B) °C.....	43
Figure 3.8: Effect of cooling profile on the accumulation of soluble solids in ‘Hayward’ kiwifruit at control storage conditions of 0 (A) and 2 (B) °C.....	45
Figure 3.9: Effect of cooling rate on ‘Hayward’ kiwifruit respiration rate at 0 (A) and 2 (B) °C.....	46
Figure 3.10: Effect of cooling rate on the incidence of rotten fruit in ‘Hayward’ kiwifruit subsequently stored at 0 or 2 °C. ....	48
Figure 3.11: Effect of storage temperature on ‘Hayward’ kiwifruit softening when direct cooled (A) or gradual cooled (B) to storage temperature at 0 or 2 °C.....	49
Figure 3.12: Effect of break in temperature control on the kiwifruit firmness after 9 or 15 weeks of storage.....	51

Figure 3.13: Effect of high temperature exposure on kiwifruit firmness after 10 weeks of storage at 0 °C.....53

Figure 3.14: Effect of humidity at 30 °C on kiwifruit firmness after 14 weeks of storage at 0 °C.....55

Figure 4.1: The proposed cooling profiles to cool fruit to storage temperature .....61

Figure 4.2: Example of severity of chilling injury found along the outer pericarp of ‘Hayward’ kiwifruit. ....63

Figure 4.3: The comparison between a fruit with chilling injury symptom (left) and a rotten fruit (right). ....64

Figure 4.4: Cooling profiles designed and created for 6 different coolchain scenarios..66

Figure 4.5: Incidence of chilling injured fruit as influence by storage time, storage conditions and fruit maturity.....72

Figure 4.6: Softening of kiwifruit harvested at different maturity as influenced by supply chain temperature conditions. ....75

Figure 4.7: Softening of kiwifruit harvested at different maturity during storage in six different supply chain conditions.....78

Figure 4.8: Softening of kiwifruit harvested at different maturity during storage in six different supply chain conditions.....79

Figure 4.9: Histograms of fruit firmness as influenced by grower line (G1, G2 and G3) and maturity .....81

Figure 4.10: Time temperature profile of control and various “switch” treatments. 0 refers to fruit stored at 0 °C.....85

Figure 4.11: (A & B) Softening of kiwifruit exposed to different temperature “switch” treatments. ....86

Figure 4.12: Percentage of fruit with chilling injury after 125, 150, 175 and 200 d of storage .....	87
Figure 5.1: Average kiwifruit firmness and soluble solids content during storage at 0 °C. ....	94
Figure 5.2: The overall conceptual model development process.....	102
Figure 5.3: (A) Modelled accumulation of soluble solids content for 2012 season, ....	105
Figure 5.4: Correlation between fruit firmness and starch content in the first 50 days of cool storage. ....	106
Figure 5.5: Estimation of average initial firmness contribution of cell wall breakdown based on at-harvest attributes.....	109
Figure 5.6: The modelled change in firmness due to breakdown of cell wall structure (%) at various temperatures. ....	112
Figure 5.7: (A) Fitted model with the incidence of chilling injured fruit collected in the 2013 harvest season. ....	114
Figure 5.8: Logistic model used to describe the incidence of chilling injured fruit .....	116
Figure 5.9: Summary of the firmness prediction model developed.....	118
Figure 5.10: Simplified overall equations explaining the variables needed to predict fruit firmness in coolchain. ....	119
Figure 6.1: The model prediction (solid lines) against the experimental data.....	124
Figure 6.2: Experimental data points and modelled average fruit firmness (solid lines) during storage.....	126
Figure 6.3: Experimental data points and modelled average firmness (solid lines) of fruit that were cooled either directly or gradually to storage temperature.....	128
Figure 6.4: Experimental data points and modelled average firmness (solid lines) of fruit exposed to a break in temperature control .....	130

Figure 6.5: Experimental data and modelled average firmness of fruit exposed to high temperature conditions after 10 weeks of storage..... 132

Figure 6.6: Experimental data points and modelled average firmness of fruit with chilling injury development..... 133

Figure 6.7: Scatter plots of initial fruit firmness ( $F_0$ ), soluble solids ( $B_0$ ) and dry matter content ( $D_{m0}$ ) ..... 135

Figure 6.8: Sensitivity analysis of model inputs on predicting the softening curve. .... 136

Figure 6.9: Scatter plots of initial starch content ( $S_0$ ), soluble solids ( $B_0$ ) and dry matter content ( $D_{m0}$ ) ..... 137

Figure 6.10: Sensitivity analysis of model inputs on predicting the softening curve. ... 139

Figure 6.11: Sensitivity analysis of model inputs on predicting the softening curve. ... 140

Figure 6.12: Sensitivity analysis of model parameter ( $a$ ) on predicting the softening curve ..... 141

Figure 6.13: Sensitivity analysis of model parameter ( $k_w$ ) on predicting the softening curve of the same grower line. .... 142

Figure 6.14: The developed model on predicting the softening of ‘Hayward’ kiwifruit ..... 143

Figure 7.1: Correlation between fruit firmness and starch content during storage period. .... 150

Figure 7.2: The linear correlation of fruit firmness and starch content is invalid when starch content is above 8 %. ..... 151

Figure 7.3: The predicted softening curve of fruit using different fruit maturity at-harvest attributes..... 153

Figure 7.4: Model validation on the experimental data collected in 2013..... 155



Figure 7.5: Difference between firmness prediction ( $F_{pred}$ ) when adopting the modified correlation between starch and fruit firmness. .... 156

Figure 7.6: The predicted softening curve of early maturity fruit from different grower lines using at-harvest attributes ..... 158

Figure 7.7: Model validation on softening pattern of late maturity fruit from three different grower lines collected in the 2013 season. .... 159

Figure 7.8: Comparison of softening prediction using at-harvest attributes of respective grower lines or an average across 3 grower lines ..... 161

Figure 7.9: Model validation on softening pattern of late maturity fruit from three different grower lines ..... 162

Figure 7.10: Comparison of the predicted and experimental data for the proportion of chilling injured fruit ..... 164

Figure 7.11: Comparison on the predicted fruit softening curve of rapidly cooled ( $R_{12h,0}$ ) against gradually cooled ( $G_{2w,0}$ ) fruit to 0 °C. .... 165

Figure 8.1: (A) Correlation of fruit firmness and starch content with a plateau when starch content is above 8 %. .... 171

Figure 8.2: The accumulation of soluble solids content ..... 174

Figure 8.3: The model prediction on the softening curve by modifying the rate of starch degradation ( $k_s$ ) to be 3 times lower when starch content is above 8 %. .... 176

Figure 8.4: Model prediction of firmness of fruit in 2012 (A) and 2013 (C) harvest season. .... 179

Figure 8.5: The possible approach to relate ethylene to fruit firmness..... 181

## List of Tables

Table 3.1: Average at-harvest attributes of kiwifruit from respective grower lines. ....	41
Table 3.2: Contingency table for the relationship of cooling rate on the incidence of rotten fruit in ‘Hayward’ kiwifruit .....	47
Table 3.3: Contingency table for the relationship of storage temperature on the incidence of rotten fruit in ‘Hayward’ kiwifruit.....	50
Table 4.1: Average at-harvest attributes of kiwifruit from 3 grower lines harvested at 3 different maturity stages.....	68
Table 4.2: Contingency table for the relationship of storage treatment on incidence of chilling injured fruit after 172 d of storage. ....	70
Table 4.3: Contingency table for the relationship of fruit maturity on incidence of chilling injured fruit after 172 d of storage. ....	71
Table 4.4: Contingency table for the relationship of grower line on incidence of chilling injured fruit after 172 d of storage. ....	72
Table 4.5: ANOVA table displaying the sum of square (SS), mean square (MS), F and P value of respective factors.....	73
Table 5.1: Summary of different mechanisms hypothesized to occur during the observed phases of kiwifruit softening.....	93
Table 5.2: The values of the model parameters with the standard deviation.....	104
Table 5.3: Values of the model parameters with the standard deviation .....	111
Table 5.4: Estimated values of activation energies for different reactions .....	112
Table 6.1: At-harvest attributes of kiwifruit from respective grower lines from 2012 season. ....	122
Table 7.1: The model inputs and parameters used to predict fruit firmness in coolchain. ....	148

Table 7.2: Average At-harvest attributes of kiwifruit from 3 growers harvested at 3 different maturity stages..... 148

Table 7.3: Average at-harvest attributes of kiwifruit from 3 different maturity stages.153

Table 7.4: The different approach to define model parameter  $a$  and  $F_{B0}$ ..... 161

Table 7.5: The comparison of the proportion of chilling injured fruit (%) between model prediction (Pred) and experimental data (Exp)..... 163

## Nomenclature

Symbol	Definition	Units
$a$	Correlation between starch and firmness	$N \%^{-1}$
$AHU_a$	AHU when 50 % of chilling injured fruit is reached	$^{\circ}C d$
$AHU$	Accumulated heat unit	$^{\circ}C d$
$B$	Soluble solid contents	$^{\circ}Brix$
$B_0$	Initial soluble solid content	$^{\circ}Brix$
$B_{final}$	Final soluble solids content	$^{\circ}Brix$
$F_{CI}$	Lowest firmness measured	$N$
$CI$	Proportion of chilling injured fruit	$\%$
$CI_{max}$	Maximum proportion of chilling injured fruit	$\%$
$CI_{min}$	Minimum proportion of chilling injured fruit	$\%$
$D_m$	Dry matter content	$\%$
$E_{a,s}$	Activation energy for starch breakdown	$J mol^{-1}$
$E_{a,p}$	Activation energy for breakdown of cell wall structure	$J mol^{-1}$
$F$	Fruit firmness	$N$
$F_0$	Initial fruit firmness	$N$
$F_A$	Firmness correlated with starch degradation	$N$
$F_B$	Firmness contributed by cell wall component	$N$
$F_{Fix}$	Underlying basal firmness	$N$
$F_{pred}$	Predicted fruit firmness	$N$
$F_{soft}$	Predicted soft fruit firmness	$N$
$k_b$	Rate constant of accumulation of soluble solids content	$d^{-1}$
$k_s$	Rate constant of starch breakdown	$d^{-1}$
$k_{s,ref}$	Rate constant of starch breakdown at $T_{ref}$	$d^{-1}$
$k_w$	Rate constant of the breakdown of cell wall structure	$d^{-1}$
$k_{w,ref}$	Rate constant of the breakdown of cell wall structure at $T_{ref}$	$d^{-1}$
$R$	Universal gas constant, 8.314	$J mol^{-1}K^{-1}$
$S$	Starch content	$\%$
$S_0$	Initial starch content	$\%$
$t$	Time	$d$
$T$	Temperature	$^{\circ}C$
$T_{abs}$	Temperature in absolute	$K$
$T_{ref}$	Reference temperature at 20 $^{\circ}C$	$K$
$T_b$	Base temperature	$^{\circ}C$
$\mu$	Rate constant of chilling injury development	$d^{-1}$

<b>Symbol</b>	<b>Definition</b>
	Storage conditions
TB 9	Break in temperature control after 9 weeks of storage
TB 15	Break in temperature control after 15 weeks of storage
HT	High temperature treatments
DH	Different humidity conditions at 30 °C
R <sub>12h,0</sub>	Fruit rapidly cooled to 0 °C within 12 hours
R <sub>12h,2</sub>	Fruit rapidly cooled to 2 °C within 12 hours
D <sub>3d,0</sub>	Fruit directly cooled to 0 °C within 3 days
G <sub>2w,0</sub>	Fruit gradually cooled to 0 °C within 2 weeks
C <sub>1w,0</sub>	Fruit rapidly and gradually cooled to 0 °C within 1 week
C <sub>1w,2</sub>	Fruit rapidly and gradually cooled to 2 °C within 1 week