

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

At the Cutting Edge: Structural Analysis and Chemical Modification of the Edges of Mechanically Cleaved Graphene Nanoribbons

A thesis presented in partial fulfilment of the requirements for the degree
of

Doctor of Philosophy in Nanoscience

at Massey University, Manawatū, New Zealand

Haidee Michaela Dykstra

2017

Abstract

The first decade of the new carbon nanomaterial graphene has been a time of great discovery and excitement as the exceptional properties of this material were uncovered and its promise for numerous applications realised. The unique properties of graphene, including its exceptional electronic structure, are now well-established, and investigations into how these properties can be manipulated and exploited are rapidly taking off. This research contributes to the emerging field by exploring the structure and chemistry of the edges of mechanically cleaved graphene nanoribbons; groundwork for the future development of edge-modified nanoribbons that could be used to form self-assembled graphene nanoribbon composite structures with potential for devices in solar energy conversion. For this purpose, a Raman microscope was built that enabled for various aspects of the structure of graphene nanoribbons to be probed, in particular the geometry and smoothness of the edges, which have important implications for the specific reactivity of the edge carbon atoms. Chemical approaches for the specific functionalisation of the edges of the nanoribbons were developed, involving reactions tailored to the reactive groups present at the edges, and these were found to be highly successful and selective.

Acknowledgements

I would like to thank my primary supervisor, A. Prof. Mark Waterland, for his extensive guidance, input, support and patience throughout the project, as well as my co-supervisors A. Prof. Gareth Rowlands and Prof. Simon Hall for their useful advice.

I would also like to thank Olaf Griewaldt from IFS Engineering Services for his assistance in the design and machining of microscope mounts; MMIC (Manawatu Microscopy and Imaging Centre) for the use of their microtome for cutting HOPG and TEM for imaging - in particular Jordan Taylor for all the hours she has put into this; Sergei Lebedkin from Karlsruhe Institute of Technology for his very helpful advice regarding optimisations of the microscope; Krishanthi Jayasundera for her advice and guidance for the nanoribbon edge chemistry; and Dominique Appadoo for his help with the Far-IR beamline at the Australian Synchrotron.

I would also like to thank Sam Brooke, Ewan Fisher and Ashley Way for their continued collaboration, advice and input throughout the project, in particular to Ewan Fisher for his invaluable assistance in developing methodologies for characterising graphene nanoribbon samples at the Far-IR beamline at the Australian Synchrotron.

Finally, I would like to thank my family, in particular my husband Liande Gao, for their continued support throughout the project.

Contents

List of Figures	vi
List of Tables	xi
Acronyms and Abbreviations	xii
1 Introduction	1
1.1 The History of Graphene	1
1.2 The Properties and Applications of Graphene.	2
1.3 Graphene Nanoribbons	4
1.4 The Chemistry of Graphene and Graphene Nanoribbons.	6
1.5 Raman Spectroscopy for Characterising Graphene Nanoribbons	8
1.6 Project Scope, Aims and Outline	8
2 Design and Implementation of a Terahertz Raman Microscope	11
2.1 Introduction	11
2.1.1 Basic Components of a Raman Spectrometer	11
2.1.2 Holographic Bragg Filters - New Raman Notch Filter Technology	12
2.2 Design, Fabrication and Assembly of a Raman Microscope	16
2.2.1 Design	16
2.2.2 Testing Components	18
2.2.3 Preliminary Setups	19
2.2.4 Microscope Alignment	24
2.2.5 Optimisations and Improvements	31
2.2.6 Benchmarking Microscope Performance	39
2.2.7 Incorporating Additional Excitation Wavelengths	43
2.2.8 Polarised Raman	46
2.3 Conclusion	50
3 Experimental methods	51
3.1 Production of GNRs	51
3.1.1 Nanotomy-Based Mechanical Cleavage of HOPG	51
3.1.2 Exfoliation of GNBs	53
3.2 Characterisation of GNRs	54
3.2.1 AFM	54
3.2.2 TEM	54
3.2.3 Raman Microscopy	54
3.3 Functionalisation of GNRs	57
3.3.1 Chemical Methods	57

3.3.2	FTIR Spectroscopy	58
3.3.3	Raman Microscopy	61
3.3.4	SERS	61
3.3.5	THz Spectroscopy	61
3.3.6	XPS	62
4	Structure of Graphene Nanoribbons	64
4.1	Introduction	64
4.1.1	Overview of the History and Development of the Understanding of Raman Processes in Graphene and Graphite	65
4.1.2	Theory of Raman Scattering Processes in Graphite and Graphene	66
4.2	Results and Discussion	73
4.2.1	Raman Spectroscopy of HOPG and Graphene Samples	73
4.2.2	Fabrication of Graphene Nanoribbons	80
4.2.3	Characterisation of GNRs by Raman Spectroscopy, AFM and TEM	82
4.2.4	Raman Spectroscopy for the Determination of Layer Number in GNRs	89
4.2.5	D:G Ratios of GNRs	92
4.2.6	Polarised Raman Microscopy of GNRs	101
4.3	Conclusion	118
5	Edge Chemistry of Graphene Nanoribbons	119
5.1	Introduction	119
5.1.1	Clar Sextet Theory and Graphene Aromaticity	119
5.1.2	Functionalisation of Graphene and GNRs	122
5.1.3	Characterisation of Functionalised Graphene Edges	126
5.1.4	Scope and Aims	127
5.2	Results and Discussion	129
5.2.1	Ester Coupling	129
5.2.2	Amide Coupling	137
5.2.3	Diazonium Chemistry	145
5.2.4	SERS	150
5.2.5	THz Spectroscopy	155
5.3	Conclusion	159
6	Conclusion and Future Perspectives	160
7	Bibliography	164
8	Appendix A	179
8.1	Calculation for Monolayer Coverage of SDS on GNRs	179

8.2	Estimation of Edge:Basal Carbon Ratios in GNRs	180
9	Appendix B	181
9.1	Additional AFM images	181
9.2	Additional TEM images	185
9.3	Additional Raman data	192
9.3.1	HOPG	192
9.3.2	Graphene test samples	194
9.3.3	GNRs	195
9.3.4	Polarised Raman	200
9.3.5	Functionalised GNR	248
9.4	Additional FTIR data	251
9.5	Additional Far-IR data	256
9.6	Additional SERS data	257

List of Figures

Introduction	1
Figure 1 (a) Zigzag and (b) armchair GNRs.	5
Design and Implementation of a Terahertz Raman Microscope	11
Figure 2 A simple Raman setup.	12
Figure 3 Reflecting (left) and transmitting (right) VBG filters.	14
Figure 4 Diagram depicting how interference can be avoided when multiple VBG filters are used in series.	14
Figure 5 The IX70 microscope after removal of the lamp head, stage, objectives and beamsplitter rail.	17
Figure 6 (a) CAD design for 20° diffraction at the BPF and (b) photograph of actual setup.	17
Figure 7 (a) Prism mirror mounted to the beamsplitter rail.	18
Figure 8 (a) Image showing diffraction of the laser beam, collimated by the FiberPort, from the BPF.	19
Figure 9 (a) Laser spectrum at higher power, without BPF cleaning.	20
Figure 10 Diagram of the preliminary Raman setup.	21
Figure 11 Spectrum of the laser background.	21
Figure 12 Raman setup with separate objectives to illuminate the sample and collect the scattered light.	23
Figure 13 Raman spectrum of cyclohexane with 90° collection geometry	23
Figure 14 Raman spectrum of cyclohexane with backscattering geometry and Rayleigh suppression with 2 BNFs.	24
Figure 15 Beam collimating and expanding optics incorporated into a cage system.	25
Figure 16 (a) Drawing of the system used for initial alignment of the BPF.	26
Figure 17 Photograph of the FiberPort positioned to collect the scattered light exiting the microscope side-port.	28
Figure 18 Initial Raman spectrum of cyclohexane, taken with a 4x microscope objective.	29
Figure 19 Drawing of the final microscope setup.	30
Figure 20 Raman spectrum of cyclohexane, with 3 BNFs rejecting the Rayleigh line.	30
Figure 21 Photograph of the collection fibre and tube lens aligned in a cage system.	32
Figure 22 Spectra of (a) fluorescent glass coverslip and (b) Si wafer	34

Figure 23	Diagram depicting how chromatic aberration occurring in the microscope objective lens can lead to different effective focal lengths	35
Figure 24	Raman spectra of Si wafer	36
Figure 25	Raman spectra of Si wafer, taken with a series of achromat objectives.	37
Figure 26	Plots of Si intensity vs NA (a) and (b) and Raman:Rayleigh ratio vs NA (c) for different microscope objectives.	38
Figure 27	Full microscope setup.	40
Figure 28	(a) Raman spectrum of HOPG, laser power 5 mW, integration time 60s.	41
Figure 29	Raman spectra of L-cystine with multi-mode collection fibres with different core diameters.	42
Figure 30	Raman spectra of Si wafer acquired with two different spectrograph gratings using 532 nm excitation.	44
Figure 31	Raman spectra of (a) sulfur and (b) L-cystine, taken with 532 nm excitation.	45
Figure 32	Polarised Raman spectra of cyclohexane, taken with a 4x objective.	46
Figure 33	Polarised Raman spectra of a linear polariser acquired with 532 nm excitation.	48
Figure 34	Polar plot of the Raman intensity of the linear polariser sample . .	49
Experimental methods		51
Figure 35	View from the top showing the water-filled knife boat and HOPG sample	52
Figure 36	Curve-fitting procedure in OPUS for a typical Raman spectrum of GNRs.	56
Figure 37	FTIR spectra of GNR edge-functionalised with sulfanilic acid. . .	60
Structure of Graphene Nanoribbons		64
Figure 38	(a) The unit cell of monolayer graphene, spanned by the vectors \vec{a}_1 and \vec{a}_2	66
Figure 39	(a) Breathing mode for the D band of graphene.	67
Figure 40	(a) Representation in real space of D peak Raman scattering at an ordered edge.	68
Figure 41	(a) Wavevectors of defects for armchair (\vec{d}_a) and zigzag (\vec{d}_z) edges in graphene.	69
Figure 42	The doubly degenerate E_{2g} LO (a) and TO (b) phonon modes of the G band in graphene.	70
Figure 43	Calculated symmetries and frequencies for the shear modes of graphene with differing layer numbers.	71
Figure 44	Raman spectra of HOPG acquired with 785 nm excitation.	73

Figure 45	Raman spectra of HOPG acquired with 532 nm excitation.	74
Figure 46	Raman spectra of HOPG and various graphene samples acquired with 785 nm excitation.	75
Figure 47	Raman spectra of (a) HOPG and (b) CVD graphene on a Ni substrate.	77
Figure 48	Raman spectra of monolayer CVD graphene on (a) a glass substrate and (b) a SiO ₂ substrate.	78
Figure 49	Raman spectra of (a) single/double CVD graphene (b) and (c) multilayer CVD graphene	79
Figure 50	Nanotomy method for the production of GNRs.	81
Figure 51	(a) AFM image of 100 nm wide GNBs on a mica substrate, acquired in tapping mode.	83
Figure 52	(a) AFM image of exfoliated GNRs.	84
Figure 53	AFM in tapping mode of 100 nm GNR on a mica substrate.	85
Figure 54	AFM in tapping mode of a GNR aggregate on mica.	86
Figure 55	TEM images of GNRs cut at (a) 300 nm and (b) 20 nm.	87
Figure 56	TEM images of GNRs cut at 100 nm.	88
Figure 57	Raman spectra acquired with 785 nm excitation of the low-frequency region of HOPG, 100 nm GNB and 100 nm GNR.	89
Figure 58	2D bands of Raman spectra of 100 nm and 300 nm wide GNRs fit with Lorentzians.	91
Figure 59	D:G ratios for (a) dry- and (b) wet-cut GNRs plotted as a function of ribbon width.	93
Figure 60	(a) Averaged D:G ratios over all sonication powers plotted as a function of ribbon width.	94
Figure 61	D:2D ratios for (a) dry- and (b) wet-cut GNRs plotted as a function of ribbon width.	95
Figure 62	(a) Averaged D:2D ratios over all sonication powers plotted as a function of ribbon width.	96
Figure 63	Averaged Raman spectra of wet- and dry-cut 100 nm GNRs acquired with 532 nm excitation.	97
Figure 64	Averaged Raman spectra of wet- and dry-cut 100 nm GNRs acquired with 488 nm excitation.	98
Figure 65	Averaged D:G ratios plotted as a function of ribbon width (see figure 60a) with error bars.	99
Figure 66	(a) Intensities of the D and G bands of HOPG plotted as a function of the degree of rotation of the plane of polarisation of the input laser beam.	104
Figure 67	D:G ratio of HOPG Raman spectra plotted as a function of polarisation, with a cos ² θ fit.	105

Figure 68	G:2D, D:G and D:2D ratios of 500 nm wet-cut GNR plotted as a function of polarisation.	108
Figure 69	G:2D, D:G and D:2D ratios of 300 nm wet-cut GNR plotted as a function of polarisation.	110
Figure 70	G:2D, D:G and D:2D ratios of 300 nm wet-cut GNR plotted as a function of polarisation.	111
Figure 71	G:2D, D:G and D:2D ratios of 100 nm dry-cut GNR plotted as a function of polarisation.	112
Figure 72	G:2D, D:G and D:2D ratios of 500 nm wet-cut GNR plotted as a function of polarisation.	113
Figure 73	G:2D, D:G and D:2D ratios of 500 nm GNR plotted as a function of polarisation.	114
Figure 74	G:2D, D:G and D:2D ratios of 500 nm GNR plotted as a function of polarisation.	115
Figure 75	G:2D, D:G and D:2D ratios of 500 nm wet-cut GNR plotted as a function of polarisation.	116
Figure 76	G:2D, D:G and D:2D ratios of 100 nm wet-cut GNR plotted as a function of polarisation.	117
Figure 77	G:2D, D:G and D:2D ratios of 100 nm dry-cut GNR plotted as a function of polarisation.	117
Edge Chemistry of Graphene Nanoribbons		119
Figure 78	Resonance structures of a graphene sheet.	120
Figure 79	The three different width-dependent Clar formulae for armchair-edged GNRs	120
Figure 80	Segment of a zigzag-edged GNR with migration of Clar sextets depicted by red arrows.	121
Figure 81	Alternative representation of the Clar structure shown in figure 80.	121
Figure 82	Illustration of the symmetry of A and B sublattices in (a) armchair and (b) zigzag GNRs.	122
Figure 83	Mechanism for covalent functionalisation of graphene with diazonium salts via aryl radical formation.	124
Figure 84	Reaction scheme for functionalisation of GNRs with 4-aminobenzoic acid via EDC/DMAP ester bond formation.	130
Figure 85	Chemical structures of the modifiers used in the ester coupling reactions.	130
Figure 86	FTIR microscope spectra of unmodified GO and GO functionalised with Rhb and 4-NBA.	131

Figure 87	Raman spectra of GO and modified GO samples acquired with 785 nm excitation.	132
Figure 88	Raman spectra of as-cut GNR, GNR reduced with NaBH ₄ and GNR modified with 4-ABA.	133
Figure 89	ATR-FTIR spectra of r-GNR, 4-ABA-GNR and RhB-GNR.	133
Figure 90	Raman spectra of the samples in figure 89 acquired with 532 nm excitation.	134
Figure 91	Raman spectra of 50 nm modified and unmodified GNR samples acquired with 532 nm excitation.	135
Figure 92	ATR-FTIR spectra of 50 nm r-GNR with various edge modifications.	136
Figure 93	Reaction scheme for the amidation of GNR COOH groups.	137
Figure 94	Rearrangement of the O-acylurea intermediate to form an N-acylurea.	137
Figure 95	Chemical structures of the modifiers used in the amide coupling reactions.	138
Figure 96	FTIR microscope reflectance spectra of GO and GO modified with SA and 4-NA.	139
Figure 97	Raman spectra of GO and f-GO samples acquired with 785 nm excitation.	139
Figure 98	FTIR microscope spectra of multiple SA-GNR samples	140
Figure 99	(a) Comparison of SA-GO and SA-GNR FTIR spectra.	141
Figure 100	Raman spectra of 20 nm GNR samples acquired with 532 nm excitation.	142
Figure 101	FTIR microscope spectra of 20 nm f-GNR.	143
Figure 102	Raman spectra acquired with 488 nm excitation.	144
Figure 103	FTIR microscope spectra of 20 nm GNR modified with SP groups.	145
Figure 104	FTIR microscope spectra of 20 nm GNR modified with NP groups	146
Figure 105	Raman spectra of 20 nm GNR modified with (a) SP and (b) NP groups.	148
Figure 106	Comparison Raman spectra of 20 nm GNR modified NP groups . .	149
Figure 107	SERS spectra acquired with 488 nm excitation.	151
Figure 108	SERS spectra acquired with 532 nm excitation.	153
Figure 109	Raman and SERS spectra of 4-NA and NP-GNR samples acquired with 488 nm excitation.	154
Figure 110	GNRs edge-functionalised with various combinations of multi-carboxylated molecules and ethylenediamine.	156
Figure 111	Far-IR spectra of 20 nm GNR with various edge modifications. . .	157

List of Tables

Experimental methods	51
1 Samples prepared for the Far-IR beamline at the Australian Synchrotron.	63
Structure of Graphene Nanoribbons	64
2 Comparison of average D:G and D:2D ratios for 100 nm wet- and dry-cut GNRs with 532 nm and 488 nm excitation.	98
3 Variation in D:G ratios for the dry-cut GNRs in figure 65.	99
4 Variation in D:G ratios for the wet-cut GNRs in figure 65.	99
5 Variation in D:G ratios for the 100 nm GNRs in figures 63 and 64.	100

List of Acronyms and Abbreviations

In order of appearance:

CVD	Chemical vapour deposition
GO	Graphene oxide
BZ	Brilluoin zone
QHE	Quantum Hall effect
FET	Field effect transistor
LED	Light-emitting diode
ITO	Indium Tin oxide
SAM	Self-assembled monolayer
DOS	Density of states
GNR	Graphene nanoribbon
CNT	Carbon nanotube
HOPG	Highly-oriented pyrolytic graphite
HNF	Holographic notch filter
VBG	Volume Bragg grating
VHG	Volume holographic filter
PTR	Photo-thermo-refractive
BNF	BragGrate TM notch filter
BPF	BragGrate TM bandpass filter
OD	Optical density
NA	Numerical aperture
OPSL	Optically pumped semiconductor laser
MFD	Mode field diameter
PM	Polarisation maintaining
PMMA	Poly(methyl) methacrylate
GNB	Graphite nanoblock
AFM	Atomic force microscopy
TEM	Transmission electron microscopy
IPA	Isopropanol
SDS	Sodium dodecylsulfate
CMC	Critical micelle concentration
THF	Tetrahydrofuran
r-GNR	Reduced GNR
4-ABA	4-aminobenzoic acid
EDC	1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide
DMAP	4-Dimethylaminopyridine

DMSO	Dimethyl sulfoxide
f-GNR	functionalised GNR
SA	Sulfanilic acid
NHS	N-Hydroxysuccinimide
DMF	Dimethylformamide
FTIR	Fourier transform infrared
ATR	Attenuated total reflection
S:N	Signal-to-noise
HDFT	Heptadecafluoro-1-decanethiol
KK	Kramers-Kronig
SERS	Surface-enhanced Raman spectroscopy
CTAB	Cetyltrimethylammonium bromide
Agnp	Silver nanoparticle
PE	Polyethylene
XPS	X-ray photoelectron spectroscopy
TA	Terephthalic acid
TMA	Trimesic acid
BTB	Benzene 1,3,5-tribenzoic acid
EDA	Ethylenediamine
TFA	Trifluoroacetic acid
4-NA	4-nitroaniline
PDOS	Phonon density of states
iTO	In-plane transverse optical
iLO	In-plane longitudinal optical
DR	Double resonance
PAH	Polycyclic aromatic hydrocarbon
NMP	N-methyl-pyrrolidone
SDBS	Sodium dodecylbenzenesulfonate
BDE	Bond dissociation energy
OTS	Octadecyltrichlorosilane
hBN	Hexagonal Boron nitride
STM	Scanning tunnelling microscopy
TERS	Tip-enhanced Raman spectroscopy
AE	Activated ester
4-NBA	4-nitrobenzoic acid
RhB	Rhodamine B
SP	Sulfophenyl
NP	Nitrophenyl
SPR	Surface plasmon resonance

R6G

HATN

TCNQ

F4-TCNQ

Rhodamine 6G

hexaazatriphenylene

7,7,8,8-tetracyanoquinodimethane

2,3,5,6-tetrafluoro-7,7,8,8-
tetracyanoquinodimethane