

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**SYNTHESIS OF AN ARGININE MIMIC FOR  
AN ANTIFUNGAL OCTAPEPTIDE LIBRARY**

This thesis is presented in partial fulfilment of the requirements for the  
degree of Master of Science in Chemistry at Massey University,  
Palmerston North, New Zealand

**Hao Sun**

**2008**

## ABSTRACT

A family of bioactive antifungal octapeptides has been established<sup>1-3</sup>. The template for this library is based on peptides composed of two predefined amino acids, three arginines, and three unknown components. The arginine contains a guanidino group (pKa =12.48) which is readily protonated under physiological conditions, and hence can bind to fungal cell walls which possess a negatively charged surface. Other components of this peptide are used to kill the fungi by inhibition of their sodium transport system<sup>1-5</sup>.

To date this story has survived on the fact that D-amino acids work best. However both D and L arginine derivatives are quite expensive. In collaboration with the research group in the University of Otago, it was planned to develop an effective and economic method to produce a series of compounds to replace arginine. These mimics were designed to be artificial amino acids as  $\text{NH}_2\text{---N}^*(\text{R}')\text{---COOR}$  where the side chain R' carries a guanidino group. Thus the chiral carbon is now replaced by an amide group hence no stereocenter, no enantiomers but the guanidino group will retain a high pKa similar to that of the natural arginine. For the synthetic strategy, the target molecule was divided into two parts, one the amino acid backbone, and the other the guanidine containing side chain. Each segment was built up separately and finally combined together.

The backbone of the arginine mimic must possess two amino groups. One is at the N-terminal for the next step in peptide synthesis, the other one (N\*) is used to connect to the guanidino containing side chain. This was reacted with selected acetate derivatives (*t*-butylchloroacetate, benzylchloroacetate, methylbromoacetate) that had potential to produce the desired backbone in good yield.

Construction of the side chain (R') to be composed of a di-protected guanidine and a carboxylic group was the most challenging and difficult part of this project. To achieve this, two different approaches were studied. One was to use a primary amino acid  $\text{NH}_2\text{---}(\text{CH}_2)_x\text{---COOH}$  (X=1,2,5) to react with a guanylating reagent to make the unprotected side chain, then two protecting groups were added to the two nitrogen containing groups of the guanidine. The other method was protection of a guanylating reagent (usually carboxamide compounds) then reaction with a primary amino acid to make a di-protected guanidine containing side chain.

In amino acid and peptide chemistry, to avoid self condensation and by-product formation, selectivity of amino and carboxylic groups is very crucial hence application of different types of protecting groups are the basis of peptide construction. Therefore in this project, employment of suitable protecting groups at both N- and C-terminals was incorporated in this study.

Finally, one arginine mimic *N*-[*N'*-((9-fluorenyl)methoxycarbonyl)-2-aminoethyl]-*N*-*t*-butyloxycarbonylmethyl-3-*N',N''*-bis(*t*-butyloxycarbonyl)carbamidinopropanamide was successfully built up. It will be used for the construction of octapeptides for the current antifungal programme. The final product will be sent to University of Otago for bio-activity tests.

## ACKNOWLEDGEMENTS

Firstly I would like to record my appreciation to Associate Professor David R. K. Harding as my research supervisor for his guidance and assistance in helping me complete my Masterate and opening the door to peptide chemistry.

I also wish to thank Mr David Lun for his constant instruction and assistance in the operation of equipment (HPLC/MS). A special thanks to Dr Krishanthi Jayasundera who taught me a lot about Flash Column Chromatography. Thanks to Mr Sebastian Parker for his suggestions and assistance in amide bond formation in this project.

Similarly I express my appreciation to Dr Wayne Campbell and Mr Adam Stephenson for their efforts in proof reading drafts of this thesis. Thanks to Mr Ross Davidson for the assistance with the calculation of bond angles of molecules in this project.

Furthermore, thanks to all the academic staff and other members of Chemistry Discipline, Institute of Fundamental Sciences, especially Professor Andrew M. Brodie, Associate Professor Carol Taylor, Associate Professor Trevor Kitson and Associate Professor Eric Ainscough for their support and suggestions throughout my two years Masterate study at Massey University.

## TABLE OF CONTENTS

Abstract	I	
Acknowledgements	III	
List of Figures	VII	
List of Tables	X	
List of Abbreviation	XI	
<b>CHAPTER ONE</b>	<b>INTRODUCTION</b>	<b>01</b>
1.1	A D-Octapeptide Combinatorial Library	01
1.2	Natural Arginine and the Guanidino Group	03
1.3	Structure of a Designed Arginine Mimic and the Synthetic Strategy	05
1.4	Strategy for the Synthesis of a Protected Guanidino Group	07
1.5	Protection of the N-Terminus in Peptide Chemistry	09
1.6	Protection of C-Terminus in Peptide Chemistry	13
1.7	Coupling Reactions	14
<b>CHAPTER TWO</b>	<b>EQUIPMENT AND MATERIALS</b>	<b>16</b>
2.1	Equipment	16
2.2	Materials	16
<b>CHAPTER THREE</b>	<b>FORMATION OF THE PROTECTED GUANIDINO GROUP</b>	<b>17</b>



4.4	Experimental	52
<b>CHAPTER FIVE</b>	<b>BACKBONE SYNTHESIS</b>	<b>54</b>
5.1	Introduction	54
5.2	Synthesis and Discussion	56
5.3	Conclusion	58
5.4	Experimental	58
<b>CHAPTER SIX</b>	<b>COUPLING OF THE GUANIDINE CONTAINING SIDE CHAIN TO THE AMINO ACID BACKBONE</b>	<b>60</b>
6.1	Introduction	60
6.2	Discussion	61
6.3	Conclusion	62
6.4	Experimental	62
<b>CHAPTER SEVEN</b>	<b>SUMMARY AND CONCLUSION</b>	<b>64</b>
<b>CHAPTER EIGHT</b>	<b>CONCLUSION</b>	<b>67</b>
<b>REFERENCES</b>		<b>69</b>



## LIST OF FIGURES

Figure 1	Designed D-octapeptide and the combinatorial library	02
Figure 2	Structure of natural L and D arginine	03
Figure 3	Resonance structures of guanidine	04
Figure 4	Structure of the target arginine mimic in this project	05
Figure 5	Synthetic strategy of arginine mimics	06
Figure 6	Structure and molecular weight of glycine, $\beta$ -alanine and 6-aminohexanoic acid	06
Figure 7	Structure of a typical guanylation reagent and the normal process of formation of a guanidino group	07
Figure 8	Four possibilities of di-peptide formation using unprotected amino acids	10
Figure 9	Design of a protected arginine mimic	11
Figure 10	Mechanisms of cleavage of Fmoc and Boc groups	12
Figure 11	The structures of DCC and HBTU	15
Figure 12	The three routes for the formation of substituted guanidines	17
Figure 13	The mechanism of Mitsunobu reaction to achieve substituted guanidine formation	19
Figure 14	Synthetic strategy for making the di-Boc guanidino acid from the guanidine hydrochloride	20
Figure 15	Formation of 1H-pyrazole-1-carboxamide hydrochloride	24
Figure 16	Reaction of 1H-pyrazole-1-carboxamide hydrochloride with a primary amino acid	24
Figure 17	Self-condensation of 1H-pyrazole-1-carboxamide and of <i>N</i> -alkyldiguanidine	26
Figure 18	Formation of guanidino acids from reaction of aminoiminomethanesulfonic acid with primary amino acids	27
Figure 19	Structure of unprotected guanidino acids	32
Figure 20	Structure of di- <i>t</i> -butyl dicarbonate and the	

	mechanism of its amine protecting reaction	34
Figure 21	Preparation of di-Boc-protected guanidino acid	35
Figure 22	Three reagents for the introduction of the Cbz group	36
Figure 23	Preparation of di-Cbz protected guanidino acids using benzyloxycarbonylsuccinimide	37
Figure 24	Di-tosyl protected arginine mimic	38
Figure 25	Attempted di-protection of a guanidino group with <i>p</i> -toluenesulfonyl chloride	39
Figure 26	Angles of the guanidino group for the unprotected and mono protected guanidino acids	41
Figure 27	Angles of the guanidino group for thiourea and S-methylisothiourea and their di-Boc products	43
Figure 28	Basic mechanism for the reaction of <i>N,N'</i> -bis- <i>t</i> -Butoxycarbonylthiourea with amines catalyzed by HgCl <sub>2</sub>	45
Figure 29	Formation of di-protected guanidine acids from S-methylisothiourea	46
Figure 30	Protonation of pyrazole	49
Figure 31	Formation of 1H-pyrazole-1-[ <i>N,N''</i> - -bis( <i>t</i> -Butoxycarbonyl)]carboxamidine	49
Figure 32	Production of guanidines from 1H-Pyrazole-1- [ <i>N,N''</i> -bis( <i>t</i> -butoxycarbonyl)]carboxamidine	50
Figure 33	Designed backbone structure	54
Figure 34	Basic process of solid phase peptide synthesis	55
Figure 35	Mechanism for removal of the 9- fluoromethyloxycarbonyl group (Fmoc)	56
Figure 36	Reaction scheme for the construction of the protected backbone amino acid	57
Figure 37	Structure and reaction scheme for the coupling reagents DCC and HBTU	61
Figure 38	The arginine mimic <i>N</i> -[ <i>N'</i> -((9-fluorenyl)methoxycarbonyl)- 2-aminoethyl]- <i>N-t</i> -butyloxycarbonylmethyl- 3- <i>N',N''</i> -bis( <i>t</i> -butyloxycarbonyl)carbamidinopropanamide produced in this project	64

Figure 39	Designed arginine mimic in which benzyl ester is used as C-terminal block group	65
Figure 40	Mechanism for the removal of the benzyl group on the C-terminal by hydrogenolysis	65

## LIST OF TABLES

Table 1	Experimental results for the reaction of 1H-Pyrazole-1-carboxamide hydrochloride with three primary amino acids	25
Table 2	Experimental results for the reaction of aminoiminomethanesulfonic acid with three primary amino acids	27
Table 3	Experimental results for the formation of di-Boc guanidino acids from the reaction of the <i>N,N'</i> -bis ( <i>t</i> -butyloxycarbonyl)- <i>S</i> -methylisothiourea with three primary amino acids	47
Table 4	Basicity of common amines	60

## LIST OF ABBREVIATIONS

Arg	arginine
Boc	<i>t</i> -butyloxycarbonyl
(Boc) <sub>2</sub> O	di- <i>t</i> -butyl dicarbonate
Cbz	benzyloxycarbonyl
(Cbz) <sub>2</sub> O	dibenzyl dicarbonate
Cbz-OSu	benzyloxycarbonylsuccinimide
DCC	dicyclohexylcarbodiimide
DCM	dichloromethane
DEAD	diethyl azodicarboxylate
DIEA	diisopropylethylamine
Fmoc	9-fluorenylmethyloxycarbonyl
Fmoc-OSu	<i>N</i> -(9-fluorenylmethoxycarbonyloxy)succinimide
HBTU	O-Benzotriazol-1-yl- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
ortho-Br-Cbz	ortho-bromobenzyloxycarbonyl
ortho-Cl-Cbz	ortho-chlorobenzyloxycarbonyl
PNA	peptide nucleic acid
PPh <sub>3</sub>	triphenylphosphine
SPPS	solid phase peptide synthesis
TFA	trifluoroacetic acid