

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**ENVIRONMENTAL EFFECTS OF DENSELY PLANTED
WILLOW AND POPLAR IN A SILVOPASTORAL SYSTEM**

*A thesis presented in partial fulfilment
of the requirements for the degree of*

Doctor of Philosophy (PhD)

in
Agroforestry



Institute of Natural Resources
Massey University
Palmerston North, New Zealand

ZAKER HUSSAIN

2007

***THIS THESIS IS DEDICATED TO MY PARENTS
WHO ALWAYS PRAYED FOR MY SUCCESS TO
GAIN HIGHER QUALIFICATION***

Abstract

New Zealand, having large areas of hilly landscapes, is subject to the risk of soil erosion, and summer and autumn droughts that limit pasture growth, which in turn affects the livestock-based economy. The nitrogen and phosphorus input in fertilisers coupled with livestock excreta and soil disturbance impose a serious threat to downstream water quality. The planting of trees is one option used to decrease soil erosion, increase the quantity of forage and manage runoff.

To date, research has mainly focused on wide spaced poplar trees for feed quality and their effects on understorey pasture growth. However, there is increasing interest in the use of densely planted willow and poplar for fodder purpose. The effects of young (< 5 yrs old) willow and poplar planted at close spacing on runoff, soil erosion, growth of understorey pasture and nutrient losses have never been studied in New Zealand.

Three field trials (two at Crop and Research Unit, Moginie, Manawatu and one at Riverside Farm, Masterton) were conducted between October 2004 and November 2006 that incorporated comparative establishment and growth of densely planted willow and poplar and their effects on soil moisture, runoff, sediment load and nutrient losses from grazed and fertilised farmland.

It was concluded that densely planted willow and poplar (3-4 yrs) reduced total nitrogen (TN) and dissolved reactive phosphorus (DRP) by 47 % each and sediment load by 52 %. Young trees reduced surface runoff and soil moisture more as they aged. However, due to their deciduous nature willow and poplar were not effective in reducing surface runoff in winter and early spring. Sheep preferred camping under trees, especially in late spring and summer, and this led to greater deposition of dung and urine under trees than open pasture. Sheep grazing, especially in winter, significantly increased sediment and nutrient loads in runoff water. The N and P fertiliser application increased nutrient load in runoff water well above the threshold level required to initiate algal growth to create eutrophication.

Densely planted willow and poplar significantly reduced understorey pasture growth by 23 % and 9 %, respectively, in their second year at Moginie, mainly due to shade, but coupled with soil moisture deficit in summer. The pasture growth in a willow browse block was 52 % of that in open pasture as a result of shade and differences in pasture species composition. Sheep browsing reduced willow leaf area significantly. Willow and poplar survival rates were similar ($P > 0.05$) after two years of establishment (100 % vs 90.5 %, respectively). However, willow grew faster than poplar in height (1.90 vs 1.35 m), stem diameter (43.5 vs 32.6 mm), canopy diameter (69 vs 34 cm) and number of shoots (8.7 vs 2.3) at the age of two years, respectively.

The research clearly demonstrated that densely planted young willow and poplar trees can reduce runoff, sediment load and nutrient losses from farmland to freshwater, but shade and soil moisture can limit pasture growth under trees. It is recommended that willow and poplar should be planted at wide spacing on the whole farm to minimise loss of pasture. Where blocks of trees are necessary, such as willow browse blocks, sheep browsing can be used as a tool to reduce shade to improve pasture growth. Livestock access to riparian strips should be minimal to avoid livestock camping that can have deleterious effects on water quality.

Key words: willow; poplar; growth; nutrients; pasture; runoff; soil erosion; soil moisture; fertiliser; water; sheep; browsing; camping.

Acknowledgements

The exploration of environmental effects of willow and poplar in a silvopastoral system was not possible without the physical, financial, technical and moral support of many individuals and some organisations. These acknowledgements are intended to express gratitude and appreciation to all those who contributed to the success of this thesis. However, undoubtedly, God almighty stands first and foremost for blessing me with the health and mental capability to undertake this study.

Sincere and heartfelt gratitude to my chief supervisor Professor Peter Kemp for his support, guidance and patience throughout my study. His sympathetic behaviour, generosity and great encouragement were instrumental to my academic career at Massey. I also thank my co-supervisor Dr David Horne for his supervision and technical input in my thesis. Dr Horne made a significant contribution through corrections and fruitful criticism of manuscripts of this thesis.

The physical and technical support by Mr. Mark Osborne, senior technician, is greatly appreciated. My thanks to Mr. Tom Dodd and Mr. Ross Wallace for their physical input in setting up experiments and data collection, Mr. Bob Toes for constructing the tipping buckets and help with laboratory analyses, Ian Furkert, and Ms. Glenys Wallace for their support in laboratory works and analyses. Thanks to INR office support staff, especially Denise Stewart, Denise Brunskill and Glenys Gilligan for their friendly behaviour and always available support.

Special thanks to Mr. James Hanley, for his technical support in instrumentation of runoff measurements and nutrient analyses. I thank Dr Alasdair Noble, Dr Komang Jaya and Anne West for their contribution in statistical analyses of the data. Thanks to Mr. Lance Currie for his support through good administration and help for analyses of soil samples. I appreciate Dr James Millner for his technical advice and provision of instruments for the collection of tree growth data. Dr Cory Matthew has been generous with his advice whenever and where ever approached. Mike Bretherton for developing the Excel Macro for

processing the tipping bucket data logger files, used to measure surface runoff. I appreciate Dr Baisen Zhang, Dr Entin Daningsih, Dr Tehseen Aslam and Edmundo Viegas for their advice and company during their doctoral studies at Massey. Thanks to Mr. Bhoj Bahadur Kshatri and Zulkefly Sulaiman for helping me in data collection and having useful discussions on my research. Thanks to Girija Page, Bijaya, Diwas and other postgraduate students for their company.

My gratitude to the Doctoral Research Committee chairman and members for granting me the 'Massey doctoral scholarship' that laid the foundation for my PhD. I also acknowledge the financial support from Sustainable Farming Fund Project, Helen E. Akers and John Hodgson pastoral science scholarships. Special thanks to AJK Forest Department for the grant of study leave.

It is a great pleasure to acknowledge the friendship, advice and kind support by Dr M. A. Choudhary, Dr Manzoor-ul Haque Awan and Mr. Zulfiqar Haider Butt that made it possible for me to join Massey University and achieve this success. I extend my gratitude to my friends in Pakistan, in particular, Mr. Tanveer Hussain Qureshi, Mr. Muhammad Shafiq, Mr. Muhammad Arif Qureshi, Mr. Abdul Hamid, Mr. Mustafa Mughal, Mr. Muhammad Tayyab, Mr. Mushtaq Qureshi, Mr. Abdul Rauf Qureshi and Abdul Rashid Jhang for their prayer and appreciation that has been instrumental to me during my study at Massey.

I acknowledge the great support and friendship of Mr. Tahir Mahmood and Mrs. Amsha Nahid from day one of my arrival in New Zealand. Their kind hospitality and their role in looking after my family were of great value to me and will never be forgotten. Similarly, Mr. and Mrs. Babar Saeed, Mr. and Mrs. Muhammad Nasim, Mr. and Mrs. Tariq Choudhry, Mr. and Mrs. Sarfraz, Mr. and Mrs. Muhammad Ali and Mr and Mrs. Ahmer Khawaja deserve great appreciation for their friendship and support.

My humble and heartfelt acknowledgement goes to my beloved Father and mother who prayed so much for my success. I can't be indifferent to my brothers and sister for their love and appreciation to my success. Finally, I appreciate the patience and support of my wife Samina. My daughters Anum and Aminah have been a source of pleasure and relief during stress.

Table of contents

	Page
Abstract.....	i
Acknowledgements.....	iii
Table of contents	v
List of Tables.....	vi
List of Figures	viii
List of Plates	xv
List of Appendices.....	xvii
Glossary and abbreviations	xviii
CHAPTER 1:	
General introduction.....	1
CHAPTER 2:	
Literature review	11
CHAPTER 3:	
Establishment of densely planted willow and poplar and their effect on understorey pasture growth	66
CHAPTER 4:	
Pasture growth in a willow browse block.....	104
CHAPTER 5:	
Effects of densely planted willow and poplar on runoff, sediment and nutrient losses.....	145
CHAPTER 6:	
General discussion and conclusions.....	216

List of Tables

Table 2.1	The distinguishing features of the genus <i>Populus</i> and <i>Salix</i> . -----	15
Table 3.1	The spring pasture species composition (%) from willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at Moginie, Manawatu. -----	81
Table 3.2	Seasonal NHA rate (kg DM ha ⁻¹ day ⁻¹) in willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at Moginie, Manawatu. -----	83
Table 4.1	Soil Olsen P (µg g ⁻¹) and pH in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm near Masterton, Wairarapa. -----	114
Table 4.2	Soil bulk density (g cm ⁻³) in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm near Masterton, Wairarapa. -----	115
Table 4.3	Pasture species composition (%) in early autumn 2005 in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm, near Masterton, Wairarapa. -----	116
Table 4.4	Pasture species composition (%) in spring 2005 in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm near Masterton, Wairarapa. -----	117
Table 4.5	Seasonal NHA rate (kg DM ha ⁻¹ day ⁻¹) for 2004-05 in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm near Masterton, Wairarapa. -----	118
Table 4.6	Least square means comparison for volumetric soil moisture content (%) between willow pasture (WP) and open pasture (OP) treatments at Riverside Farm near Masterton, Wairarapa. -----	123
Table 5.1	Least square means for volumetric soil moisture content (%) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. -----	159
Table 5.2	Seasonal runoff (mm) comparison between open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. -----	162

Table 5.3	Seasonal suspended solid losses (kg ha^{-1}) in surface runoff (mm) from open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu.	165
Table 5.4	Least square means for suspended solids concentrations (mg L^{-1}) and losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu.	166
Table 5.5	Total P concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu.	168
Table 5.6	Seasonal comparison for total P and DRP losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu.	169
Table 5.7	Least square means of total P and DRP concentrations (mg L^{-1}) and losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu.	170
Table 5.8	Seasonal comparison of total nitrogen (TN), ammonium-N ($\text{NH}_4^+\text{-N}$) and nitrate-N ($\text{NO}_3^-\text{-N}$) losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu.	177
Table 5.9	Least square means of concentrations (mg L^{-1}) and losses (kg ha^{-1}) of total nitrogen (TN), total kjeldahl nitrogen (TKN), ammonium-N ($\text{NH}_4^+\text{-N}$) and nitrate-N ($\text{NO}_3^-\text{-N}$) in surface runoff in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu.	180
Table 5.10	Least squares means for suspended solids concentrations (mg L^{-1}) and losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and willow pasture (WP) treatments at Riverside Farm near Masterton, Wairarapa.	188
Table 5.11	Least squares means for total P and DRP concentrations (mg L^{-1}) and losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and willow pasture (WP) treatments at Riverside Farm near Masterton, Wairarapa.	190
Table 5.12	Least squares means for TN and $\text{NH}_4^+\text{-N}$ concentrations (mg L^{-1}) and losses (kg ha^{-1}) at Riverside Farm near Masterton, Wairarapa.	193

List of Figures

Figure 1.1	Outline of the Thesis. -----	7
Figure 2.1	P can be released from soil and plant material to surface and subsurface runoff water or lost by erosion.-----	35
Figure 3.1	The canopies of four nuclei trees (red circles) planted as a square grid (1.2 ×1.2 m spacing). The square with blue lines indicates the actual spacing at which trees were planted while the areas of red circles inside the square show the canopy. -----	71
Figure 3.2	Total monthly rainfall (mm) and mean air and soil temperature (°C) during the experimental period in Palmerston North, New Zealand. -----	76
Figure 3.3	Data pattern shows the relationship between the tree stem diameter (mm) and tree height (m) of willow and poplar at Moginie, Manawatu.-----	78
Figure 3.4	Data pattern shows linear relationship between initial tree stem diameter (mm) and final stem diameter (mm) of willow and poplar at Moginie, Manawatu (n = 342, P < 0.0001). -----	78
Figure 3.5	Month wise mean daily NHA rate (kg DM ha ⁻¹ day ⁻¹) from willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at Moginie, Manawatu. Values for July and August were equally distributed. Bars show the SEM for each treatment. * denotes significance at P < 0.05. -----	82
Figure 3.6	Volumetric soil moisture content (%) from willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at 0-150 mm depth at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at P < 0.05. -----	84
Figure 3.7	Volumetric soil moisture content (%) from willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at 0-300 mm depth at Moginie, Manawatu. The line bars show the SEM for each treatment. * denotes significance at P < 0.05. -----	85

Figure 3.8	Volumetric soil moisture content (%) from willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at 0-450 mm depth at Moginie, Manawatu. The line bars show the SEM for each treatment. -----	86
Figure 3.9	Relationship between soil moisture at 0-150 mm depth and NHA rate (Kg DM ha ⁻¹ month ⁻¹) from September-April. Moisture data is the mean for each month from willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at Moginie, Manawatu. -----	87
Figure 4.1	Monthly rainfall (mm) for Massey's Riverside Farm, near Masterton, Wairarapa. Data were obtained from Wairarapa College Air Quality Site (Coord NZMG 1949 E2732767 N6024886) -----	115
Figure 4.2	Net herbage accumulation (NHA) rate from November 2004 to October 2005 in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes a significant difference between treatments at P < 0.05. -----	118
Figure 4.3	Cumulative herbage accumulation rate (kg DM ha ⁻¹ month ⁻¹) from November 2004 to July 2005 in willow pasture (WP) treatment at Riverside Farm near Masterton, Wairarapa. Line bars show the SEM. -----	119
Figure 4.4	Leaf area difference (m ² tree ⁻¹) between browsed (1st year) and unbrowsed (2nd year) willow coppice trees in willow pasture (WP) treatment at Riverside Farm near Masterton, Wairarapa. Line bars show the SEM for each date. * denotes P < 0.05 significance. -----	121
Figure 4.5	Linear relationship between willow leaf area (m ² tree ⁻¹) and understory net herbage accumulation rate (kg DM ha ⁻¹ day ⁻¹) in willow pasture (WP) treatment from December to February at Riverside Farm near Masterton, Wairarapa. -----	121
Figure 4.6	Volumetric soil moisture content (%) at 0-150 mm depth at Massey's Riverside Farm near Masterton, Wairarapa. Line bars show the SEM for each date. * denotes significance at P < 0.05. -----	123

Figure 4.7	Volumetric soil moisture content (%) at 0-300 mm depth in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm near Masterton, Wairarapa. Line bars show the SEM for each date. * denotes significance at $P < 0.05$. -----	124
Figure 4.8	Volumetric soil moisture content (%) at 0-450 mm depth in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm near Masterton, Wairarapa. Line bars show the SEM for each date. * denotes significance at $P < 0.05$. -----	125
Figure 4.9	Relationship between mean monthly volumetric soil moisture content (%) in the 0-150 mm depth and NHA rate ($\text{kg DM ha}^{-1} \text{ month}^{-1}$) from November 2004 to March 2005 in willow pasture (WP) and open pasture (OP) treatments at Riverside Farm near Masterton, Wairarapa. -----	126
Figure 5.1	Total monthly rainfall (mm) and mean air and soil temperature ($^{\circ}\text{C}$) during the experimental period in Palmerston North, New Zealand. -----	157
Figure 5.2	Volumetric soil moisture content (%) at 0-150 mm depth in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment * denotes significance at $P < 0.05$. -----	158
Figure 5.3	Volumetric soil moisture content (%) at 0-300 mm depth in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	159
Figure 5.4	Volumetric soil moisture content (%) at 0-450 mm depth in open pasture (OP) and tree pasture (TP) treatments at Moginie, manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	160
Figure 5.5	Event-wise surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	162
Figure 5.6	Total suspended solids (TSS) concentration (mg L^{-1}) in surface runoff in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	163

Figure 5.7	Inorganic suspended solids (ISS) concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	164
Figure 5.8	Organic suspended solids (OSS) concentrations (mg L^{-1}) in surface runoff in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	164
Figure 5.9	Total suspended solids (TSS) losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	166
Figure 5.10	Inorganic suspended solids (ISS) losses (kg ha^{-1}) in surface runoff from OP and TP treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	167
Figure 5.11	Organic suspended solids (ISS) losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. • denotes significance at $P < 0.05$. -----	167
Figure 5.12	Total phosphorus losses (kg ha^{-1}) in surface runoff in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	171
Figure 5.13	DRP concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	172
Figure 5.14	Relationship between total P and DRP concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) treatment at Moginie, Manawatu. (n = 38). -----	173
Figure 5.15	Relationship between total P and DRP concentrations (mg L^{-1}) in surface runoff (mm) in tree pasture (TP) treatment at Moginie, Manawatu. (n = 76). -----	173

Figure 5.16	DRP losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	175
Figure 5.17	TN concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	176
Figure 5.18	TN losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	177
Figure 5.19	NH_4^+ -N concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	178
Figure 5.20	NH_4^+ -N losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	180
Figure 5.21	NO_3^- -N concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	181
Figure 5.22	Total NO_3^- -N losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	182
Figure 5.23	DRP concentrations (mg L^{-1}) in soil water samples taken from suction cups in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment.-----	185
Figure 5.24	Event-wise surface runoff (mm) in open pasture (OP) and tree pasture (WP) treatments at Riverside Farm, Masterton. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	186

Figure 5.25	Total suspended solids (TSS) concentrations (mg L^{-1}) in surface runoff in open pasture (OP) and willow pasture (WP) treatments at Riverside Farm, Masterton. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	187
Figure 5.26	Total suspended solids (TSS) losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	188
Figure 5.27	Total P concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and willow pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	189
Figure 5.28	Total P losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and willow pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. -----	190
Figure 5.29	DRP concentrations (mg L^{-1}) in surface runoff in open pasture (OP) and tree pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	191
Figure 5.30	DRP losses (kg ha^{-1}) in surface runoff in surface runoff (mm) in open pasture (OP) and tree pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	191
Figure 5.31	TN concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and willow pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	192
Figure 5.32	TN losses (kg ha^{-1}) in surface runoff in open pasture (OP) and tree pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	193

Figure 5.33 NH_4^+ -N concentrations (mg L^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. ----- 194

Figure 5.34 NH_4^+ -N losses (kg ha^{-1}) in surface runoff (mm) in open pasture (OP) and tree pasture (WP) treatments at Riverside Farm, near Masterton, Wairarapa. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. ----- 195

List of Plates

Plate 2.1	Tangoio willow growing at the Moginie block, Massey University, Palmerston North, New Zealand. -----	17
Plate 2.2	Kinyoyanagi willow growing at Massey University's Riverside farm, Near Masterton, New Zealand. -----	18
Plate 2.3	Veronese poplar growing at the Moginie block, Massey University, Palmerston North, New Zealand. -----	19
Plate 2.4	Dudely poplar growing at Moginie block, Massey University, Palmerston North, New Zealand. -----	20
Plate 2.5	Stabalising hillslopes with poplar in New Zealand. -----	22
Plate 2.6	Willow fodder block ready to be grazed during summer drought period at Massey University's Riverside Farm near Masterton, New Zealand. -----	24
Plate 3.1	(a) A view of the experimental site with willow and poplar stakes planted at Moginie, Manawatu. (b) Early sprouting and growth (December, 2004) of willow and poplar at Moginie, Manawatu. -----	72
Plate 3.2	(a) Willow shoots at the end of first year growth in willow pasture (WP) treatment at Moginie, Manawatu. (b) Poplar shoots at the end of first year growth in poplar pasture treatment (PP) at Moginie, Manawatu. -----	79
Plate 3.3	(a) View of poplar defoliation on May 29, 2006 in poplar pasture (PP) treatment at Moginie, Manawatu. (b) View of willow defoliation on May 29, 2006 in willow pasture (WP) treatment at Moginie, Manawatu. -----	80
Plate 3.4	(a) Poplar leaf sprouting as on 30th September 2006 in poplar pasture (PP) treatment at Moginie, Manawatu. (b) Willow leaf sprouting as on 30th September 2006 in willow pasture (WP) treatment at Moginie, Manawatu. -----	80
Plate 3.5	(a) Irrigation in willow pasture (WP) and poplar pasture (PP) treatments using sprinkler system during drought in summer 2005 at Moginie, Manawatu. (b) Willow gall saw fly (<i>Pontania proxima</i>) attack (reddish brown galls) on leaves during summer at Moginie, Manawatu. -----	90

Plate 4.1	Willow gall sawfly attack on Kinuyanagi willow (<i>S. viminalis</i>) leaves in willow pasture (WP) treatment at Riverside Farm near Masterton, Wairarapa. -----	120
Plate 5.1	(a) View of a tree pasture (TP) plot showing plot borders with wooden boards, down pipe and pit during setup for runoff measurement at Moginie, Manawatu. (b) View of a tipping bucket fixed in a frame and mounted on a slab in the pit at Moginie, Manawatu. -----	153
Plate 5.2	(a) Trench excavated to discharge surplus water from the main pit at Riverside Farm. Novaflow pipe is ready to be laid into the trench. (b) Spouting laid into the trench at the bottom end of a tree pasture (TP) plot at Moginie, Manawatu. -----	154
Plate 5.3	(a) Tree damage from summer sheep grazing. Broken and bent branches are visible in a tree pasture (TP) plot. (b) Main tree stem broken from summer sheep grazing in a tree pasture (TP) plot at Moginie, Manawatu. -----	183
Plate 5.4	(a) Day time sheep camping in a tree pasture (TP) plot in summer at Moginie, Manawatu. (b) Early winter sheep grazing in a tree pasture (TP) plot at Moginie, Manawatu. -----	184
Plate 5.5	(a) A open pasture (OP) plot after winter sheep grazing. Dung and urine spots are not visible at Moginie, Manawatu. (b) A tree pasture (TP) plot after winter sheep grazing. Dung and urine spots are visible at Moginie, Manawatu. -----	184

List of Appendices

Appendix 3.1	Volumetric soil moisture content (%) at 150-300 mm depth in willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at Moginie Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	233
Appendix 3.2	Volumetric soil moisture content (%) at 300-450 mm depth in willow pasture (WP), poplar pasture (PP) and open pasture (OP) treatments at moginie Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	233
Appendix 4.1	Volumetric soil moisture content (%) at 150-300 mm depth in willow pasture (WP), and open pasture (OP) treatments at Riverside Farm. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	234
Appendix 4.2	Volumetric soil moisture content (%) at 300-450 mm depth in willow pasture (WP), and open pasture (OP) treatments at Riverside Farm. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$.-----	234
Appendix 5.1	Volumetric soil moisture content (%) at 150-300 mm depth in open pasture (OP) and tree pasture (TP) treatments at moginie, manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	235
Appendix 5.2	Volumetric soil moisture content (%) at 300-450 mm depth in open pasture (OP) and tree pasture (TP) treatments at Moginie, Manawatu. Line bars show the SEM for each treatment. * denotes significance at $P < 0.05$. -----	235

Glossary and Abbreviations

ANOVA: analysis of variance.

Blanket forestry: plantation of whole farm with a single species, in New Zealand this usually refers to *Pinus radiata*.

°C: centigrade.

cm⁻³: per cubic centimetre.

cm: centimetre.

Canopy: the part of a tree consisting of branches and foliage.

DAP: diammonium phosphate.

DM: dry matter.

d.p: decimal point.

DOP: dissolved organic phosphorus.

DIP: dissolved inorganic phosphorus.

DRP: dissolved reactive phosphorus.

Densely planted: in this thesis it refers to pastureland planted with >6000 trees per hectare.

dia: diameter.

g: gram.

GLM: general linear model.

h: hour.

ha: hectare.

ISS: inorganic suspended solid.

Kg: kilogram.

L: litre.

LSD: least significant difference.

m: metre.

mg: milligram.

mm: millimetre.

N: nitrogen.

NHA: net herbage accumulation.

NH₄⁺-N: ammonical-nitrogen.

NO₃⁻-N: nitrate-nitrogen.

OP: open pasture.

OSS: organic suspended solid.

P: phosphorus.

PAR: photosynthetically active radiation.

PP: poplar pasture....pastureland planted with poplar trees
RCB: refers to experimental design as randomised complete block
SAS: statistical analysis system.
SEM: standard error of mean.
SSP: single superphosphate.
TKN: total kjeldahl nitrogen.
TN: total nitrogen.
TP: tree pasture...pastureland planted with trees.
TP: total phosphorus.
TSS: total suspended solid.
WP: willow pasture....pastureland planted with willow trees
yr: year.
yrs: years.
Zone 3: the centre of four nuclei trees planted at a square grid.
 θ : soil moisture content.
 ω : soil volume at a given depth.
 μg : microgram.