

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Massey University Library

Thesis Copyright Form

Title of thesis: An investigation of the agronomic value of fine grinding and granulating reactive phosphate rocks

- (1) (a) I give permission for my thesis to be made available to readers in the Massey University Library under conditions determined by the Librarian.
- (b) I do not wish my thesis to be made available to readers without my written consent for _____ months.
- (2) (a) I agree that my thesis, or a copy, may be sent to another institution under conditions determined by the Librarian.
- (b) I do not wish my thesis, or a copy, to be sent to another institution without my written consent for _____ months.
- (3) (a) I agree that my thesis may be copied for Library use.
- (b) I do not wish my thesis to be copied for Library use for _____ months.

Signed

S. J. Offier

Date

11/2/90

The copyright of this thesis belongs to the author. Readers must sign their name in the space below to show that they recognise this. They are asked to add their permanent address.

NAME AND ADDRESS

DATE

AN INVESTIGATION OF THE AGRONOMIC VALUE
OF FINE GRINDING AND GRANULATING
REACTIVE PHOSPHATE ROCKS.

*A thesis presented in partial fulfilment
of the requirements for the degree of
Master of Horticultural Science
in Soil Science at Massey University
New Zealand.*

Sally Jane Officer

1989

ERRATA

Page	Paragraph	Line	Error	Correction
Whole Text			p ³²	³² p
IV	3	8	K ₂ PO ₄	K ₂ HPO ₄
IV	3	16	cylindar	cylinder
XVI		32	made a	made using a
6	3	3	which the	which form the
10	3	4	mutually one	mutually replace one
14	1	6,7	fluorine	fluoride
16, 17, 18	all three pages		DGdo	ΔG ^o
"			H ⁺	H ⁺
"			PO ₄ ³⁻	PO ₄ ³⁻
"			CO ₃ ²⁻	CO ₃ ²⁻
"			F ⁻	F ⁻
"			H ₂ PO ₄ ⁻	H ₂ PO ₄ ⁻
"			CO ₂	CO ₂
"			H ₂ O	H ₂ O
"			DG _{ro}	ΔG _r ^o
"			DG _{no}	ΔG _n ^o
17	2	1	Chien (1977b)	Chien (1977a)
17	3	1	Dole	Doll
17	3	6	6.36	6.3δ
17	3	6	where b	where δ
17	3	7	CO ₃	CO ₃ ²⁻
17	3	9	pH ₂ PO ₄	pH ₂ PO ₄
17	3	10	on b	on δ
17	4	1	if b = 0	δ = 0
17	4	1	if b = 1	δ = 1
22	4	11	subtitution	substitution
23	2	2	relationship(9)	relationship(10)
24	2	4	applied the	applied to the
24	2	8	in thesis	in this thesis
25	1	6	vsolvent	mlsolvent
29	3	4	CaH ₂ PO ₄ ⁻	CaH ₂ PO ₄ ⁺
30	3	2	redrawn	taken
31	1	2	days	days (Mt/Ma)
33	2	4	Nye did	Nye (1986d) did
35	4	2	Nye predicted	Nye (1986d) predicted
36	1	1	Nye tested	Nye (1986b) tested
38	2	9	640 lbsP/acre	716.8 kg P/ha
39	1	5	Nye,	Nye (1986d)
39	1	7	of Kirk and Nye the	Kirk and Nye (1986d)
39	2	1	Fried were	Fried (1958) were
39	2	8	Chin were	Chin (1974) were

39	2	9	Fried,	Fried (1958),
39	3	9	Nye, combined	Nye (1986d) combined
40	2	12	Nye appeared	Nye (1986d)
40	3	5	Nye,	Nye (1985a,b, 1986 a-d)
43	1	7	as predicted	to that predicted
51	4	1	x _B	δ
58	1	9	formic solubility	formic acid solubility
59	2	1	North	The north
61	4	6	x _B	δ
84	1	1	was a significant effect of time	was no significant effect of treatment
84	1	2	and no	and a
84	1	3	time on	treatment on
84	1	4	time effect	treatment effect
84	1	5	of P ³²	of P ³¹
89	2	1	equation 6	equation 7
89	2	3	equation 4	equation 6
123	2	title	segential	sequential
137	1	5	K ₂ HPO ₄	K ₂ PO ₄
138	1	8	106 mgP/pot(106 mgP/pot (where the x axis is the mg P dissolved per pot)
142			K ₂ HPO ₄	K ₂ HPO ₄
175	1	14	in the ratio to formic acid volume basis	ratio of phosphate rock to extractant solution
193	equation 3		Division line missing over P ³² c + P ³² ex	
193	equation 4, upperline		P ³² c.Pex	Pc.P ³² ex
194	equation 5		-P ³² c-	Pc
194	equation 7		-P ³² c-	Pc
194	equation 8		-P ³² c-	Pc

ABSTRACT

The future trends in the use of reactive phosphate rocks in New Zealand may be dependent on improving the handling characteristics of these fine sand- and powder-like materials. Granulation of these materials has been suggested as one option. The effect of fine grinding and granulating reactive phosphate rocks on their agronomic performance was evaluated using a range of phosphate rocks, in laboratory studies and in field and glasshouse trials.

North Carolina, Arad, Sechura and White Youssafia phosphate rocks, in forms normally imported into New Zealand (sand sized material, majority <2mm particle size), were characterised in terms of origin, composition, particle size, and solubility in 2% formic acid. In a 30 minute formic acid extraction of the imported material, White Youssafia phosphate rock at 44% solubility was found to be less reactive than the other phosphate rocks, which ranged from 47% to 55% in formic solubility

In preliminary field trials a very finely ground North Carolina phosphate rock (100% <42 μ m particle size) was granulated with K₂SO₄. The ungranulated phosphate rock, and granules of 0.5-1mm, 1-2mm and 2-4mm diameter, were evaluated on permanent pasture on the Tokomaru silt loam, using an inverse isotopic dilution technique in which the field soil, at the 1.5-6cm depth, was uniformly labelled with P³² by a novel injection method. No plant yield response to fertiliser was observed but significant differences in herbage phosphate content and specific activity indicated a phosphate uptake response to fertiliser. Despite careful selection of areas of sward which had a similar plant content and vigour, the large variability in data from replicate treatments limited the amount of information which could be drawn from the results but the data indicated that the agronomic performance of the finely ground North Carolina phosphate rock was not limited by granulating to 0.5-1mm (mini-granules).

A further range of granulation agents, including neutral salts, organic and mineral acids, their salts, and tallow, were tested for their ability to form strong mini-granules from unground North Carolina phosphate rock. The best granulation agent was a 1:0.6 mixture of citric acid and magnesium sulphate, producing 0.5-1mm mini-granules which had an arbitrary crushing strength of 0.8kg/granule. The production of mini-granules involved pre-drying a phosphate rock/granulation agent slurry until it was just unsaturated, followed by cutting the wet mix through a 0.710mm sieve, granulation at high speed for 30 seconds, and drying of the granules at 80°C for approximately 2 hours. This granulation process was then used to manufacture granules from unground Sechura and Arad phosphate rocks, as well as ground North Carolina and Arad phosphate rocks. Ground North Carolina phosphate rock was also granulated using tallow, by melting the fat and mixing in the phosphate rock, followed by setting the mix in a mould.

Granulated materials, including a commercially prepared product ("Hyphos"), and ungranulated phosphate rocks (including White Youssafia), were evaluated in a glasshouse pot trial. The fertiliser was applied to the surface of pots of established "Nui" perennial ryegrass, with 7 harvests over three and half months.

In general, at the common application rate of 60kgP/ha, the phosphate rock materials were never more than 70% as effective as mono calcium phosphate. The yields derived from unground, ungranulated Sechura, North Carolina, Arad and White Youssafia phosphate rocks were similar, the only significant difference being that the yield derived from Sechura phosphate rock was greater than the yield derived from North Carolina phosphate rock.

The effect of mini-granulation on agronomic performance varied with the type and particle size of the phosphate rock used to make the granules. For example, mini-granulation of "as received" North Carolina and Sechura phosphate rocks caused no reduction in phosphate

availability from these materials, however, mini-granulated "as received" and works ground Arad phosphate rock caused a significant reduction in phosphate availability.

The agronomic performance of North Carolina phosphate rock was improved by grinding to less than 250 μ m in particle size but no further improvement occurred if the phosphate rock was more finely ground (<42 μ m particle size). The agronomic performance of Arad phosphate rock was not improved by grinding.

The sequential fractionation of soil phosphate (1MNaOH followed by 1MHCl) indicated that only approximately 8% of the works ground North Carolina phosphate rock fertiliser had dissolved in the soil at the 5th harvest (10 weeks). A comparison of yields derived from pots fertilised with different rates of K₂HPO₄ sprayed onto chromite (which had a similar particle size distribution to the unground phosphate rocks) indicated that the dissolved phosphate in the soil from the phosphate rock had a similar agronomic value to the K₂PO₄. The low amount of phosphate rock dissolution and the absence of increased of yield response when works ground North Carolina phosphate rock was applied to soil at rates greater than 40 kgP/ha indicated that soil factors were limiting the dissolution of phosphate rock in this experiment. The extent of the limitation varied depending on the phosphate rock type and also the type of pot used (the black polythene bag used for the majority of treatments was enclosed in a galvanised steel cylinder^o for an inverse isotopic dilution experiment). The variable effects of grinding and granulation were attributed to the limitation of the phosphate rock dissolution.

The type of granulation agent (including partial acidulation) had no significant effect on the agronomic performance of the granulated materials, except when tallow was used as a granulation agent and reduced the availability of works ground North Carolina phosphate rock. Unground White Youssafia phosphate rock requires further testing under more rigorous conditions before conclusions can be made about its agronomic availability.

Two isotopic techniques were utilised in the glasshouse experiment in an attempt to quantify the extent of phosphate rock dissolution in the soil. The surfaces of some phosphate rock treatments were sprayed with a carrier free solution of P^{32} , and the inverse isotopic dilution technique used in the field was used again on some treatments.

The use of labelled $K_2HP^{32}O_4$ as a control for the surface labelled experiment provided sufficient information to allow differentiation of phosphate in the plant which was derived from soil and the fertiliser but the model developed could not be directly applied to results from the phosphate rock treatments. The dissolution of different forms of phosphate rock could not be compared using this labelling technique.

The inverse isotopic dilution technique was re-evaluated in the glasshouse trial, by uniformly injecting the pots of ryegrass with a carrier free P^{32} solution. The fertiliser treatments unpredictably stimulated uptake of labelled soil phosphate, so that the changes in herbage specific activity provided little meaningful information.

These two unsuccessful attempts to derive quantitative information from the introduction of the P^{32} isotope into the phosphate rock-soil-plant system demonstrated the difficulties involved in using isotopic dilution techniques to examine phosphate rock dissolution in field soils.

ACKNOWLEDGEMENTS

I would like to gratefully acknowledge the patient support and guidance of my supervisors, Dr M.J. Hedley and Dr N.S. Bolan.

I would like to thank Mr M. Eggels, Mrs A. Rouse, Dr S. Rajapakse, and particularly Mr P. Phommasone, for their technical help.

Thank you to Mr I.J. Barnes, Ms R. Phillips, Mr A.G. Rasmussen and Miss L.K. Heng for their proofreading, and all my family and friends for their support and encouragement.

*The Lord is my strength
and my shield;
My heart trusts in him, and
I am helped;
Therefore my heart exults*

CONTENTS

	Page No.
ABSTRACT.....	ii
ACKNOWLEDGEMENTS.....	vi
TABLE OF CONTENTS.....	vii
LIST OF FIGURES.....	xv
LIST OF TABLES.....	xvii

CHAPTER 1

INTRODUCTION.....	1
-------------------	---

CHAPTER 2

LITERATURE REVIEW

2.1 The Geology of Phosphate Rock	
2.1.1 World deposits.....	4
2.1.2 The formation of sedimentary phosphate rock deposits...6	
2.1.3 Specific conditions for the formation of francolite deposits.....	7
2.1.4 Post precipitation processes.....	9
2.2 Chemical Characteristics of Phosphate Rock	
2.2.1 Characteristics of the apatite group.....	10
2.2.2 Sedimentary apatite.....	10
2.2.3 The association of fluorine with carbonate.....	12
2.2.4 Modelling the composition of <u>C</u> arbonate-fluor-apatite..	15
2.2.5 Further possible substitution into carbonate-fluor- apatite.....	16
2.2.6 The effect of carbonate substitution on reactivity....	16
2.2.7 The composition of <u>F</u> rancolite ore.....	18
2.2.8 The effect of variable substitution on reactivity.....	21

2.3	The Availability of Phosphate Rocks in Agronomic Systems.	
2.3.1	Formic solubility and agronomic availability.....	23
2.3.2	Factors influencing the dissolution of phosphate rock in a natural soil environment: The Kirk and Nye mechanistic model.....	28
	(a) <i>The influence of phosphate rock composition on dissolution.....</i>	30
	(b) <i>The influence of particle size and application rate on phosphate rock dissolution.....</i>	32
	(c) <i>Soil and plant factors influencing phosphate rock dissolution.....</i>	33
2.3.3	The dissolution model and experimental measurements of phosphate rock dissolution.....	35
	(a) <i>The diffusion assumption.....</i>	36
	(b) <i>The soil chemical characteristics++++.....</i>	36
	(c) <i>Particle size.....</i>	38
	(d) <i>Application rate.....</i>	40
	(e) <i>Environmental factors.....</i>	40
2.3.4	The question of granulation.....	42
2.4	Summary and conclusions.....	44
2.5	Objectives of the research programme.....	47

CHAPTER 3

GENERAL EXPERIMENTAL TECHNIQUES AND METHODS

3.1	Characterisation of phosphate rocks	
3.1.1	Sieving.....	48
3.1.2	Grinding.....	48
3.1.3	Total phosphate content.....	48
3.1.4	Water soluble phosphate.....	49
3.1.5	Formic acid soluble phosphate.....	49
3.1.6	pH.....	49
3.1.7	Granule crush strength.....	50

3.2	Dry matter analysis	
3.2.1	Root extractions.....	50
3.2.2	Digestion of herbage for total phosphate analysis.....	50
3.2.3	Analysis of digest by auto analyser for phosphate.....	51
3.3	Radioactive phosphate technique	
3.3.1	Isotope preparation.....	51
3.3.2	Soil injection.....	53
3.3.3	Cerenkov counting method.....	53
3.3.4	Quench correction.....	54
3.4	Soil description.....	54
3.5	Soil analysis	
3.5.1	Water content.....	56
3.5.2	Water extractable phosphate.....	56
3.5.3	Bicarbonate extractable phosphate.....	56
3.5.4	Soil phosphate fractionation.....	56
3.6	Statistical methods.....	57

CHAPTER 4

THE ORIGINS AND CHARACTERISTICS OF FOUR PHOSPHATE ROCKS

4.1	Introduction.....	58
4.2	Information pertaining to the phosphate rocks in existing literature	
4.2.1	North Carolina phosphate rock.....	58
4.2.2	Sechura phosphate rock.....	60
4.2.3	Arad phosphate rocks.....	61
4.2.4	White Youssafia phosphate rock.....	62

4.3	Physical and chemical characteristics	
4.3.1	Introduction.....	62
4.3.2	Source and appearance of the materials	
	(a) <i>Material in the same form as it was when it arrived in New Zealand, ("as received" or AR).....</i>	63
	(b) <i>Factory ground materials (works ground or WG).....</i>	64
4.3.3	Particle size analysis.....	64
4.3.4	Solubility of phosphate rock in dilute formic acid....	66
4.4	Conclusions.....	69

CHAPTER 5

FIELD EXPERIMENT

5.1	Introduction.....	70
5.2	Materials and Methods	
5.2.1	Phosphate rock.....	71
5.2.2	Granulation method.....	71
5.2.3	Field trial design.....	71
5.2.4	Herbage analysis.....	73
5.2.5	Theoretical and methodological background of the "A" value concept.....	73
5.3	Results	
5.3.1	Herbage yield.....	76
5.3.2	Uptake of Phosphate by Pasture.....	77
5.3.3	Uptake of labelled phosphate.....	80
5.3.4	Specific activity.....	83

5.4	Discussion; Agronomic evaluation.	
5.4.1	Evaluation of the "A" value inverse isotopic dilution technique.....	84
5.4.2	Agronomic evaluation.....	88
5.5	Conclusions.....	91

CHAPTER 6

MINI-GRANULATION OF PHOSPHATE ROCK

6.1	Introduction.....	93
6.2	Literature review	
6.2.1	The advantages of granulating phosphate rock.....	93
6.2.2	Suitable granulating agents.....	95
6.3	Materials and methods	
6.3.1	Phosphate Rocks.....	96
6.3.2	Granulation apparatus.....	97
6.3.3	Granulation method.....	97
6.4	Evaluation of granulation agents and granulation method	
6.4.1	Evaluation of granulation agents/ using NCAR.....	99
6.4.2	Evaluation of granulation agents using NCWG.....	101
6.4.3	Refining the granulation method.....	102
	(a) Granulation agent.....	103
	(b) Effect of moisture content.....	103
	(c) Effect of pre-drying.....	103
	(d) Cutting out.....	104
	(e) Granulation.....	104
	(f) Effect of final drying.....	105
6.4.4	Secondary trials using citric acid as a granulation agent.....	105

6.5	Characterization of the granulated materials	
6.5.1	Effect of citric acid as a granulation agent.....	109
6.5.2	Stability to water.....	110
6.5.3	Effect on granulation of different phosphate rocks...	110
6.6	Conclusions.....	111

CHAPTER 7

GLASSHOUSE EXPERIMENT

7.1	Introduction.....	113
7.2	Materials and methods	
7.2.1	Glasshouse trial preparation and maintenance.....	114
7.2.2	Fertiliser treatments.....	116
	(a) <i>Phosphate rock treatments.....</i>	116
	(b) <i>Labelling phosphate rock with P^{32}.....</i>	117
	(c) <i>Controls for the surface labelling technique.....</i>	117
	(d) <i>Inverse isotopic dilution.....</i>	119
7.2.3	Harvesting of herbage and soil sampling.....	120
7.3	Results and discussion	
7.3.1	Dry matter yields produced by the "as received" phosphate rock fertilisers.....	123
7.3.2	Dry matter yields produced by phosphate rock fertilisers of different particle size.....	126
7.3.3	Drymatter yields produced by granulated and ungranulated phosphate rocks.....	127
	(a) <i>Granulated "as received" phosphate rocks.....</i>	127
	(b) <i>Ground phosphate rocks granulated with $MgSO_4$ and citric acid/$MgSO_4$.....</i>	127
	(c) <i>Ground NCPR granulated with H_2SO_4 and tallow.....</i>	131
7.3.4	The effect of granulation agent on dry matter yield..	132

7.3.5	Dry matter yields produced by different fertiliser application rates	
(a)	<i>"Works ground" North Carolina phosphate rock</i>	134
(b)	<i>Potassium dihydrogen phosphate and monocalcium phosphate</i>	137
7.3.6	The effect on dry matter yield of radioactively labelling the surface of the phosphate rock.....	139
7.3.7	Herbage phosphate content (P^{31}).....	139
7.3.8	P^{32} uptake from surface labelled chromite and phosphate rock.....	140
(a)	<i>Chromite controls</i>	140
(b)	<i>labelled phosphate rock treatments</i>	141
(c)	<i>Interpretation of the P^{31} and P^{32} uptake from the surface labelled materials</i>	142
(d)	<i>Discussion</i>	147
7.3.9	Fractionation of soil.....	149
(a)	<i>Recovery of phosphate fertiliser applied to soil</i>	150
(b)	<i>Recovery of fertiliser applied to pots in the surface labelling experiment</i>	151
(c)	<i>Distribution of ^{32}P and ^{31}P in the extractant solutions</i>	153
7.3.10	Fractionation of roots.....	156
7.3.11	Inverse isotopic dilution experiment.....	159
(a)	P^{32} uptake.....	160
(b)	<i>Estimates of the amount of phosphate in the soil which has been released from fertiliser</i>	161
(c)	<i>Estimates of the amount of phosphate in the herbage which was derived from the fertiliser</i>	165
(d)	<i>Fractionation of soil labelled by injection</i>	168
7.4	Conclusions.....	171

CHAPTER 8

SUMMARY OF CONCLUSIONS.....	175
-----------------------------	-----

APPENDIX 1

TREATMENT LIST OF THE GLASSHOUSE EXPERIMENT.....	181
--	-----

APPENDIX 2

EXPERIMENTAL DATA; RATE OF DRY MATTER PRODUCTION.....	184
---	-----

APPENDIX 3

EXPERIMENTAL DATA; PHOSPHATE UPTAKE BY HERBAGE.....	188
---	-----

APPENDIX 4

ALGEBRAIC PROOF FOR SPECIFIC ACTIVITY EQUATIONS.....	191
--	-----

APPENDIX 5

THE PROBLEM OF NEMATODES.....	195
-------------------------------	-----

REFERENCES.....	197
-----------------	-----

LIST OF FIGURES

	Page No.
FIGURE	
2.1: World phosphate reserves and resources, (Stangel, 1978) with the locations of some important deposits.....	5
2.2: The theoretical effects of the most important variables on the dissolution of phosphate rock (carbonate-fluor-apatite) and the subsequent plant uptake of fertiliser phosphate (Kirk and Nye, 1986d).....	31
3.1: A cross sectional diagram of the injector system used to inject soil cores with radioisotope in the inverse isotopic dilution experiments.....	52
5.1: The site of the field experiment, showing the three caged groups of <i>in situ</i> pasture cores which formed the three treatment replicates.....	72
5.2: The uptake of phosphate by fertilised pasture over the duration of the field experiment. Presented as the increase in phosphate taken up per hectare per day above the unfertilised control...78	78
5.3: The P ³² uptake by pasture from injected soil cores (average for treatments) in the field experiment (a), and the resulting herbage specific activity in different treatments over the duration of the experiment (b).....	81
7.1: Selected experimental pots from the glass house experiment demonstrating the plant response after 12 days growth to various fertilisers. The pot type for these treatments was "surface labelled" and the surface of the fertilisers applied to these pots (except MCP) was labelled with P ³²	121
7.2: Selected experimental pots from the glasshouse experiment which were used for the inverse isotopic dilution technique. Note the galvinised metal cylinders around the pots.....	122

FIGURE

- 7.3: The accumulated plant yield (102 days) of ryegrass growing in pots fertilised with various "as received" phosphate rock fertilisers (respectively Sechura, Arad, White Youssafia, North Carolina).....125
- 7.4: The effect of grinding "as received" (AR) North Carolina (NC) and Arad phosphate rock fertilisers to a "works" particle size (WG) on the accumulated yield (102 days) of ryegrass.....125
- 7.5: The effect of mini-granulating (Gr) "as received" North Carolina phosphate rock (NCAR), with either $MgSO_4$ (Mg) or a citric acid/ $MgSO_4$ mixture (CS), on the accumulated yield (102 days) of ryegrass.....128
- 7.6: The effect of mini-granulating (Gr) "as received" Sechura phosphate rock (SAR), with either a citric acid/ $MgSO_4$ mixture (CS), or $MgSO_4$ (Mg), on the accumulated yield (102 days) of ryegrass.....128
- 7.7: The accumulated yield (102 days) of ryegrass grown on pots fertilised with works ground North Carolina phosphate rock (NCWG ungran.), compared with NCWG that had either been partially acidulated, or granulated with either a citric acid/ $MgSO_4$ - mixture (CS), or $MgSO_4$ (Mg), or tallow.130
- 7.8: The effect of mini-granulating "as received" (AR) and "works" ground (WG) Arad phosphate rock, with either a citric acid/ $MgSO_4$ mixture (CS) or $MgSO_4$ (Mg), on the accumulated yield (102 days) of ryegrass.....130
- 7.9: The effect of the granulation agents, $MgSO_4$ (Mg) and a citric acid/ $MgSO_4$ mixture (CS), on the accumulated yield (102 days) of pots of ryegrass fertilised with K_2HPO_4 (5mg P per pot).....133
- 7.10: The accumulated yield (102 days) of ryegrass in the inverse isotopic dilution experiment, from pots fertilised with finely ground (FG) and works ground (WG) North Carolina phosphate rock (NC), and mini-granules (Gr) of NC WG made with $MgSO_4$ (Mg) and mini-granules of "as received" NC made a contaminated source of citric acid (CA).....133

FIGURE

- 7.11: The effect of pot type (inverse isotopic dilution pots or surface labelled treatment pots) on the rate of growth of ryegrass fertilised with works ground North Carolina phosphate rock.....135
- 7.12: The effect of different rates (kgP/ha) of works ground North Carolina phosphate rock fertiliser on the accumulated yield (102 days) of ryegrass.....136
- 7.13: The effect of different rates water soluble phosphate (5, 10, 20 mgP/pot as K_2HPO_4) and MCP (106mgP/pot) on the accumulated yield (102 days) of ryegrass.....136
- 7.14: Comparison of the accumulated (69 days) plant yield from pots fertilised with different rates of North Carolina phosphate rock, different rates of K_2HPO_4 , and also MCP. Also included for comparison are soil fractionation results (taken at 69 days) of the amount of fertiliser which had dissolved from three works ground North Carolina phosphate rock treatments applied at 106mgP/pot.....138
- 7.15: The relationship between the phosphate in plant herbage which was derived from labelled K_2HPO_4 fertiliser (PdfK, equation 1) and the accumulated (69 days) herbage phosphate.....143
- 7.16: The distribution of root mass (g root DM/cm³ soil) at different soil depths in unfertilised pots (control), in pots fertilised with works ground North Carolina (NC) phosphate rock, and in pots fertilised with mini-granules of works ground North Carolina phosphate rock made using $MgSO_4$ (Gran. NCWG+ $MgSO_4$).
.....157
- 7.17: A comparison of the P^{31} uptake (a) and P^{32} uptake (b) in the inverse isotopic dilution experiment.....162
- 7.18: The vertical distribution of P^{32} activity ($\mu CiP^{32}/gsoil$) in a pot from the inverse isotopic dilution experiment which was dissected 69 days after injection of the label into approximately the 1-5cm depth zone.....169

LIST OF TABLES

	Page No.
TABLE	
2.1: The length of the "a" axis dimension of various apatitic minerals.....	14
2.2: Unit cell compositions of a selection of carbonate-fluor-apatites (Chien and Hammond, 1978b).....	15
2.3: Contents of the major elements in a selection of purified apatites and unpurified phosphate rock ores.....	19
2.4: Solubility of the phosphate in a selection of phosphate rocks, using 2% w/w formic acid as an extractant for 60 minutes (Chien and Hammond, 1978b).....	24
4.1: Particle size analysis of the materials used. The results are the amount of material collected on each sieve, expressed as a percentage of the total amount of material recovered from all the sieves.....	65
4.2: Solubility in formic acid of the phosphate rock materials used in this work. The results were found by the standard method (see section 3.1.5). Extractions were carried out for both one hour and half an hour.....	67
5.1: Accumulated plant yield and herbage phosphate 131 days after fertiliser was applied to pasture.....	77
5.2: Estimates of the size of the labile pool in each pot (mgP/pot) and the "A" values derived for fertilisers, at each harvest (mgP/pot).....	86
5.3: PUD and RAID values calculated using accumulated phosphate content of herbage.....	89
6.1: Summary of granulation agents tried with "as received" North Carolina phosphate rock. The proportion of granules in the 0.5-1mm diameter size range is given as an estimated percent of the total yield of granules.....	100

TABLE

6.2:	The proportion of granules in the 0.5-1mm diameter size range is given as an estimated percentage of the total yield of granules.....	102
6.3:	Granulation results using "as received" North Carolina phosphate rock and the full granulation method as stated in section 6.3.3.....	107
6.4:	Extraction of water soluble phosphate from fertilisers granulated with 10% 1:0.6 mix of citric acid and $MgSO_4$	109
6.5:	Analysis of the strength, total phosphate content, binder content (derived from the total phosphate content) and solubility in formic acid (30 minutes) of a range of granulated materials.....	111
7.1:	Particle size analysis of the chromite. The results are expressed as a percentage of the total amount of material recovered from the sieves by a dry brushing method.....	118
7.2:	Dates and intervals of harvesting pot herbage, after the fertiliser application on 25/8/88.....	120
7.3:	Accumulated P^{32} and P^{31} uptake per pot (ten weeks, 5th harvest) from treatments involving water soluble phosphate added to chromite.....	141
7.4:	Calculation of the amount of phosphate rock derived phosphate in the plant available pool, using equation 11. All results are on a per pot basis, accumulated to the 5th harvest.....	146
7.5:	The percentage of fertiliser P^{31} and P^{32} recovered by sequential 1MNaOH and 1MHCl extracts of soil (mean of four replicates).....	150
7.6:	Recovery of fertiliser (% of total applied) based on six replicate measurements. These results are based on the more detailed results presented in 7.7.....	152
7.7:	Extraction of phosphate by 1MNaOH and 1MHCl from fertilised soil samples taken from pots destructively sampled at four weeks and ten after fertiliser application.....	154
7.8:	Distribution of phosphate throughout a ryegrass plant eleven weeks after fertiliser application.....	156

TABLE

7.9:	Recovery of P^{32} from all roots plus shoots harvested from pots (also presented as a percentage of the total P^{32} applied to the pots), and total recovery of P^{32} from plant and soil fractionations, four weeks after fertiliser application.....	158
7.10:	Accumulated (fourteen and a half weeks) dry matter, P^{31} and P^{32} in herbage grown in the inverse isotopic dilution experiment.	160
7.11:	Estimate of the size of the plant available pool in unfertilised pots, and the amount of phosphate released from the fertiliser into the plant available pool, based on the difference in plant available pool size of fertilised and unfertilised pots, as calculated by equation 12.....	163
7.12:	Estimates of the amount of fertiliser phosphate taken up from treatments where the inverse isotopic dilution technique was employed, based on the model developed in Appendix 4.....	166
7.13:	The accumulated amount of fertiliser phosphate in plant herbage in pots in the inverse isotopic dilution experiment as found by two different models (equations 7 yielded PdfPR(1) and equation 12 PdfPR(2)). A comparison is also presented of the actual accumulated P^{32} uptake in the plant herbage and the amount of P^{32} uptake predicted based on the results of PdfPR(1). The difference between the actual and predicted P^{32} uptake is presented as a percentage of the predicted uptake.....	167

CHAPTER 1: INTRODUCTION

Phosphate rock has become the general term for all rocks that contain approximately 2% (or more) phosphorus as orthophosphate (PO_4^{3-}) (Notholt, 1980). The fertilising value of these materials has been recognised for a long time (Morfit, 1873), but usually only in terms of their suitability as raw materials for superphosphate manufacture.

Recently reactive phosphate rocks have been recognised as a distinctive type of phosphate rock which can have a high agronomic performance when applied directly to the soil as fertiliser. These materials have been called "reactive" because they exhibit relatively high solubility in weak organic acid solutions, compared to other phosphate rocks.

The use of reactive phosphate rocks as fertilisers in New Zealand has been limited, but interest in these materials has increased as the cost of superphosphate has risen, and the ore deposits which have been traditionally used for superphosphate manufacture become depleted. There has also been interest generated in reactive phosphate rocks as slow release "natural" fertilisers, more suited to the "organic husbandry" philosophy.

Most reactive phosphate rocks are imported into New Zealand as light, dry, free flowing, sand-like products, which become fine powder when ground. The unground sand sized materials are proving difficult to handle efficiently in current transporting systems. In some cases considerable portions of loads have been lost as the material sifts

through any small openings in the transport (Mr N. Charteris, Ravensdown, pers.comm. 1989). The unground phosphate rocks are also difficult to apply evenly to soil by aerial topdressing in anything but very calm conditions (Gillingham *et al.* 1987), although the small quantities at present being applied allow top-dressing operators to wait for optimum weather conditions (Mr. Bernie Haskill, Jet Spread. pers.comm. 1989). These problems will be accentuated if the phosphate rocks are ground, which is highly probable in view of the strong recommendation to grind these materials to improve their agronomic value. Several researchers have shown that the agronomic performance of these materials is significantly improved by grinding (Khasawneh and Doll, 1978; Rajan *et al.* 1987), and fertiliser regulations in the EEC already require that such materials be ground before application.

The solution to the handling and spreading problems of the reactive phosphate rocks is to granulate these materials, which would also allow the addition of other elements to the fertiliser. Conventional trials of granulated phosphate rocks have found that granulation reduces the amount of phosphate available from these materials (Buchan *et al.* 1970; Chien and Hammond, 1978a), but recent work suggests that on New Zealand pastures, where fertiliser is usually applied to the soil surface, the effects of granulation on agronomic performance may be negligible (Gillion *et al.* 1978; MacKay *et al.* 1981).

The objectives of this thesis are:

1. To examine research literature concerning reactive phosphate rocks, and their use as fertilisers directly applied to the soil (chapter 2), in such a way that a range of rocks can be selected and characterised (chapter 4) for the manufacture of granulated fertiliser.
2. To evaluate methods for the granulation of unground and ground reactive phosphate rocks (chapter 6).
3. To evaluate the effects that grinding and granulation have on the agronomic performance of the selected reactive phosphate rocks (chapters 5 and 7).