

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

INHIBITORY METABOLITES PRODUCTION BY THE
CYANOBACTERIUM *Fischerella muscicola*

A thesis presented in partial fulfilment of
the requirements for the degree of
Doctor of Philosophy in Biotechnology
and Bioprocess Engineering at Massey University.

VISHAL C. SRIVASTAVA

1996

Dedication

This study is dedicated to my Guru
Pandit Sri Ram Sharma Acharya and Vandaniya Mataji.

Without their blessings this study would have been impossible.

ABSTRACT

Studies have been carried out on the production and toxicity of inhibitory metabolites by cyanobacteria.

In the present study a detailed screening system was used to detect antimicrobial substances produced by the cyanobacteria. A range of cyanobacterial species were screened for the production of inhibitory metabolites. However, a majority of the cyanobacterial species screened showed no evidence of production of antimicrobial substances under the culture conditions used. However, *Fischerella muscicola* (UTEX 1829) produced antimicrobial metabolites and this cyanobacterium became the subject of further studies.

Preliminary characterization of both, intracellular and extracellular crude methanolic preparation from the cultures of cyanobacterium *Fischerella muscicola* (UTEX 1829) showed the inhibition of a range of cyanobacteria, eukaryotic algae and eubacteria, using an agar zone inhibition technique.

Using HPLC and GCMS techniques it was possible to demonstrate unique characteristics of these inhibitory metabolites in the crude methanolic cell extracts prepared from cultures of *F. muscicola*. The application of the GCMS technique confirmed the absence of fischerellin from the cultures of *F. muscicola* grown under various laboratory conditions used in these studies.

The effect of culture age on the production and leakage of inhibitory (toxic) metabolites was investigated in batch cultures. The total toxicity of *F. muscicola* was found to reach its maximum during the late exponential or early stationary phase of the growth. The higher toxicity was always found to be associated with the cell extracts rather than the cell-free culture broths indicating that a lower amount of leakage of the toxic material into the culture medium had occurred. However, there was a higher amount of toxicity found in the cell free culture broth extract when the culture was in a state of senescence.

Factorial experiments with the split plot design for the study of effects of various environmental factors such as light intensity, temperature, nitrogen and phosphorus on the production of inhibitory (toxic) metabolites, revealed that low light intensity ($10 \mu\text{mol m}^{-2} \text{s}^{-1}$) together with low temperature (10°C) significantly enhanced the toxicity of *F. muscicola*. Variations in the concentrations of the major nutrients, nitrogen and phosphorus did not have any significant effect on the toxicity of the cyanobacterium over the concentration ranges investigated.

The effect of various environmental factors on the release of inhibitory (toxic) metabolites from the cells of *F. muscicola* was investigated and it was found that the release of metabolites was enhanced under high light intensity ($60 \mu\text{mol m}^{-2} \text{s}^{-1}$) and higher temperature (30°C). However, at low light intensity ($10 \mu\text{mol m}^{-2} \text{s}^{-1}$) and low temperature (10°C) the release was minimal.

Under sets of conditions which generated the highest observed toxicity and the lowest observed toxicity in this cyanobacterium, it was found that growth of the organism was limited by the temperature.

This research contributes to an increased in the understanding of the physiology of *F. muscicola* in particular and to the toxic cyanobacteria in general. Furthermore, this new knowledge will contribute to a deeper understanding of the ecology of these important microorganisms.

Acknowledgements

My greatest thanks goes to Dr Graham Manderson my supervisor who taught me so much and so well. Without his assistance this thesis would still be a muddle of ideas in my mind. My sincere thanks goes to my chief supervisor Prof. Rao Bhamidimarri for his invaluable guidance and encouragement during this study.

John Sykes and Anne-Marrie Jackson have been extremely helpful with my experimental work. Thanks also extended to;

- the rest of the microbiology team, Mike Stevens, Judy Collins and Mike Sahayam.
- the engineering team, John Alger, Bruce Collins and Don McLean.
- the secretarial staff, Glenda Withers, Carol Lewis and Maureen Oemcke.

Special thanks to Dr Daryl Rowan for GC-MS analysis and interpretation.

Thanks also to Dr S. Ganeshnandam for helping with statistical analysis, one of the only people I know who can understand it.

The facility provided by Assoc. Prof. David Fountain and the Plant Biology Department have been very important to the successful completion of this project, as are its staff, Elizabeth Nickless, Edwin Smith, Paul Barret, Stephanie Prince and Carolyn Evans. I would like to thank Dr C.J. O'Kelly for his help during the initial stages of this project.

My appreciation to Ross Davies, John Bronlund, Sirichom Leangon, Lilian Ferreira and Tetsu Matsumura for the excellent job in helping me to prepare this thesis.

I would like to show my appreciation to Vini Natha, Dr Satya Pasumarty and Dulcie Humphrey for their friendship and help while I have been in New Zealand.

Dr Rocky Hudson has performed the difficult task of proof reading this manuscript, the most tedious job of all. Thanks very very much for this and the friendship that you and your husband, Dr Andrew Hudson have given to me over the years.

Finally, I would like to thank my family in India, Mum, Dad, Preeti and Neeraj, for their unfailing support, love and encouragement.

Table of Contents

| | |
|--|--------------|
| Dedication | I |
| Abstract | II |
| Acknowledgements | V |
| Table of Contents | VI |
| List of Tables | XV |
| List of Figures | XVIII |
| Chapter 1 Introduction | 1 |
| Chapter 2 Literature Review | 5 |
| 2.1 CYANOBACTERIA | 6 |
| 2.1.1 Ecology | 6 |
| 2.1.1.1 Factors influencing distribution | 6 |
| 2.1.1.2 Light and its importance | 8 |
| 2.1.1.3 Temperature and growth | 8 |
| 2.1.1.4 pH | 9 |
| 2.1.1.5 Nutrient preferences | 9 |
| 2.1.2 The Toxic cyanobacteria | 11 |

| | |
|---|----|
| 2.1.2.1 Activity and structure of cyanobacterial toxins | 15 |
| 2.1.2.2 Cytotoxins | 15 |
| 2.1.2.2.1 Cyanobacterin | 16 |
| 2.1.2.2.2 Tolytoxin | 18 |
| 2.1.2.2.3 Scytophycins | 19 |
| 2.1.2.2.4 Acutiphycins | 20 |
| 2.1.2.2.5 Hapalindole | 21 |
| 2.1.2.3 Biotoxins | 24 |
| 2.1.2.3.1 Neurotoxins | 26 |
| 2.1.2.3.2 Hepatotoxins | 30 |
| 2.1.2.3.2.1 Microcystins | 30 |
| 2.1.2.3.2.2 Nodularins | 32 |
| 2.1.2.4 Lipopolysaccharides (Endotoxins) | 35 |
| 2.1.2.5 Marine cyanobacterial toxins | 36 |
| 2.1.2.6 Regulation of toxin production | 38 |
| 2.1.2.6.1 Field studies | 39 |
| 2.1.2.6.2 Laboratory studies | 41 |
| 2.1.2.7 Fate and biological significance of cyanobacterial toxins . . | 43 |
| 2.1.2.7.1 Release of toxins into the water | 43 |
| 2.1.2.7.2 Effect of toxins on aquatic life and food chains | 44 |
| 2.1.2.7.3 The biological functions of toxins | 45 |
| 2.1.2.8 Procedures for detection of cyanobacterial toxins | 46 |
| 2.1.3 Aims of this investigation | 49 |

| | | |
|------------------|--|----|
| Chapter 3 | General Materials and Methods | 52 |
| 3.1 | ORGANISMS | 53 |
| 3.1.1 | Cyanobacteria | 53 |
| 3.1.2 | Characteristics of genus <i>Fischerella</i> | 53 |
| 3.1.3 | Chlorophyceae (Green algae) | 56 |
| 3.1.4 | Xanthophycean alga | 56 |
| 3.1.5 | Eubacteria | 56 |
| 3.2 | CULTURE MEDIA | 57 |
| 3.2.1 | For cyanobacteria (fresh water) | 57 |
| 3.2.2 | For cyanobacteria (marine) | 59 |
| 3.2.3 | For chlorophycean and xanthophycean algal cultures | 62 |
| 3.2.4 | For eubacterial cultures | 62 |
| 3.3 | CHEMICALS | 63 |
| 3.4 | STERILIZATION OF CULTURE MEDIA | 63 |
| 3.5 | CLEANING OF GLASSWARE | 64 |
| 3.6 | GROWTH CONDITIONS FOR CYANOBACTERIAL AND ALGAL CULTURE | 64 |
| 3.7 | INOCULUM PREPARATION OF CYANOBACTERIAL CULTURES FOR EXPERIMENTAL FLASKS | 65 |
| 3.8 | GROWTH CONDITIONS FOR BACTERIAL CULTURES | 66 |
| 3.9 | GROWTH MEASUREMENT PARAMETERS | 66 |
| 3.9.1 | Determination of Biomass Concentration | 66 |
| 3.9.2 | Optical density measurement of cyanobacterial growth | 67 |
| 3.9.3 | Cell counting for chlorophycean alga <i>Scenedesmus</i> sp. | 67 |

| | |
|---|-----------|
| | IX |
| 3.9.4 Protein determination | 67 |
| 3.10 OTHER ANALYTICAL METHODS | 68 |
| 3.10.1 Measurement of Dissolved Oxygen | 68 |
| 3.10.2 pH measurement | 69 |
| 3.10.3 Microscopic examination | 69 |
| 3.11 CYANOBACTERIAL TOXICITY TESTING | 69 |
| 3.11.1 Preparation of cyanobacterial samples | 70 |
| 3.11.2 Agar zone diffusion method | 71 |
| 3.11.3 Toxicity testing of crude methanolic cell extracts using liquid culture | 72 |
| 3.12 TOXICITY TESTING OF PURE HEPATO- AND NEURO- TOXIN OF CYANOBACTERIA AGAINST GREEN ALGAE | 73 |
| 3.13 HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC) ANALYSIS OF METHANOLIC CRUDE EXTRACT OF <i>F. muscicola</i> | 74 |
| Chapter 4 Preliminary Screening for Toxin Producing Cyanobacteria . . . | 75 |
| 4.1 INTRODUCTION | 76 |
| 4.1.1 Liquid culture method | 77 |
| 4.1.2 Agar zone diffusion method | 77 |
| 4.2 MATERIAL AND METHODS | 80 |
| 4.2.1 Pure toxins | 80 |
| 4.2.2 Organisms and growth conditions | 80 |

| | |
|---|------------|
| | X |
| 4.2.3 Sample collection and preparation | 80 |
| 4.2.4 Antimicrobial assays | 81 |
| 4.2.5 Growth measurement parameters | 82 |
| 4.2.5.1 Cell Counts | 82 |
| 4.2.5.2 Optical density measurement | 82 |
| 4.2.5.3 Calculation of specific growth rates | 83 |
| 4.3 RESULTS | 84 |
| 4.3.1 Effect of commercial toxins on the target organism | 84 |
| 4.3.1.1 Effect of Anatoxin-a | 84 |
| 4.3.1.2 Effect of Microcystin-lr and -rr on the growth of <i>Scenedesmus sp.</i> | 87 |
| 4.3.2 The screening of NZ isolates of cyanobacteria for toxin production | 91 |
| 4.3.2.1 Shake flask (liquid) culture detection of toxins | 91 |
| 4.3.2.2 An agar diffusion zone assay system for toxin detection . . | 97 |
| 4.3.2.3 Screening of UTEX cultures for toxin production | 98 |
| 4.3.2.3.1 <i>Anabaena flos-aquae</i> (UTEX 1444) | 99 |
| 4.3.2.3.2 <i>Fischerella muscicola</i> (UTEX 1829) | 101 |
| 4.3.2.3.3 <i>Fischerella ambigua</i> (UTEX 1903) | 102 |
| 4.4 DISCUSSION | 106 |
| Chapter 5 An Analysis of Toxic Material from <i>F. muscicola</i> | 110 |
| 5.1 INTRODUCTION | 111 |
| 5.1 MATERIALS AND METHODS | 113 |

| | |
|--|-----|
| 5.1.1 Organisms and growth conditions | 113 |
| 5.1.2 Antimicrobial activity determination | 113 |
| 5.1.3 High Performance Liquid Chromatography (HPLC) of methanolic crude cell extract of <i>F. muscicola</i> | 113 |
| 5.1.3.1 Isocratic run | 113 |
| 5.1.4 Gas Chromatography-Mass Spectrometric determination of <i>fischerellin</i> | 115 |
| 5.1.4.1 Experimental 1. | 115 |
| 5.1.4.2 Experimental 2. | 115 |
| 5.2 RESULTS | 116 |
| 5.2.1 Results from high performance liquid chromatography (HPLC) | 116 |
| 5.2.2 Results from gas chromatography/mass spectrometry (GC- MS) | 120 |
| 5.3 DISCUSSION | 129 |

Chapter 6 Growth and toxin production by *Fischerella muscicola* in batch

| | |
|---|-----|
| culture | 131 |
| 6.1 INTRODUCTION | 132 |
| 6.2 MATERIAL AND METHODS | 135 |
| 6.2.1 Organisms and growth conditions | 135 |
| 6.2.1.1 Culture source | 135 |
| 6.2.1.2 Culture maintenance | 135 |
| 6.2.1.3 Inoculum preparation | 135 |

| | |
|---|------------|
| 6.2.2 Measurement of growth parameters | 135 |
| 6.2.3 pH measurement | 136 |
| 6.2.4 Toxicity measurement | 136 |
| 6.2.5 Presentation of Data | 136 |
| 6.3 RESULTS | 138 |
| 6.3.1 Preliminary determination of growth and toxicity of <i>F.</i> <i>muscicola</i> | 138 |
| 6.3.2 Quantifying Growth of <i>F. muscicola</i> (Choice of Method) . . . | 144 |
| 6.3.3 Growth and toxicity of <i>F.muscicola</i> in batch culture | 147 |
| 6.3.3.1 Extractable Intracellular Toxicity | 147 |
| 6.3.3.2 Specific Toxicity | 148 |
| 6.3.3.3 Diffusible (Supernatant) Toxicity | 150 |
| 6.3.3.4 Supernatant pH and Dissolved Oxygen Concentrations . . . | 150 |
| 6.4 DISCUSSION | 153 |
| | |
| Chapter 7 Effect of Environmental Factors on Toxins Production. | 156 |
| 7.1 INTRODUCTION | 157 |
| 7.2 MATERIAL AND METHODS | 162 |
| 7.2.1 Inoculum preparation | 162 |
| 7.2.2 Sample collection and culture preparation | 163 |
| 7.2.3 Toxicity measurement | 163 |
| 7.2.4 Measurement of growth parameters | 164 |
| 7.2.5 Experimental design | 164 |
| 7.2.6 Analysis of a split plot experiment | 169 |

| | |
|--|---------|
| 7.3 RESULTS | 170 |
| 7.3.1 Effect of light and Temperature | 170 |
| 7.3.1.1 The main effect of temperature on the toxicity | 172 |
| 7.3.1.2 The main effect of light on the toxicity | 173 |
| 7.3.2 The main effect of nitrogen on the toxicity | 175 |
| 7.3.3 The main effect of phosphorus on the toxicity | 175 |
| 7.3.4 Effect of various physical and chemical factor interaction terms on the toxicity | 175 |
| 7.3.4.1 The interaction effects of nitrogen and phosphorus on culture toxicity | 176 |
| 7.3.4.2 Interaction effects of light and nitrogen on culture toxicity | 176 |
| 7.3.4.3 The interaction effects of temperature and nitrogen on culture toxicity | 178 |
| 7.3.4.4 The interaction effects of light, temperature and phosphorus | 180 |
| 7.3.5 The leakage of <i>F. muscicola</i> toxins into the culture medium | 187 |
| 7.3.6 Growth curve of <i>F. muscicola</i> under conditions leading to highest and lowest toxicity in the cyanobacterium | 190 |
| 7.4 DISCUSSION | 192 |
| Chapter 8 General Discussion | 199 |

| | | |
|---------------------------|---|-----|
| Chapter 9 | Conclusions | 208 |
| Chapter 10 | Recommendations | 211 |
| Bibliography | | 214 |
| Appendices | | 246 |
| Appendix A | Standard curve of protein | 247 |
| Appendix B | SAS Programming | 249 |
| Appendix C | Pairwise comparison of LIT*TMP*PH | 253 |

List of Tables

| | |
|--|-----|
| Table 2.1. Countries in which toxigenic cyanobacteria have been reported in Fresh or Marine Waters (adapted from Carmichael, 1989). | 13 |
| Table 2.2. Characteristics of water-soluble toxins from cyanobacteria (Ikawa & Sanser, 1990). | 34 |
| Table 3.1. Composition of Allen's medium (Star & Zeikus, 1987). | 57 |
| Table 3.2. PIV metal solution (Star & Zeikus, 1987). | 58 |
| Table 3.3. Composition of Bold's 3N Bristol's (Star & Zeikus, 1987). | 59 |
| Table 3.4. ES-Enrichment for sea water medium (Star & Zeikus, 1987). | 60 |
| Table 3.5. Composition of PII metal solution (Star & Zeikus, 1987). | 61 |
| Table 3.6. Composition of Brain Heart Infusion (BHI) medium. | 62 |
| Table 4.1. The specific growth rates of <i>Scenedesmus</i> sp. in the presence of Anatoxin-a. | 85 |
| Table 4.2. The specific growth rates of <i>Scenedesmus</i> sp. in the presence of microcystins. | 90 |
| Table 4.3. The specific growth rates of <i>Scenedesmus</i> sp. in the presence of cyanobacterial extracts. | 96 |
| Table 4.4. The specific growth rates of a <i>Scenedesmus</i> sp. growing in the presence of <i>Nodularia</i> extracts. | 97 |
| Table 4.5 Antimicrobial activity of cyanobacterial cell extracts or culture broth tested against green alga <i>Scenedesmus</i> sp. using an agar diffusion method. | 98 |
| Table 4.6. The specific growth rates of <i>Scenedesmus</i> sp. in the presence of extracted material from <i>Anabaena flos-aquae</i> | 101 |

| | |
|--|-----|
| Table 4.7. Inhibition of growth of different indicator organisms by <i>F.musciicola</i> | 104 |
| Table 4.8. Inhibition of growth of different indicator organisms by <i>F.musciicola</i> | 105 |
| Table 5.1. Inhibition of growth of indicator strain <i>Anabaena flos-aquae</i> by HPLC fractions of methanolic crude extracts prepared from the culture of <i>F.</i> <i>musciicola</i> | 117 |
| Table 6.1. Distribution of microcystin during laboratory culture of <i>Microcystis</i> <i>aeruginosa</i> (from Anon., 1990). | 134 |
| Table 6.2. Cytotoxicity of cell extracts of <i>F. musciicola</i> grown in shake flask and tested against the indicator organism <i>A. flos-aquae</i> | 139 |
| Table 6.3. Ratio of Extractable Toxicity (inhibition zone diameter) versus biomass of <i>F. musciicola</i> | 148 |
| Table 6.4. Cytotoxicity of extracts of <i>F. musciicola</i> grown in shake flask and tested against the indicator organism <i>A. flos-aquae</i> | 152 |
| Table 7.1 Values of light and temperature used. | 167 |
| Table 7.2 Concentration of each nutrient factor at their low, original and high levels. | 168 |
| Table 7.3 Arrangement of all possible combinations of nitrogen and phosphorus. | 169 |
| Table 7.4. ANOVA Table showing the overall effect of various factors and their combinations on the toxicity of <i>F. musciicola</i> | 171 |
| Table 7.5. Pair-wise comparison of LS Means of Light and Temperature combinations. | 172 |

| | |
|---|-----|
| Table 7.6. LS Means of light, temperature and phosphorus combinations. | 182 |
| Table 7.7. LSMeans for LIT*TMP*NG*PH showing all possible treatment combinations and their mean toxicity values. | 184 |
| Table 7.8 Mean inhibition zone diameters (Toxicity) of methanol extracted cell free culture broths from <i>F. muscicola</i> grown under different environmental conditions. | 188 |

List of Figures

| | |
|--|----|
| Figure 2.1. Diagrammatic representation of a cyanobacterial cell. CM, Cell membrane; TH, thylakoid; PB1 & PB2, front and side views, respectively, of two rows of phycobilisomes; GG, glycogen granules; CY, cyanophycin granule; P polyphosphate granule; R, 70S ribosomes; C, carboxysome; G, gas vesicle. (adapted from Stanier and Cohen-Bazire, 1977) | 7 |
| Figure 2.2. Molecular structure of the halogenated metabolite "cyanobacterin" produced by the freshwater cyanobacterium <i>Scytonema hofmanni</i> (Gleason & Wood, 1987). | 17 |
| Figure 2.3. Molecular structure of tolytoxin produced by the cyanobacteria <i>Scytonema sp.</i> and <i>Tolypothrix sp.</i> (Patterson & Carmeli, 1992). | 19 |
| Figure 2.4 Molecular structure of scytophycin B. | 20 |
| Figure 2.5. The cytotoxins of <i>Oscillatoria acutissima</i> , namely acutiphycin and didehydroacutiphycin (Barchi <i>et al.</i> , 1984). | 21 |
| Figure 2.6. Molecular structures of hapalindole A and hapalindolinone A and B, the cytotoxins produced by the filamentous cyanobacteria <i>Hapalosiphon fontinalis</i> and <i>Fischerella</i> (ATCC 53558) (Moore <i>et al.</i> , 1984; Schwartz <i>et al.</i> , 1987). | 23 |
| Figure 2.7. Molecular structures for the three lipophilic toxins from <i>Fischerella ambigua</i> : ambigol A (1); ambigol B (2) and tjipanazole D (3). All molecules are bioactive (Falch <i>et al.</i> , 1995). | 25 |
| Figure 2.8. Molecular structure of Anatoxin-a produced by <i>Anabaena flos-aquae</i> (Carmichael, 1992). | 27 |

| | |
|--|----|
| Figure 2.9. Molecular structure for Anatoxin-a(s) of <i>Anabaena flos-aquae</i> and other species of <i>Anabaena</i> (Carmichael, 1994). | 28 |
| Figure 2.10. Molecular structure for Aphantoxins produced by <i>Aphanizomenon flos-aquae</i> (Carmichael, 1992, 1994). | 29 |
| Figure 2.11. Molecular structure for the most commonly occurring microcystin. This is one of a number of toxic peptides produced by species of <i>Microcystis</i> (Kotak <i>et al.</i> , 1995). | 32 |
| Figure 2.12. Three naturally occurring nodularins originally discovered in <i>Nodularia spumegina</i> | 33 |
| Figure 2.13. Structure of debromoaplysiatoxin, lyngbyatoxin A and teleocidin B, all produced by marine cyanobacterium <i>Lyngbya majuscula</i> (Moore, 1981). | 37 |
| Figure 2.14. General scheme of toxin pathways through the food chain (adapted from Ikawa and Sanser 1990) | 50 |
| Figure 3.1. Light micrograph of filamentous cyanobacterium <i>Fischerella muscicola</i> (UTEX 1829), grown in Allen's medium (5 days old culture) at light intensity of $10 \mu\text{mol m}^{-2}\text{s}^{-1}$ and temperature 10°C (400 x magnification). | 55 |
| Figure 4.1. The growth of the target green alga <i>Scenedesmus</i> sp. in the presence of varying concentrations of anatoxin-a. (a) Control flasks for each experimental flask, (b) test flasks which contained up to $20 \mu\text{g/ml}$ of anatoxin-a. | 86 |
| Figure 4.2. The growth of the target green alga <i>Scenedesmus</i> sp. in the presence of varying concentrations of the hepatotoxin microcystin-lr. (a) Control flasks, (b) test flasks which contained the microcystin-lr. | 88 |
| Figure 4.3 The growth of the target green alga <i>Scenedesmus</i> sp. in the presence of | |

| | |
|---|-----|
| varying concentrations of the hepatoin microcystin-rr. | 89 |
| Figure 4.4 Growth of the indicator alga <i>Scenedesmus</i> sp. in Allen's medium containing material from cyanobacterium <i>Scytonema</i> prepared as methanol extracts. (a) No added cell extract, (b) cell extract present. | 93 |
| Figure 4.5 Growth of the indicator alga <i>Scenedesmus</i> sp. in the presence of methanolic crude extracts prepared from <i>Oscillatoria</i> species (a) M193rt and (b) M194rt. | 94 |
| Figure 4.6 Growth of <i>Scenedesmus</i> sp. in Allen's medium, Allen' medium with methanol (+) and Allen's medium containing methanol extract of <i>Nodularia spumigena</i> | 95 |
| Figure 4.7 The growth of <i>Scenedesmus</i> sp. (a) control flasks, (b) methanolic crude extracts prepared from <i>Anabaena flos-aquae</i> | 100 |
| Figure 5.1 HPLC chromatogram (C_{18} isocratic) of the methanolic extract from a culture of <i>F. muscicola</i> showing toxic fractions No. 2 & No. 3 at λ 267nm; mobile phase MeOH:H ₂ O, 99:1 (v/v); flow rate 1 ml/min; 100 μ l injection volume | 118 |
| Figure 5.2 HPLC chromatogram (C_{18} isocratic) of the methanolic extract from a culture <i>F. muscicola</i> showing the presence of absorbance at λ 415nm corresponding with toxic fraction No.2 & 3.HPLC condition as indicated in Fig 5.1. | 118 |
| Figure 5.3 HPLC chromatogram of fraction No. 5 (collected between 7 to 9 min) at λ 267 nm. HPLC conditions as indicated in Fig. 5.1 Injection volumes 10 μ l. | 119 |
| Figure 5.4 HPLC chromatogram of fraction No. 5 (collected between 7 to 9 min) at λ 267 nm and tested for absorbance at λ 415 nm. The peak at 7.89 min is absent. HPLC conditions as indicated in Fig. 5.1. Injection volume 10 μ l. | 119 |
| Figure 5.5 GC-MS chromatogram of toxic fraction No. 3. | 122 |

| | |
|---|---------|
| Figure 5.6 Molecular ion scan of fraction No. 5, using DCI probe method for the detection of a signal at m/z 408. | 123 |
| Figure 5.7 GC-MS chromatogram of fraction No. 5 | 124 |
| Figure 5.8 Molecular ion scan of toxic fraction No. 3, using DCI probe method for detection of a signal at m/z 408. | 125 |
| Figure 5.9 Molecular ion scan of toxic fraction No. 2, using DCI probe method for detection of a signal at m/z 408. | 126 |
| Figure 5.10 GC-MS chromatogram of toxic fraction No. 2. | 127 |
| Figure 5.11 Full scan of toxic fraction No. 2, using DCI probe method for the detection of a signal associated with m/z 408. | 128 |
| Figure 6.1 Toxicity of <i>F. muscicola</i> towards <i>Anabaena flos-aquae</i> | 140 |
| Figure 6.2 Growth curve of <i>F. muscicola</i> | 140 |
| Figure 6.3 Agar plates demonstrating the agar zone diffusion method for toxin production. Growth of <i>Anabaena flos-aquae</i> UTEX 1444 is drawn on Allen's Agar. (a) No cytotoxin in central well, (b), (c), (d), (e) cytotoxin in central well producing a qualifiable zone on inhibition. | 141-143 |
| Figure 6.4 Growth of <i>F. muscicola</i> as measured by (a) dry weight (mg/l). | 145 |
| Figure 6.4b Growth of <i>F. muscicola</i> followed by Protein measurements | 146 |
| Figure 6.5 Standard curve of Optical Density vs Dry weight. Correlation Coefficient (r) =0.94 | 146 |
| Figure 6.6a Growth of <i>F.muscicola</i> as measured by OD changes at wavelength 750nm | 149 |

Figure 6.6b Toxicity of *F. muscicola* culture towards *A. flos-aquae*. 149

Figure 6.7 pH and Oxygen curve of *F. muscicola* culture. 151

Figure 7.1 Diagrammatic representation of experimental runs showing the main
plot factors Light (L) and Temperature (T). 166

Figure 7.2. Toxicity variation at different levels of light and temperature. 174

Figure 7.3 Effect of nitrogen on the toxicity at two levels of light. 177

Figure 7.4. Effect of nitrogen on the toxicity at two levels of temperature. 179

Figure 7.5 Light intensity variation at low temperature (10°C). 183

Figure 7.6 Light intensity variation at high temperature (30°C) 183

Figure 7.7. Toxicity (Inhibition zone diameters) results of *F. muscicola* culture
broths as a response to different environmental growth conditions. 189

Figure 7.8. Growth curve of *Fischerella muscicola* under different environmental
conditions. 191