

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

The impact of stormwater discharges on freshwater, marine water and marine sediments and the implications for environmental management of the Pauatahanui Inlet, Porirua, New Zealand.

**A thesis prepared in partial fulfilment of Masters of Applied Science (Natural
Resource Management), Massey University.**

**Kathryn L. Hooper
19 October 2002**

ABSTRACT

A survey of Cu, Pb, Zn and Cr levels in sediment cores taken at four locations in the Pauatahanui Inlet has been undertaken in conjunction with a freshwater and marine water quality survey.

Levels of heavy metals in sediment were within the Australian and New Zealand Guidelines for Fresh and Marine Water Quality for the protection of Aquatic Ecosystems, [Interim Sediment Quality Guideline (ISQG) (high level of protection)]. Sediments in the vicinity of discharges from suburban catchments showed higher levels of Cu, Pb, Zn and Cr than those in the vicinity of rural catchments.

Levels of Cu, Pb and Zn in streams in the suburban catchments exceeded guideline levels for 99% species protection on occasions, and guidelines for Cu, Pb and Cd were exceeded in marine water. Elevated concentrations of heavy metals in marine and freshwater coincided with rainfall events and increased suspended sediment levels, indicating stormwater discharges as a contributing source of the contaminants.

Levels of N and P exceeded guideline trigger values in freshwater tributaries on occasions. There was no distinct difference between the rural and suburban catchments in terms of nutrient levels detected.

Levels of bacteria present (E-Coli and Enterococci) exceeded NZ guideline levels for contact recreation purposes after rainfall events.

Heavy metals in stormwater were attributed to transport sources. Control of these discharges will likely be 'end of pipe' in nature due to the diffuse nature of the heavy metal inputs, and the difficulties in controlling emissions from vehicles at source.

CONTENTS

1. INTRODUCTION	9
2. RESEARCH PURPOSE	10
2.1 MEETING THE NEED FOR INFORMATION	10
2.2 SCIENTIFIC INFORMATION	11
2.3 PRIOR RESEARCH	12
2.3.1 <i>INLET ECOLOGY</i>	14
2.3.2 <i>URBAN STORMWATER RUNOFF</i>	15
2.3.3 <i>INDUSTRIAL DISCHARGES</i>	20
3. THE STUDY AREA	22
3.1 PORIRUA HARBOUR	22
3.2 PORIRUA ARM	22
3.3 PAUATAHANUI INLET	22
3.3.1 <i>HISTORY OF PAUATAHANUI INLET</i>	25
3.3.2 <i>POPULATION</i>	26
3.3.3 <i>LAND USE</i>	29
3.4 THE RURAL CATCHMENTS	33
3.4.1 <i>HOROKIRI STREAM</i>	33
3.4.2 <i>KAHAO, RATION AND PAUATAHANUI STREAMS</i>	33
3.5 THE SUBURBAN CATCHMENTS	34
3.5.1 <i>DUCK CREEK</i>	34
3.5.2 <i>BROWNS BAY STREAM</i>	35
4. OBJECTIVES	36
5. METHODOLOGY	37
5.1 WATER QUALITY SAMPLING	37
5.2 SEDIMENT CORES	40
5.3 HISTORICAL DATA	40
6. RESULTS	42
6.1 HEAVY METALS IN WATER	42
6.1.1 <i>ZINC</i>	42
6.1.2 <i>LEAD</i>	43
6.1.3 <i>MANGANESE</i>	43
6.1.4 <i>IRON</i>	43
6.1.5 <i>COPPER</i>	44
6.1.6 <i>CHROMIUM</i>	44
6.1.7 <i>CADMIUM</i>	44
6.2 NUTRIENTS IN WATER	45
6.2.1 <i>NITROGEN</i>	45
6.2.2 <i>PHOSPHORUS</i>	45
6.3 PHYSIOCHEMICAL PROPERTIES - WATER	49
6.4 BACTERIA LEVELS	50
6.5 SEDIMENT CORES	52
6.5.1 <i>HEAVY METALS IN SEDIMENT</i>	57
6.5.2 <i>CHROMIUM</i>	62
6.5.3 <i>COPPER</i>	62
6.5.4 <i>LEAD</i>	63
6.5.5 <i>ZINC</i>	63
6.5.6 <i>COMPARISON – GLASBY et al & CURRENT STUDY</i>	64
7. DISCUSSION	65
7.1 A NOTE ON DETECTION LIMITS	65
7.2 HEAVY METALS IN PAUATAHANUI INLET	65
7.3 NUTRIENTS & PHYSIOCHEMICAL STATUS OF PAUATAHANUI INLET	68

7.4	BACTERIA	69
7.5	IMPLICATIONS FOR MANAGEMENT OF THE INLET	70
7.5.1	<i>WHO IS RESPONSIBLE FOR STORMWATER MANAGEMENT IN NEW ZEALAND?</i>	71
7.6	SUSTAINABLE URBAN STORMWATER MANAGEMENT	73
7.6.1	<i>WHAT IS SUSTAINABLE?</i>	73
7.6.2	<i>IS CURRENT MANAGEMENT OF THE INLET SUSTAINABLE?</i>	74
7.7	SCIENCE AND PLANNING	74
7.8	STORMWATER MANAGEMENT OPTIONS	76
7.8.1	<i>PLANNING AND REGULATORY METHODS</i>	76
7.8.2	<i>EDUCATION</i>	76
7.8.3	<i>SOURCE CONTROLS</i>	77
7.8.4	<i>STRUCTURAL SOLUTIONS</i>	78
8.	CONCLUSIONS	80
	BIBLIOGRAPHY AND REFERENCES	82

APPENDICES

Appendix 1	Map Showing Porirua City area units (Statistics NZ)
Appendix 2	Water Quality Data Tables

LIST OF FIGURES

FIGURE 1. LOCATION OF THE STUDY AREA, PAUATAHANUI INLET, NEW ZEALAND.....	24
FIGURE 2. USUALLY RESIDENT POPULATION – PAUATAHANUI INLET CATCHMENT, 1991-2001.	26
FIGURE 3. INCREASE IN TOTAL DWELLINGS (OCCUPIED & UNOCCUPIED) – PAUATAHANUI INLET SURROUNDING AREA, 1991-2001.	27
FIGURE 4. EXPANSION OF URBAN AREAS IN THE PAUATAHANUI INLET CATCHMENT. SOURCE: WELLINGTON REGIONAL COUNCIL, 2000.	30
FIGURE 5. PAUATAHANUI INLET CATCHMENT MAP, FROM HEALY, 1980.....	32
FIGURE 6. THE TWO ARMS OF THE UPPER DUCK CREEK CATCHMENT.....	34
FIGURE 7. WATER QUALITY SAMPLING LOCATIONS – PAUATAHANUI INLET	38
FIGURE 8. SEDIMENT SAMPLING LOCATIONS – PAUATAHANUI INLET	41
FIGURE 9. COMPARISON OF NITROGEN LEVELS IN THE HOROKIRI STREAM (RURAL CATCHMENT) AND DUCK CREEK (URBAN CATCHMENT).	48
FIGURE 10. COMPARISON OF PHOSPHORUS LEVELS IN THE HOROKIRI STREAM (RURAL CATCHMENT) AND DUCK CREEK (URBAN CATCHMENT).	48
FIGURE 11. DUCK CREEK EAST CORE – PHOTO AND DESCRIPTION.....	53
FIGURE 12. DUCK CREEK WEST CORE – PHOTO AND DESCRIPTION.....	54
FIGURE 13. HOROKIRI STREAM EAST CORE – PHOTO AND DESCRIPTION.	55
FIGURE 14. HOROKIRI STREAM WEST CORE – PHOTO AND DESCRIPTION.	56
FIGURE 15. TRACE METAL CONCENTRATIONS IN SEDIMENT AS A FUNCTION OF DEPTH IN CORES FROM EAST AND WEST OF THE HOROKIRI STREAM AND DUCK CREEK DISCHARGE POINTS, IN THE PAUATAHANUI INLET.	61

LIST OF TABLES

TABLE 1. MAIN CONTAMINANTS FOUND IN URBAN RUNOFF	17
TABLE 2. DESCRIPTION OF WATER SAMPLING SITES.....	37
TABLE 3. MAXIMUM HEAVY METAL LEVELS IN FRESHWATER ($\mu\text{g/L}$) OVER 5 SAMPLING RUNS, FEB- MAY 2001 - COMPARISON WITH NEW ZEALAND & CANADIAN GUIDELINE LEVELS.	42
TABLE 4. MAXIMUM HEAVY METAL LEVELS IN MARINE WATER ($\mu\text{g/L}$) OVER 5 SAMPLING RUNS, FEB- MAY 2001 - COMPARISON WITH NEW ZEALAND & CANADIAN GUIDELINE LEVELS.	44
TABLE 5. RANGE OF TOTAL NITROGEN AND PHOSPHORUS LEVELS IN FRESHWATER ($\mu\text{g/L}$) OVER 5 SAMPLING RUNS, FEB-MAY 2001 - COMPARISON WITH NEW ZEALAND GUIDELINE LEVELS.	46
TABLE 6. RANGE OF TOTAL NITROGEN AND PHOSPHORUS LEVELS IN MARINE WATER ($\mu\text{g/L}$) OVER 5 SAMPLING RUNS, FEB-MAY 2001 - COMPARISON WITH NEW ZEALAND GUIDELINE LEVELS.	47
TABLE 7. PHYSIOCHEMICAL PARAMETERS, RANGE IN FRESH WATER OVER 5 SAMPLING RUNS, FEB-MAY 2001 - COMPARISON WITH NEW ZEALAND & CANADIAN GUIDELINE LEVELS.	49
TABLE 8. PHYSIOCHEMICAL PARAMETERS, RANGE IN MARINE WATER OVER 5 SAMPLING RUNS, FEB- MAY 2001 - COMPARISON WITH NEW ZEALAND & CANADIAN GUIDELINE LEVELS.	50
TABLE 9. BACTERIA LEVELS, RANGE IN FRESHWATER OVER 5 SAMPLING RUNS, FEB-MAY 2001 - COMPARISON WITH NEW ZEALAND GUIDELINE LEVELS (CFU/100ML).	51
TABLE 10. BACTERIA, RANGE IN MARINE WATER OVER 5 SAMPLING RUNS, FEB-MAY 2001 - COMPARISON WITH NEW ZEALAND GUIDELINE LEVELS. (CFU/100ML).....	51
TABLE 11. HEAVY METAL LEVELS IN SEDIMENT, MARCH 2001 - COMPARISON WITH NEW ZEALAND AND CANADIAN GUIDELINE LEVELS. LEVELS SHADED GREY = MAXIMUM LEVEL DETECTED IN INLET DURING SURVEY.	59
TABLE 12. HEAVY METAL LEVELS IN SEDIMENT, GLASBY <i>ET AL</i> , 1990- COMPARISON WITH NEW ZEALAND AND CANADIAN GUIDELINE LEVELS. RESULTS OF 30 SAMPLES TAKEN FROM PAUATAHANUI INLET.	60

LIST OF PHOTOS

PHOTO 1. SUBURBAN DEVELOPMENT IN PROGRESS, WHITBY.	28
PHOTO 2. VIEW ACROSS PAUATAHANUI INLET TO THE 'SOUTHERN' (SUBURBAN) SIDE OF THE INLET.	28
PHOTO 3. VIEW ACROSS THE PAUATAHANUI INLET TO THE 'NORTHERN' (RURAL) SIDE OF THE INLET. ...	29
PHOTO 4. SUBDIVISION DEVELOPMENT, WHITBY, 2000.....	31

ACKNOWLEDGEMENTS

This study was commissioned by the Wellington Regional Council (WRC), upon identification of the need for more data about the contaminants in the urban runoff which is discharging into the Pauatahanui Inlet. The significance of this information when identifying environmental pressures on the Inlet is clear. The reason for this study was a need for site specific information upon which to base policy, planning and environmental management decisions. I would like to take this opportunity to thank the WRC for supporting this research.