Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

SPACE AND TIME RELATIONS IN THE
COAT OF THE NEW ZEALAND ROMNEY
LAMB.

NANCY M. GALPIN, M. Sc.

MASSEY AGRICULTURAL COLLEGE,

PALMERSTON NORTH.

NEW ZEALAND.

Massey University Library
New Zealand & Pacific Collection



A.

C.

The Distribution of the Fibre Type Arrays over the Body of the New Zealand Romney Lamb.

- B.

 The Prenatal Development of the Coat of the Romney Lamb.
- Relative Abundance of Fibres of the Halo-Sickle Group on Different Body regions.

MAJOR SECTIONS :

- 1. The Distribution of the Fibre Type Arrays.
- 2. The Prenatal Development.
- 3. The Distribution of the Pre-Curly-tip fibres.
- 4. General Summary.
- 5. Reference.
- 6. Description of Plates.

SECTION A :

THE DISTRIBUTION OF THE FIBRE TYPE ARRAYS OVER THE BODY OF THE NEW ZEALAND ROMNEY LAMB.

- Introduction. 1.
- 11. Material and Methods.
- 111. Distribution of the Fibre Type Arrays.
 - (1) (2) (3)

Escarpment - a new fibre type array.
The Fibre Type Arrays.
The Distribution of the Fibre Type Arrays.

Minor Variations. Summary.

- (4) Minor Variations.
 (5) Summary.
 (6) Prevalence of Several Fibre Type Arrays on different positions.
- Comparative Studies in two finer woolled breeds: Ryeland and Southdown. IV.
 - (1) Southdown.
 - (2) Ryeland.
- The Shedding of Fine Fibres. V.
- Vl. Summary and Discussion.