Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

THE TONSILLAR CARRIAGE OF YERSINIA SPECIES BY PIGS

A THESIS PRESENTED IN PARTIAL FULFILMENT (40%) OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF PHILOSOPHY IN VETERINARY SCIENCE AT MASSEY UNIVERSITY

CLAUDE ADRIAN De.ALLIE

April, 1994

"Let those who labour hold the reins" M.R. Bishop

In memory of a Grenadian Hero and Martyr who endeavoured to uplift the standards of the working class through mass education and social reforms. His spirit lives on!

To my family and children Maurice and Yenifer.

ABSTRACT

The impetus for this study arose due to the increasing isolation of species of *Yersinia* from people, with pigs being suspected as major reservoirs of human pathogenic strains of the organism in New Zealand. The general aims of the study, conducted in two phases, among pigs from several herds sent for slaughter at an abattoir in Palmerston North were:

- (i) to determine the presence of human pathogenic strains of *Yersinia* in the tonsils of slaughtered pigs and their distribution among selected herds,
- to determine the seasonal effect on prevalence of isolation and type of organism isolated, and
- (iii) to determine the *in vitro* virulence characteristics of strains of the organism isolated from the tonsils of slaughter pigs, and their potential public health implications.

The first phase involved a cross-sectional study, conducted between August and September, 1993. Tonsils were collected from 124 pigs from eight farms and were examined for the presence of species of Yersinia. A total of 77 (62.1%) strains of Yersinia were isolated, this consisted of 42 (33.9%), 27 (21.8%), 7 (5.6%) and 1 (0.8%) strains of Y. enterocolitica, Y. pseudotuberculosis, Y. frederiksenii and Y. kristensenii respectively. Yersinia enterocolitica serotypes 0:3, 0:5,27 and Y. pseudotuberculosis comprised 26 (33.8%), 12 (15.6%) and 27 (35.1%) of the total number of isolates respectively. Yersiniae were isolated from all eight farms with individual farm prevalences ranging from 20% to 100%, while the number of species per farm ranged from 1 to 3. The pyrazinamidase activity test correctly identified 48 of the isolates as pathogenic or non-pathogenic yersiniae, (a specificity of 96%).

The second phase, a longitudinal study, was conducted over a period of twelve months (February 1993 - January 1994), among pigs from four farms, selected according to the particular strain of *Yersinia* prevailing in the herd. A total of 705 pigs were examined for the carriage of species of *Yersinia* in their tonsils. A total of 264 isolates were obtained, consisting of 198 (75%), 55 (20.8%), 5 (1.9%), and 1 (0.4%) strains of *Y. enterocolitica*, *Y.*

pseudotuberculosis, Y. intermedia, Y. frederiksenii and Y. kristensenii respectively. Yersinia enterocolitica serotypes 0:5,27 and 0:3 comprised 105 (39.8%) and 78 (29.5%) of the total number of isolates respectively. Yersinia pseudotuberculosis comprised 55 (20.9%) with serotype III, 39 (14.8%) the most consistently isolated serotype.

Yersiniae were isolated throughout the year particularly in the colder months. *Yersinia* enterocolitica serotypes 0:3 and 0:5,27 were found throughout the year with the lowest prevalence in the warmer months. However, a seasonal variation existed among serotypes of *Y. pseudotuberculosis*, with serotypes I and II found only in the winter and spring. Serotype III was found throughout the year, except for February.

During phase two of the study, 150 isolates of Yersinia were tested for in vitro virulenceassociated characteristics. The autoagglutination test, CR-MOX agar, and the pyrazinamidase assay, coupled with salicin and aesculin tests, were highly successful in separating pathogenic from non-pathogenic strains of Y. enterocolitica. Likewise, the three assays successfully identified virulence activity in the majority of strains of Y. pseudotuberculosis with specificity among the three assays ranging between 90-100% for both Y. pseudotuberculosis and Y. enterocolitica.

The study also revealed marked variation in prevalence and type of *Yersinia* species isolated from pigs from different farms. The fact that particular serotypes predominate and persist on specific farms strongly suggest that there are factors such as source of pigs, management parctices or contact with other animals which determine their status. Identification of these determinates could lead to control or eradication of important yersiniae from pig farms.

The overall prevalence of 41.1% ranks New Zealand among countries with reported high isolation rates of the organism and further emphasises the fact that pigs constitute major reservoirs for human pathogenic strains of *Yersinia* worldwide. The infection among slaughter pigs in New Zealand may be of human health concern and this warrants further investigation particularly to determine whether the strains isolated from pigs are identical to those involved in human disease.

ACKNOWLEDGEMENTS

I would like to record my sincere gratitude to the government and people of New Zealand for permitting me to pursue higher studies in Veterinary Science, under the Commonwealth Scholarships and Fellowship Plan.

My research would have been jeopardised, without the excellent supervision of my Chief Supervisor, Professor C R Wilks. Colin is a genius. His accessible personality, motivation, and constructive criticism enabled the termination of my study and my successful stay in New Zealand.

My other supervisors, Mr Per Madie and Mr Stanley Fenwick, contributed significantly throughout the duration of my study. Per's introduction to the working environment at the abattoir, established a solid foundation for carrying out the research. Stan's quest for new strains of *Yersinia* enlightened my interest in working with this organism. There constructive critical comments during editing of the manuscripts has been greatly appreciated.

Several other persons contributed to my learning experience at Massey University.

Associate Professor M R Alley supervised the pathology section of my course and from whom I acquired tremendous skills in Diagnostic Veterinary Pathology.

Professor B W Manktelow, with his vast experience in Veterinary Pathology, and from whom I learnt a lot during weekly departmental tutorials.

The other members of staff in the department, who unselfishly assisted whenever it became necessary.

Dr Alister Johnstone and his staff at the Ministry of Agriculture and Fisheries (MAF) from whom I gained a lot during pathology seminars.

This modest work would not have terminated without the excellent secretarial skills of Mrs Gillian Budge, who typed the manuscripts and Mrs Sheryll Crawford, for her tremendous

assistance in correcting and formatting the manuscripts. Her willingness to assist was greatly appreciated.

My special thanks to Mr Malcolm Rice for his excellent preparation of the graphs. Associate Professor M R Alley for the photomicrographs. Mr Tony Watts for the photographs taken at the abattoir. Mr Peter Wildbore for arranging transportation during the collection of samples. Mrs Jan Schrama for preparation of special microbiological media. Mr N K Pillai for calculation of molar volumes in the preparation of special media.

Finally, to all who contributed with social gestures, which made my stay in New Zealand a memorable one. Not forgetting, Mr Stanley Fenwick, Dr Eammon Gormley and Dr / David Palmer.

To all, my sincere appreciation.

TABLE OF CONTENTS

		Page
ABSTRACT		iii
ACKNOWLEDGEM	ENTS	v
TABLE OF CONTEN	VTS	vii
LIST OF FIGURES		xi
LIST OF TABLES		xii
LIST OF PLATES		xv
GENERAL INTROD	UCTION	xvi
CHAPTER ONE:	REVIEW OF THE LITERATURE	1
SECTION A:	YERSINIOSIS	1
HISTO	DRICAL AND GEOGRAPHICAL ASPECTS	2
GENE	RAL CHARACTERISTICS OF YERSINIA SPECIES	6
	The bacterium	6
	Growth requirements	7
	Isolation	8
	Identification	9
	Biotypes and serotypes	10
EPIDI	EMIOLOGY	13
	Reservoirs	13
	Transmission	14
	Susceptible species	16
PATH	OGENESIS AND CLINICAL PRESENTATION	17
	Yersinia pseudotuberculosis infection	17
	Yersinia enterocolitica infection	18
YERS	INIOSIS IN PIGS	23
	Infections with Y. enterocolitica, Y. pseudotuberculosis	
	and Y. pestis	23
	Anatomical distribution	27
	Seasonal occurrence	29
VFRS	INIA ENTEROCOLITICA IN FOOD HYCIENE	30

	viii
Occurrence in pork, related products and other	
foodstuffs	30
Foodborne outbreaks	22
YERSINIOSIS IN NEW ZEALAND	24
Human infections	24
Animal infections	34
VIRULENCE ACTIVITY	36
Pyrazinamidase Test	37
Calcium dependent growth at 37 °C	38
Congo Red uptake	38
Autoagglutination	38
DIAGNOSTIC METHODS	39
TREATMENT, PREVENTION AND CONTROL	41
Treatment	41
Prevention and control	42
PUBLIC HEALTH IMPLICATIONS	45
SECTION B: THE PORCINE PALATINE TONSILS	47
INTRODUCTION	47
LOCATION AND ONTOGENESIS	47
CLASSIFICATION OF TONSILS	48
MICROSCOPIC ANATOMY	48
PATHOLOGY	49
CHAPTER TWO: CROSS-SECTIONAL STUDY OF THE OCCURRENCE OF	
YERSINIA SPECIES IN THE TONSILS OF SLAUGHTER	
PIGS IN THE MANAWATU	54

54

55

55

55

56

56

INTRODUCTION

MATERIALS AND METHODS

Determination of sample sizes

Slaughter of pigs and sample collection

Processing of samples for cold-enrichment

Cultivation of species of cold-enriched samples

		ix
	Screening for Yersinia species	57
	Identification and serotyping of Yersinia isolates	57
	Virulence assay	58
	Biochemical media used	58
RESU	LTS	60
	Occurrence and Distribution of Yersinia isolates	60
	Biochemical and serological characterization of isolates	60
	Pathogenic Yersinia isolates	61
	Virulence assay	62
DISCU	USSION	62
CONC	CLUSIONS	64
CHAPTER THREE:	THE EFFECT OF SEASON ON PREVALENCE AND	
	SPECIES OF YERSINIA ISOLATED FROM THE	
	TONSILS OF SLAUGHTERED PIGS	66
	INTRODUCTION	66
MATE	ERIALS AND METHODS	67
	Farm selection	67
	Determination of sample sizes	67
	Animal reference	68
	Sampling plan	68
RESU	LTS	68
	Distribution and prevalence of species	
	of Yersinia	72
	Monthly isolation of yersiniae	72
	Yersinia enterocolitica	72

Distribution	and	seasona	l incid
of Yersinia o	n in	dividual	farms

Other Yersinia species

Yersinia psuedotuberculosis

average daily temperatures

Relationship between monthly isolation of Yersiniae and

and seasonal incidence of species

DISCUSSION

75 83

72

74

74

75

CONCLUSIONS

CHAPTER FOUR:	IN VITRO ASSESSMENT OF VIRULENCE AMONG	
	STRAINS OF YERSINIA ENTEROCOLITICA	
	AND YERSINIA PSEUDOTUBERCULOSIS ISOLATED	
	FROM THE TONSILS OF SLAUGHTERED PIGS.	
	PUBLIC HEALTH SIGNIFICANCE OF POTENTIAL	
	PATHOGENS	88
INTR	ODUCTION	88
MAT	ERIALS AND METHODS	90
	Bacterial strains	90
	Growth, isolation and identification procedures	90
	Evaluation of sample tests used to define pathogenic	
	serotypes of Y. enterocolitica	90
	Temperature-dependent autoagglutination	91
	Congo red, magnesium oxalate agar (CR-MOX)	92
	Pyrazinamidase test	92
	Salicin fermentation-aesculin hydrolysis	92
RESU	JLTS	93
	Virulence-associated characteristics in strains of	
	Y. enterocolitica	93
	Virulence-associated characteristics in strains of	
	Y. pseudotuberculosis	94
DISC	USSION	94
CON	CLUSIONS	98
CHAPTER FIVE:	GENERAL DISCUSSION	99
RECO	OMMENDATIONS	103
APPENDICES		109
REFERENCES		135

x 86

LIST OF FIGURES

		Page
Figure 1.1	The proposed route of infection for Yersinia enterocolitica.	15
Figure 1.2	Flow diagram for production and processing of pork products.	44
Figure 3.1	Individual farm prevalence of species of Yersinia isolated from the tonsils of pigs during the period February 1993-January 1994.	69
Figure 3.2	Relationship between monthly isolations of species of Yersinia and average daily temperatures.	76
Figure 3.3	Seasonal distribution and prevalence of Y. pseudotuberculosis serotype I from Farm K.	77
Figure 3.4	Seasonal distribution and prevalence of Y. pseudotuberculosis serotype II from Farm K.	77
Figure 3.5	Seasonal distribution and prevalence of Y. pseudotuberculosis serotype III from Farm K).	78
Figure 3.6	Seasonal distribution and prevalence of Y. enterocolitica serotype 0:3 from Farm T.	80
Figure 3.7	Seasonal distribution and prevalence of Y. enterocolitica serotype 0:5,27 from Farm W.	82

LIST OF TABLES

Table 1.1	Interrelationships between serogroup, biovar, phagevar and geographical distribution of the most common human	
	pathogenic strains of Y. enterocolitica.	5
Table 1.2	Relationship between biotype, serogroups, pathogenicity and geographic distribution of Y. enterocolitica.	6
Table 1.3	Biochemical differentiation of species within the genus Yersinia.	12
Table 1.4	Biochemical differentiation of Y. enterocolitica biogroups.	13
Table 1.5	Cardiovascular infections with Y. enterocolitica.	22
Table 1.6	The prevalence of <i>Yersinia enterocolitica</i> in pigs based on the findings of various investigators.	27
Table 1.7	Summary of some documented outbreaks of Yersiniosis.	33
Table 1.8	Methods used to define virulence in cultures of Yersinia enterocolitica.	37
Table 2.1	Flow diagram used in the isolation of Yersinia species from tonsils.	59
Table 2.2	Occurrence and distribution of Yersinia species from tonsils.	60
Table 2.3	Biochemical and serological characterization of 77 isolates of Yersinia.	61

Table 2.4	Prevalence of pathogenic Yersinia from the tonsils of pigs.	62
Table 2.5	Presence of pyrazinimidase activity (PYZ) in the isolates of <i>Y. enterocolitica</i> and related species.	
Table 3.1	Farm distribution and prevalence of species of Yersinia isolated from the tonsils of pigs during the period February 1993 - January 1994.	70
Table 3.2	Species of Yersinia isolated in the study.	71
Table 3.3	Monthly isolation of species of Yersinia from the tonsils of pigs during the period February 1993 - January 1994.	73
Table 3.4	Serological characterization of 253 strains of Y. enterocolitica and Y. pseudotuberculosis isolated during the study.	74
Table 3.5	Monthly isolation of species of Yersinia from Farm H.	75
Table 3.6	Monthly isolation of species of Yersinia other than Y. pseudotuberculosis from Farm K.	79
Table 3.7	Monthly isolation of species of Yersinia other than Y. enterocolitica serotype 0:3 from Farm T.	81
Table 3.8	Monthly isolation of species of Yersinia other than Y. enterocolitica serotype 0:5,27 from Farm W.	83
Table 4.1	Properties of Y. enterocolitica strains.	91

xiii

Table 4.2	Occurrence of virulence-associated characteristics among	
	strains of Yersinia enterocolitica (Ye) isolated from	
	the tonsils of pigs.	93
Table 4.3	Occurrence of virulence-associated characteristics among	
	serotypes of Yersinia pseudotuberculosis isolated from the	
	tonsils of pigs.	94

xiv

LIST OF PLATES

Plate 1	I	The palatine tonsils of the pig; paired lympho-epithelial organs.	52
Plate]	п	Histologic section of pig's tonsil, showing a tonsillar follicle, consisting of a cypt (C) with its assoicated lymphatic centres (L) H & E stain (x20).	52
Plate 1	III	Histologic section of pig's tonsil, showing invagination of surface epithelium to form a crypt H & E stain (x40).	53
Plate 1	IV	Histological tonsil of pig's tonsil, showing active lymphoid epithelium of a crypt H & E (x60).	53
Plate	V	Porcine tongue, tonsils and pharynx. Location of tonsils are indicated.	65
Plate	VI	Circumanal incision and removal of the intestines.	106
Plate V	VII	Excision of tongue, pharynx and particularly the tonsils.	106
Plate V	VIII	Post-mortem meat inspection involving incision of mandibular lymph nodes.	108
Plate 1	IX	Deboning of head meat.	108

GENERAL INTRODUCTION

Yersiniosis, a zoonotic disease caused by Yersinia enterocolitica and Y. pseudotuberculosis, is now recognised worldwide. The species Y. enterocolitica is an important cause of gastroenteritis in humans, especially in temperate countries (Mollaret et.al., 1979; WHO, 1981, 1987). Yersinia enterocolitica is considered to be a foodborne pathogen, despite the fact that attempts to isolate the bacterium from foods implicated in cases of disease in humans have rarely proved successful. However, some large foodborne outbreaks caused by Y. enterocolitica have been reported in the U.S.A., Canada and Japan (see Table 1.7). Pork products are considered to be the most likely source of infection (Hurvell, 1981; Lee et.al., 1981; Morris and Feeley, 1976), although some aspects of the epidemiology still remain to be clarified.

Pigs appear to constitute an important reservoir for Y. enterocolitica infection (Hurvell, 1981; Kapperud, 1991; Schiemann, 1989), and are the only food animal which regularly harbours pathogenic Y. enterocolitica. Pigs are often healthy carriers of Y. enterocolitica biotype 4/0:3 and biotype 2/0:9 strains which cause disease in humans. Biotype 1B/0:8, the predominant human pathogen in the U.S.A., appears to be rare in pigs. This serotype may have entirely different reservoirs and ecology (Schiemann, 1989). Serotype 0:3 and 0:9 are both faecal commensals and inhibitants of the oral cavity of pigs, especially the tonsils and tongues. Serotype 0:3 is also frequently encountered as a surface contaminant on freshly slaughtered pig carcasses (Andersen, 1988; Nesbakken, 1988; Nesbakken, et.al., 1985).

Many surveys, reviewed elsewhere (Table 1.6) have demonstrated the common occurrence of Y. enterocolitica and related microbes in the intestinal tract and oral cavity of healthy slaughter pigs. The earlier reports of Y. enterocolitica in pigs were based on examination of faeces or intestinal contents. It was later demonstrated that the isolation frequency of these bacteria was approximately ten times greater from the tongues or tonsils than that obtained from faeces (Pedersen, 1979; Schiemann, 1980; Wauters, 1979). The reported isolation rates range up to and in excess of 56.0% (Table 1.6) depending on the type of samples examined (tongues, tonsils, throat swabs), geographical origin, and efficacy of the isolation methods.

In Belgium, which is the country with the highest reported incidence of Y. enterocolitica infection of people, a case control study has shown that the infection was strongly associated with eating raw pork (Tauxe et.al., 1987). The apparent rareness of Y. enterocolitica infection in Moslem countries (Samadi et.al., 1982) also supports the potential role of pork as the vehicle of Y. enterocolitica infection.

In New Zealand, reports of human disease due to yersiniosis, and of isolations from healthy subjects have been sporadic (Henshall, 1963; Lello and Lennon, 1992; Malpass, 1981; McCarthy and Fenwick, 1990; Rose, 1976), and an active search of possible hosts of species of *Yersinia* has only begun in earnest in recent years.

Prior to this study, only one published report of isolation of yersiniae from pigs in this country existed (Hodges *et.al.*,1984). However, unpublished data by Fenwick (*pers.comm* 1989) suggested that pigs may be carriers of human pathogenic yersiniae in their tonsils.

This study was therefore conducted as a follow up to the former, with the aim of confirming the findings and establishing some epidemiological aspects of the occurrence of yersiniae in the tonsils of slaughtered pigs.

The study was conducted in two phases. The first phase involved a cross-sectional study to determine the presence of species of *Yersinia* in the tonsils of slaughtered pigs and their distribution among farms supplying pigs for slaughter. The second phase, a longitudinal study, which was based on findings from the first, involved a selection of farms in relation to their particular carriage of species of *Yersinia*. Abattoir sampling was carried out on a monthly basis for twelve months, with the objective of investigating the seasonal effects on the occurrence of species of *Yersinia* in the tonsils of slaughtered pigs. During this phase, isolates were tested for possible virulence-associated characteristics with the aim of determining the role of pigs as possible reservoirs for human infection with yersiniae and thus the potential public health significance of pathogenic strains which may be harboured as free-living commensals in the tonsils of slaughtered pigs.