

Variation at the common polysaccharide antigen locus drives lipopolysaccharide diversity within the *P. syringae* species complex

Jay Jayaraman^{1,2}, William T. Jones³, Dawn Harvey³, Lauren M. Hemara^{1,2,4}, Honour C. McCann^{5,†}, Minsoo Yoon¹, Suzanne L. Warring⁶, Peter C. Fineran^{2,6}, Carl H. Mesarich^{2,7}, Matthew D. Templeton^{1,2,4*}

¹Bioprotection Technologies, The New Zealand Institute for Plant and Food Research Limited, Auckland, New Zealand, ²Bioprotection Centre for Research Excellence, New Zealand, ³Bioprotection Technologies, The New Zealand Institute for Plant and Food Research Limited, Palmerston North, New Zealand, ⁴School of Biological Sciences, University of Auckland, New Zealand, ⁵Institute of Advanced Studies, Massey University, Auckland, New Zealand, ⁶Department of Microbiology and Immunology, University of Otago, Dunedin, New Zealand, ⁷School of Agriculture and Environment, Massey University, Palmerston North, New Zealand.

*correspondence to Matthew Templeton (matt.templeton@plantandfood.co.nz)

[†]present address Max Planck Institute for Developmental Biology, Max-Planck-Ring 9, D-72076 Tübingen, Germany.

ABSTRACT

The common polysaccharide antigen (CPA) from the lipopolysaccharide (LPS) component of cell walls from the species complex *Pseudomonas syringae* is highly variable both in structure and immunological specificity, but the genetic basis for this is not well understood. We have characterised the CPA locus from *P. syringae* pv. *actinidiae* (*Psa*). This locus has a modular structure with genes for both L- and D-rhamnose (Rha) biosynthesis and that of an unknown sugar. It also contains an operon coding for ABC transporter subunits, a bifunctional glycosyltransferase and an O-methyltransferase. This operon is predicted to have a role in transport, elongation and termination of the Rha backbone of the CPA oligosaccharide and is referred to as the TET operon. This is the first report of the identification of this operon in *P. syringae*. Two alleles of the TET operon were present amongst the different biovars of *Psa* and lineages of the closely related pathovar *P. syringae* pv. *actinidifoliorum*. This allelic variation was reflected in the electrophoretic properties of purified LPS from the different isolates. Gene knockout of the TET operon allele from biovar 1 and replacement with that from biovar 3, demonstrated the link between the genetic locus and the electrophoretic and immunogenic properties of the LPS molecules in *Psa*. Sequence analysis of the TET operon from a wide range of *P. syringae* and *P. viridiflava* isolates displayed a phylogenetic history which is incongruent with core gene phylogeny, but correlates with previously reported tailocin sensitivity, suggesting a functional relationship between LPS structure and tailocin susceptibility.

INTRODUCTION

Pseudomonas syringae pv. *actinidiae* (*Psa*) is the causal agent of canker disease in kiwifruit (*Actinidia* Lindl spp.). Outbreaks of the disease were first observed in Japan and Korea in the 1980s and 1990s respectively, but isolates of the bacterium responsible for these outbreaks did not spread from their country of origin (Koh *et al.*, 1994; Serizawa *et al.*, 1989; Takikawa *et al.*, 1989). Between 2008 and 2010, a pandemic clone spread around the world, devastating the majority of regions growing cultivars of *A. chinensis* var. *chinensis* (gold kiwifruit) (Scortichini *et al.*, 2012). A large number of *Psa* isolates from a range of geographical origins have been sequenced, and phylogenies generated from the core genomes show that the three emergences of the disease are closely related, but form distinct clades (McCann *et al.*, 2017; McCann *et al.*, 2013). Single Nucleotide Polymorphism (SNP) analysis indicates that the recent pandemic isolate originated in China, but the location of the source population of this pathovar has not been identified (McCann *et al.*, 2017). Despite the close phylogenetic relationship between *Psa* isolates, their accessory genomes vary significantly in their effector complement and secondary metabolite portfolios (Butler *et al.*, 2013; Marcelletti *et al.*, 2011; Mazzaglia *et al.*, 2012; McCann *et al.*, 2013). Based on both the phylogeny of the core genome and accessory gene variation, *Psa* isolates have been designated as biovars (BVs) (Cunty *et al.*, 2015; Vanneste *et al.*, 2013). The isolates that only cause leaf spots on kiwifruit, initially described as BV4, are closely related to *Psa*, but have now been given their own pathovar designation; *P. syringae* pv. *actinidifoliorum* (*Pfm*), and consist of four distinct lineages (Cunty *et al.*, 2015). Recently, two more *Psa* BVs have been discovered in Japan (Fujikawa and Sawada, 2016; Sawada *et al.*, 2016). Variation in the accessory genome of *P. syringae* pathovars and *Psa* BVs, and its

role in host specificity has been well documented (Dillon *et al.*, 2019a; McCann *et al.*, 2013). In addition, *P. syringae* isolates have been shown to have a high degree of structural and serological variation in their lipopolysaccharide (LPS) (Zdorovenko and Zdorovenko, 2010). However, the genetic basis for this is not well understood. Both bacteriophages and tailocins utilise LPS as a receptor to recognise and bind to their host. Tailocins are derivatives of bacteriophages, comprising predominantly the tail proteins that function to depolarise the bacterial membrane (Riley and Wertz, 2002). Isolates of *P. syringae* use tailocins (also known as R-type syringacins) to target and outcompete closely related strains that presumably occupy a similar ecological niche (Hockett *et al.*, 2015). It has been shown recently that there is a high degree of variation in the sensitivity to tailocins within *P. syringae* (Baltrus *et al.*, 2019). Understanding the molecular basis of this variation is important if we are to implement novel methods such as the use of phage therapy or tailocins for biological control of this pathogen (Baltrus *et al.*, 2019; Frampton *et al.*, 2015; Frampton *et al.*, 2014; Pinheiro *et al.*, 2019; Rooney *et al.*, 2019).

LPS are complex glycolipids that make up the surface leaflet of the outer membrane of gram-negative bacteria, forming a physical protective barrier. LPS have three distinct domains: Lipid A that tethers the molecule to the outer membrane, the core domain, and finally the O-polysaccharide (OPS). The last domain includes the O-specific antigen (OSA) and/or the common O-polysaccharide antigen (CPA), and is structurally the most variable component of the LPS. The distinction between the two OPS polysaccharides is that the CPA is exported via an ABC transporter-dependent pathway, while OSA synthesis follows the Wzx/Wzy-dependent pathway (Raetz and Whitfield, 2002). In many bacterial species, the OPS are responsible for the immunological variation between isolates and, for human and

animal pathogens, subspecies classification is based on serotypes (DebRoy *et al.*, 2016; King *et al.*, 2009). Similarly, the structures of OPS in plant-pathogenic bacteria such as *P. syringae* are highly variable at the pathovar level (Ovod *et al.*, 1997a). For *P. syringae*, the OPS appears to be comprised of the CPA (Kutschera *et al.*, 2019; Mesarich *et al.*, 2017) with a backbone consisting of a tri- or tetra-saccharide repeating unit, comprising either L- or D-Rhamnose (Rha), or a combination of the two, with various linkages. These are decorated with side chains consisting of different sugars such as *N*-acetyl-glucosamine (GlcNAc), fucose (Fuc), Rha, and *N*-acetyl-fucosamine (FucNAc) (Ovod *et al.*, 1997a; Zdrovenko and Zdrovenko, 2010). The OPS can also be methylated or acetylated to varying degrees (Zdrovenko *et al.*, 2001). At least nine serotypes of *P. syringae* have been identified, and each differs in the structure of the OPS backbone and side-chain modifications (Zdrovenko and Zdrovenko, 2010). Although considerable effort in the past was devoted to attempts at relating serotype to pathovar identification and taxonomy, correlations were not consistently observed (Ovod *et al.*, 1997b). Little is known about the biosynthesis of LPS in *P. syringae* or the genetic basis for the observed structural and immunological variability within the genus.

Transposon mutagenesis of *Psa* BV3 using Tn5 identified a set of mutations with a rough (R) colony phenotype (Mesarich *et al.*, 2017). This phenotype is often associated with mutations in LPS biosynthesis. Indeed, the majority of the Tn5 inserts in these R-LPS mutants mapped to a gene cluster in *Psa* orthologous to the CPA biosynthetic pathway from *P. aeruginosa* (Mesarich *et al.*, 2017). Furthermore, LPS was not observed in the mutants, suggesting CPA is the sole OPS in *Psa* (Hockett *et al.*, 2017). Other recent papers have directly identified components of

121 CPA biosynthesis in *P. syringae* through gene knockouts (Kutschera *et al.*, 2019)
 122 and increased tailocin resistance (Kandel *et al.*, 2019; Kutschera *et al.*, 2019).

123 Here, we investigated the genetics of LPS biosynthesis in *Psa*. We
 124 characterised the CPA locus and the orthologous region in the different *Psa* BVs, *P.*
 125 *syringae* and the closely related *P. viridiflava* isolates. We show an operon from this
 126 locus is hyper-variable and accounts for the high degree of structural and
 127 immunological variation observed in this species complex. Furthermore, the
 128 relationship between this component of the CPA locus and sensitivity to tailocins was
 129 highly correlated.

RESULTS

Major LPS biosynthesis genes in *Psa* BV3 are shared with *P. aeruginosa*

In *P. aeruginosa* PAO1, the genes and operons responsible for the biosynthesis of Lipid A, the core LPS and the OPS tend to be co-located and have been well annotated and characterised (King *et al.*, 2009; Lam *et al.*, 2011). We identified the key loci containing the majority of the LPS biosynthetic genes in *Psa* BV3 using BLASTx with *P. aeruginosa* PAO1 orthologs (Figure 1A & Table 1). Amino acid similarity to *P. aeruginosa* PAO1 orthologs of the Lipid A, core LPS, and Rha biosynthetic pathways in the CPA locus is relatively high (Table 1). In contrast, *Psa* genes putatively coding for other components of the CPA, such as the *wzm/wzt/wbpX* orthologues, are more divergent and have poor or no significant BLASTx hits. Evidence for an OSA biosynthetic cluster was not found, apart from genes in the *wbpK-M* locus, which is also involved in CPA biosynthesis (Table 1). The co-location of genes annotated as ABC transporters and glycosyltransferases (GT) with both the D- and L-Rha biosynthetic pathways suggests that the CPA is the predominant OPS in *Psa* BV3 (Figure 1A).

Bioinformatic identification of the CPA locus in *Psa* BV3

Identifying the CPA locus in *P. syringae* has been challenging due to the lack of similarity to *wzm/wzt/wbpX* orthologues (Kutschera *et al.*, 2019). Previous work identified five R-LPS mutants by Tn5 mutagenesis that had inserts in a region with homology to the CPA biosynthetic pathway from *P. aeruginosa* PAO1 (Mesarich *et al.*, 2017). Bioinformatic analysis of this region revealed a locus of 18 genes (Figure 1A). These genes were arranged in operons predicted to be involved in the synthesis

of L-Rha, D-Rha, an unknown sugar, an ABC transporter complex involved in the transport of the CPA, a bifunctional glycosyltransferase, an O-methyltransferase, and a region of unpredicted function (Table 2). Of the two genes annotated as ABC transporters, the first (IYO_023015) has homology to the ABC permease superfamily gene *wzm*, while the second (IYO_023010) has an N-terminal ATPase domain with a C-terminal *wzt*-like domain (hereafter called *wzt*). The latter domain binds to the non-reducing terminal modification of the sugar chain and is thus specific for the transported molecule (Cuthbertson *et al.*, 2007). The GT gene (IYO_023000) on the same operon as the two ABC transporters has two GT4 domains, implying it catalyses two distinct glycosyl transfer reactions. It is homologous to *wbdA* (formerly known as *mtfB*), a functionally characterised mannosyltransferase in *E. coli*, which has been shown to direct the extension of the O9-specific polysaccharide chain by synthesising, then polymerising, the tetra-saccharide repeat (Liston *et al.*, 2015). The adjacent gene (IYO_023005) on the *Psa* BV3 operon is annotated as an O-methyltransferase. In *E. coli*, the LPS oligosaccharide chain extension, catalysed by WbdA, is terminated by a dual methyltransferase/kinase enzyme (WbdD) (Mann *et al.*, 2019). It appears therefore that this operon possesses genes required for the transport, extension, and termination of the backbone oligosaccharide chain of the CPA in *Psa* BV3 (hereafter referred to as the TET operon). This is the first time this operon has been identified in *P. syringae*.

The TET operon from *Psa* and *Pfm* is encoded by two different alleles

The CPA locus from representatives of all five *Psa* BVs and two lineages of *Pfm* were identified and compared. The TET operon, genes for D-Rha biosynthesis, and

regions coding for a chemotactic receptor, a DnaJ molecular chaperone and a tRNA-Met were shared and could be aligned (Figure 1B). The latter three genes were syntenic in all genomes but are not part of the CPA locus. The region including the D-Rha biosynthetic pathway and the first gene (*wzm*) in the TET operon (5' region) is well conserved with few SNPs and a phylogenetic tree (Figure 1C) closely matched the topology of a tree generated from seven conserved genes (Sawada *et al.*, 2016). In contrast, the 3' region, from the gene encoding the ABC transporter ATPase (*wzt*) of the TET operon is more variable, representing two groups of divergent sequences. One group includes *Psa* BVs 1 and 2, and *Pfm* L3, while the other includes *Psa* BVs 3, 5, and 6, and *Pfm* L1. A phylogenetic tree of this region (Figure 1D) resolved into two clades that did not match the topology of Figure 1C, or the phylogeny generated from the core genome (Supplementary Figure S1). These results suggest that variation at the TET operon is driving diversity at the CPA locus in *Psa* and *Pfm*.

LPS from *Psa* and *Pfm* has different biochemical properties

To assess whether genetic variation in the TET operon results in variation of the biochemical properties of LPS, we extracted LPS from *Psa* and *Pfm* using the method of Westphal and Jann (1965). Resolution of the extracted LPS by SDS-PAGE revealed the classic ladder pattern characteristic of bacterial LPS molecules, a reflection of the length and charge micro-heterogeneity of the O-linked saccharides (Jann *et al.*, 1975). Comparison of the LPS between the five different BVs of *Psa* and two lineages of *Pfm* revealed two patterns based on differences in the molecular weight range and spacing between bands (Figure 2A). LPS from *Psa* BV1, BV2 and

Pfm L3 formed one group, which was characterised by a broad range of molecular weights, especially in the higher molecular mass range. In contrast, *Psa* BV3, BV5, BV6 and *Pfm* L1 formed a second group that had a lower range of molecular weights and narrower spacing between bands (Figure 2A). The two electrophoretic pattern groups reflect the broad clusters evident in the TET operon tree (Figure 1D), suggesting the TET operon plays a significant role in defining the electrophoretic properties of LPS. Banding patterns within a group were not identical, suggesting that they may differ in decoration of the back-bone with side-chain sugars and/or acylation. Therefore, the genetic and biochemical data indicate that the CPA locus identified in *Psa* and *Pfm* is likely responsible for LPS biosynthesis in these *P. syringae* pathovars.

LPS from *Psa* and *Pfm* are immunologically distinct

Western blots of the various LPS preparations for *Psa* and *Pfm* isolates were probed with each isolate-specific polyclonal antibody preparation (Figure 2B). All purified polyclonal antibodies showed a high degree of specificity, with little cross-reactivity against closely related isolates. The anti-*Psa* BV3 antibody preparation was the most specific, reacting with BV3, BV5, and only weakly to BV6 LPS (Figure 2Bi). The weak binding to the *Psa* BV6 LPS and the slightly different *Psa* BV6 LPS profile (Figure 2A) suggest a subtle difference in the CPA decoration. There was no cross-reactivity of the *Psa* BV3 antibody to the *Psa* BV1, BV2, or *Pfm* LPS preparations (Figure 2B). The lack of binding to *Pfm* L1 LPS (which appears more similar to BV3 LPS than *Pfm* L3) is likely due to the fact that antibodies that bind to both BV3 and *Pfm*L1 were depleted by the affinity chromatography purification process used. The

anti-BV1 antibody preparation reacted strongly with both *Psa* BV1 and BV2 LPS, as expected based on the close genetic identity between their TET loci (Figure 2Bii). There was also weak cross-reactivity of *Psa* BV1 with *Psa* BV3 and BV5 LPS preparations (Figure 2Bii). Interestingly, the *Pfm* L1 antibody had a slight cross-reactivity with *Pfm* L3, but not with the *Psa* isolates in the same grouping (Figure 2Biii). These results further support the different grouping of LPS between these strains, but also indicate that antibody specificity is influenced by both backbone structure and side-chain decorations.

The TET operon determines the variability of LPS banding in *Psa*

To confirm whether the TET operon, expressing the ABC transporter (*wzt*), putative oligosaccharide chain elongation GT (*wbdA*), and the O-methyl transferase (*wbdD*), are responsible for the different LPS profiles, this operon was deleted in *Psa* BV1 (Figure 3A; dark green region, Supplementary Figure S2) and replaced with that from *Psa* BV3 (Figure 3A, blue region, Supplementary Figure S2). The *Psa* BV1 TET operon knock-out (Δ LPS KO) lacked LPS and possessed a rapid sedimentation phenotype, but had a very weak R-LPS colony morphology, as observed for the BV3 mutants (Figure 3B; Supplementary Figures. S3A & S4). *Psa* BV1 with the BV3 LPS knocked back in (four knock-in [KI] lines), but not for the KO revertants that failed to retain this region, restored these LPS-associated phenotypes (Figure 3B, Supplementary Figures. S3B & S4). Furthermore, the LPS profile was identical to that of wild-type BV3, proving this locus codes for the genes involved in generating the BV-specific LPS oligosaccharide ladder. A Western blot of the LPS gel of these strains was probed with the *Psa* BV3 antibody (Figure 3C), which showed that

transfer of the TET locus from *Psa* BV3 to the *Psa* BV1 Δ LPS KO also transferred the antibody specificity. In summary, this genetic TET-swap analysis provided direct evidence that these genes were necessary and sufficient for the differences in LPS between these strains.

LPS is required for full pathogenicity of *Psa* in kiwifruit plantlets

Previously, LPS mutations have been found to affect host colonization by *P. syringae* (Kutschera *et al.*, 2019). To investigate the role of LPS in the virulence or pathogenicity of *Psa*, we infected kiwifruit plantlets with *Psa* BV3 R-LPS Tn5 mutants (Δ wbpL, Δ wzm, Δ wbdD, Δ wzt, and Δ gmd) (Mesarich *et al.*, 2017) and the *Psa* BV1 knock-out (Δ LPS), two knock-in (Δ LPS+LPS_BV3) and two knock-out revertant (Δ LPS-LPSrev) mutants, along with their wild-type counterparts. All LPS mutants showed at least a log reduction in growth *in planta* compared to wild-type strains, indicating that LPS is required for host colonization (Figure 4A-B) and symptom development (Supplementary Figure S5A-B). Meanwhile, the BV1 Δ LPS+LPS_BV3 strains with complemented LPS production restored wild-type growth to the Δ LPS mutant (Figure 4B; Supplementary Figure S5B) and this did not occur in the strains that had reverted to Δ LPS. This correlated with the strains' ability to stay suspended in inoculation buffer (Supplementary Figure S3A-B).

The TET operon from *P. syringae* isolates is highly variable but correlates with LPS

Having identified the TET operon in *P. syringae* for the first time we set out to analyse this operon in the *P. syringae* species complex and closely related plant

associated *Pseudomonas* species. In *P. syringae*, the composition of the CPA backbone (D- and/or L-Rha), the number of sugars in the repeat, and the bonds linking the sugars within the oligosaccharide, vary between different *P. syringae* isolates (Ovod *et al.*, 1997a; Zdorovenko and Zdorovenko, 2010). BLASTx and Conserved Domain Database (CDD) searches identified the CPA locus in other *P. syringae* pathovars and epiphytic isolates. We also included several *P. viridiflava* isolates (See Supplementary Table 1 for details of all isolates analysed). Comparison of these loci revealed variation in the sugar biosynthetic pathways comprising the CPA locus, including the TET operon. In some *P. syringae* isolates, the GT and O-methyltransferase genes were fused. The TET operon is the only one consistently present at the CPA locus, and it showed a high degree of sequence variation between *P. syringae* pathovars and isolates. Variation in CPA sugar biosynthetic pathways and the sequence divergence of the TET operon made the identification of this region in *P. syringae* challenging, but as noted previously, the CPA locus was consistently located adjacent to genes encoding a chemotactic receptor, DnaJ molecular chaperone and a tRNA-Met (Figure 1A). A gene encoding an integrase was often located adjacent to the tRNA-Met in an orientation reminiscent of an integron (Domingues *et al.*, 2012; Gillings, 2014). Given that tRNAs are often sites for integration of mobile elements, this might explain the modular and variable nature of the CPA locus in *P. syringae*. However, a canonical integron structure was not found using programs such as integron finder (Cury *et al.*, 2016).

Phylogenetic analysis of the concatenated proteins from the TET operon generated a tree with five main clades (including the two clades identified in *Psa* and *Pfm*), each representing different sets of alleles of this operon from the *P. syringae*

and *P. viridiflava* species complexes (Figure 5A). As observed with *Psa* and *Pfm* isolates, the topology of the tree made from these sequences is very different from the corresponding tree generated from five core genes from the respective genomes (Supplementary Figure S1). This indicates that the TET operon has an evolutionary history driven by recombination or horizontal gene transfer that is distinct from the evolution of the core genome.

Since genetic variation in the CPA between even closely related isolates of *Psa* and *Pfm* is reflected in different LPS profiles, we performed a broader analysis of LPS profiles across different clades of *P. syringae* (Figure 5A). Eleven isolates of *P. syringae* were chosen, representing four TET operon phylogenetic clades. The banding patterns were different for all isolates, but similarity within clades was observed. Notably, the spacing between bands was clearly similar within clades, but varied between them (Figure 5B). Interestingly, the range of molecular weights of different LPS molecules within clades was highly variable, suggesting the efficiency of the chain terminating enzyme (WbdD) varies within a clade. While considerable variability has been demonstrated in *P. syringae* CPA, the functional relevance of this is less well understood.

Correlation between tailocin sensitivity and TET operon variability

Like phages, many R-type syringacins (tailocins) use LPS as a receptor. Recently the R-syringacin killing and sensitivity spectra for a diverse range of *P. syringae* isolates were characterised (Baltrus *et al.*, 2019). A feature of both spectra is that dendrograms generated from the killing and sensitivity matrices do not match MLST phylogenetic trees (Baltrus *et al.*, 2019). Since LPS is a likely receptor for R-

syringacins, we hypothesised that there might be a correlation between the patterns of variation in tailocin sensitivity within *P. syringae* isolates and the genetic variation in the TET operon responsible for the structure of the CPA backbone.

To test this hypothesis, we generated a UPGMA phylogenetic tree from the TET operons from 29 of the isolates tested for tailocin killing and sensitivity by Baltrus *et al.* (2019). This was compared to a hierarchical clustering tree generated from the sensitivity profile of the same isolates using a tanglegram (Figure 6A). Tanglegrams compare phylogenetic trees to determine how similar they are and correlation coefficients can be calculated from the similarity between different pairs of trees (Galili, 2015). The correlation coefficient between the sensitivity and TET operon trees was 0.987, indicating a high degree of congruence (Supplementary Figure S6). Both trees comprise two main clades, and there was a 100% correlation among the isolates between the clades in both trees. In contrast, tanglegrams between either the sensitivity profile or the TET operon and an MLST phylogenetic tree had low correlation coefficients of 0.011 and -0.003, respectively, indicating poor congruence (Figure 6B-C; Supplementary Figure S5). These results suggest a strong relationship between tailocin sensitivity and the structure of the CPA oligosaccharide of LPS in *P. syringae*. They provide further evidence that LPS is a tailocin receptor in *P. syringae*, explaining the complex host range observed for these proteins (Baltrus *et al.*, 2019) and providing a plausible explanation for the diversification pattern of these loci.

DISCUSSION

LPS is a highly complex macromolecule located on the surface of the bacterial cell wall that interfaces with the environment and other organisms. It is often used as a receptor or ligand for recognition by other bacteria, phages, bacteriocins and the defence system of host organisms. The modular structure can be used to generate structural variability, particularly in the OPS, which can help bacteria avoid parasitic organisms, toxins and host defences (King *et al.*, 2009). The structure and function of LPS has been well explored in human and animal bacterial pathogens where it has been shown that LPS interacts directly with the host immune system and hence immunity is a strong driver for genetic variation at loci coding for LPS (Lerouge and Vanderleyden, 2002; Maldonado *et al.*, 2016).

In contrast, the role of LPS in plant-pathogenic and epiphytic bacteria is less clear. In this study, we have identified the CPA locus in *Psa* and shown it is responsible for the biosynthesis of the predominant OPS in *P. syringae*. The locus is highly modular with operons predicted to be involved in the biosynthesis of various sugars, although only those for L-Rha and D-Rha have been convincingly identified through similarity. The TET operon that codes for proteins involved in the polymerisation and transport of the backbone oligosaccharide through the inner membrane was also identified for the first time in *P. syringae*. This was made possible by the use of LPS screens of our transposon library (Mesarich *et al.* 2017). This operon was present in the majority of *P. syringae* and *P. viridiflava* CPA loci examined. Phylogenetic analysis of three of the proteins in the TET operon revealed five distinct clades, which most likely reflect the different unit structures of the CPA Rha-backbone oligosaccharide. There are two elements that contribute to the structural variability of the CPA in *P. syringae*. One is the nature of the backbone

oligosaccharide, which can be tri- or tetra-saccharides of either D-Rha, L-Rha or both, linked by α 1-2, α 1-3, or β 1-4 bonds (Ovod *et al.*, 1997a). The other is the side-chain decoration, which is usually a single sugar residue attached by a variety of different linkages. In addition, acetylation is a common back-bone modification. Unfortunately it is not possible to directly correlate the structural information accurately with the clades resolved in Figure 5 because the isolates for which structural LPS information is known are not those that have been sequenced (Zdorovenko and Zdorovenko, 2010).

The mechanism by which the different operons within the CPA locus are shuffled between *P. syringae* isolates to generate the observed structural variability could occur via homologous recombination or insertion via an integron-like cassette. The structure of the CPA locus in *Psa* BV3 resembles that of an integron, although an intact phage-like integrase adjacent to the tRNA-Met was not present in all other *P. syringae* isolates. In addition, a classic integron promoter and *attC* insertion sequences were not found using programs such as integron finder. The presence of operons from *P. viridiflava* in the phylogenetic trees suggest that the mechanism of recombination between isolates is not exclusive to the *P. syringae* species complex, but also includes other closely related species such as *P. viridiflava* that share the same ecological (plant-associated) niche.

The variation at the TET locus is also reminiscent of that at the capsule locus within the *Klebsiella pneumoniae* clonal group 258 (Wyres *et al.*, 2015). In this case it has been shown that extensive intra-clonal variation at the capsule polysaccharide synthesis locus is mediated by large-scale recombination events. This has been responsible for generation of the 78 immunologically distinct capsule variants which have been described in *K. pneumoniae* (Wyres *et al.*, 2015).

Tailocins are a subset of bacteriocins derived from phage tail proteins that have been co-opted by host bacteria as a means for killing closely related bacterial isolates occupying the same ecological niche (Ghequire and De Mot, 2015). What has been demonstrated recently within *P. syringae* is the depth and complexity of both the killing and sensitivity spectra generated from a wide range of isolates (Baltrus *et al.*, 2019). While it is clear that LPS is a receptor for some tailocins, a link between the sensitivity to tailocins and the structural variation in LPS has not been demonstrated until now. Our results indicate a strong relationship, suggesting that killing by tailocins (and by inference infection by phage) may drive diversity in LPS structure. It is interesting to note that the genetic variation driving these differences appears in both cases to be mediated by localised recombination events (Baltrus *et al.*, 2019).

As has been reported previously for other *P. syringae* LPS mutants (Kutschera *et al.*, 2019), those in both the *Psa* BV3 and BV1 backgrounds displayed significantly reduced *in planta* growth compared to their wild-type strains at both 6- and 12-days post-inoculation (Figure 4). The inability of the LPS knockout isolates to remain suspended in solution may also affect the ability of these mutants to move *in planta* and could be responsible for reduced pathogenicity. Although mutants lacking LPS have been shown to be highly resistant to particular bacteriophages and/or tailocins, the loss of pathogenicity probably ensures that strain completely lacking LPS are strongly selected against *in planta* (Kandel *et al.*, 2019). This means that gene replacements at this locus should be favoured over simple gene deletion events.

There did not appear to be any obvious geographical bias for any of the CPA clades. While this is unlikely to be an exhaustive identification of TET operon

variants, a significant proportion of available *P. syringae* whole genome sequences were analysed in this study. Since members of four clades were isolated recently from kiwifruit in New Zealand, the TET operon alleles identified are likely ubiquitous among *P. syringae* populations. These findings, firstly that LPS is a trait conferred by a predictable locus, and secondly that LPS is required for pathogenicity, offer intriguing possibilities for a durable disease resistance program using bacteriophage/tailocin-mediated control. While the locus is open to recombination, the discovery of the population of TET operon alleles present on a plant species may facilitate a comprehensive mixture of bacteriophage/tailocin that covers the most common alleles to be deployed to increase efficacy and durability of this treatment by circumventing escape through recombination at the TET operon locus.

EXPERIMENTAL PROCEDURES

Bacterial isolates used in this study

Metadata for the bacterial isolates and genomes used in this study are listed in Supplementary Table 1. Lysogeny broth (LB), and M9 minimal media (M9) (Sambrook and Russell, 2001) were supplemented with agar 1.5% (w/v) and 50 µg/mL kanamycin (Km) where required. *Escherichia coli* and *Pseudomonas* isolates were routinely cultured at 37°C and 28°C, respectively. All kits and reagents were used, except where specified, in accordance with the manufacturer's instructions.

Bioinformatics

Bioinformatics analysis was carried out using the Geneious R10 platform (www.geneious.com). For unannotated genomes, the RAST pipeline was used for gene calling (Aziz *et al.*, 2008). GenBank searches were carried out using BLAST (Boratyn *et al.*, 2013). Sequences were searched for *attI* and *attC* motifs using Integron Finder https://github.com/gem-pasteur/Integron_Finder.

LPS purification

For small-scale preparations of LPS, the hot-phenol method was used (Westphal and Jann, 1965). Overnight bacterial cultures were resuspended in PBS (500 µL) to an A₆₀₀ of between 2 and 5. An equal volume of Tris-saturated phenol (Sigma-Aldrich, Mo, USA, P4557) was added and the sample was incubated at 68°C for 90 min with intermittent vortexing. The supernatant was sequentially treated with DNase I (37°C) and RNase (65°C), each for 1 h. The samples were treated with

phenol/chloroform/isoamyl alcohol, followed by chloroform/isoamyl alcohol and stored at -20°C. To generate antigen used for immunization and polyclonal antibody generation, large-scale extraction of LPS from *Psa* BV1 and BV3, and *Pfm* L1 was performed using the hot-phenol method of Maier and Brill (1978). Cells were harvested in late exponential phase, collected by centrifugation and freeze-dried for storage at -20°C.

LPS were resolved using 4–12% premade NuPAGE™ Bis-Tris SDS gels on a Novex system with MOPS as the running buffer. Gels were stained using the Pro-Q™ Emerald 300 Lipopolysaccharide Gel Stain Kit (Molecular Probes, OR, USA). For Western blotting, gels were transferred to a polyvinylidene difluoride (PDVF) membrane (Millipore, Burlington, MA). The anti-*Psa* antibodies (see below) were used at 1:5,000 dilution and the Anti-Rabbit IgG-Peroxidase antibody (Sigma-Aldrich, St Louis, MO) was used at 1:10,000 dilution. Chemiluminescence was developed using the Clarity Max Western ECL (Bio-Rad, Hercules, CA).

Production of antibodies to *Psa* biovars

Heat-killed cells from *Psa* BV1 (ICMP 9617), *Psa* BV3 (ICMP 18884), and *Pfm* L1 (ICMP 18803) in Freund's complete adjuvant were each used to individually immunise two rabbits by subcutaneous injection of 10⁸ cells. Rabbits were rested for 4 weeks and immunised every 2 weeks by three further injections. Blood was collected 1 week after the last injection, and rabbit IgG was purified from serum using ammonium sulphate fractionation and protein-A sepharose chromatography (Ey *et al.*, 1978). These polyclonal antibodies were further purified by affinity chromatography through matrices attached to LPS extracted from the above isolates

not used for immunisation in each of the three cases. The aim of this step was to remove antibodies that bound to antigens in common with other isolates, to give an isolate-specific polyclonal antibody preparation. For example to purify antibodies specific for *Psa* BV3, IgG were passed through affinity columns with LPS from *Psa* BV1 and *Pfm* L1. For preparation of the matrices LPS was dissolved in buffer and oxidised using periodate. Excess periodate was removed by gel filtration using a PD10 column. Oxidised LPS was reacted with biotin hydrazide to biotinylate the LPS as per method sheet 28020 (Thermo Fischer Scientific). LPS matrices for affinity chromatography were prepared by attachment of biotin-LPS to streptavidin-labelled sepharose columns. Biovar-specific LPS antibodies were purified by standard affinity chromatography as described by the manufacturer (GE Life Sciences, PA USA).

LPS locus knock-out and knock-in

To generate the LPS TET operon (*wzt* – *wbdA*; ~5.3 kb) deletion in *Psa* BV1 (ICMP 9617), a modified pK18mobsacB vector-based method was utilized (Kvitko and Collmer, 2011). DNA fragments containing the upstream (~1 kb) and downstream (~1 kb) regions of the LPS knock-out region were amplified using PCR with primer pairs *PsaJ_LPS-KO_UP-F/PsaJ_LPS-KO_UP-R* and *PsaJ_LPS-KO_DN-F/PsaJ_LPS-KO_DN-R*, respectively, with the UP-R and DN-F primers carrying added *Xba*I restriction enzyme sites (Supplementary Table 2). Each PCR fragment was gel-purified using an EZNA gel extraction kit (Custom Science, Auckland, NZ) and digested with *Xba*I, re-purified with an EZNA PCR product purification kit and ligated (to the *Xba*I-cut overhangs) to form a 2 kb KO fragment. The 2 kb fragment containing both the upstream and downstream fragments for the KO region was then

re-amplified by PCR using primers PsaJ_LPS-KO_UP-F and PsaJ_LPS-KO_DN-R. This PCR product was cloned into the *Eco53kl* blunt-end restriction enzyme site of pK18mobsacB that had first been mutagenized to remove the non-MCS *Eco53kl* site to generate pK18BΔE (Schafer *et al.*, 1994). The pK18BΔE vector carrying the ~2 kb KO fragment, called pΔLPS, was transformed into *E. coli* DH5α, plated on X-gal/IPTG/kanamycin LB agar for blue/white selection. Positive transformants were confirmed by Sanger sequencing (Macrogen, Seoul, South Korea). *Psa* BV1 was electroporated with the pΔLPS construct and transformants were selected on LB agar with nitrofurantoin, cephalixin and kanamycin. Selected colonies were subsequently streaked onto LB agar containing 10% (w/v) sucrose to counter-select plasmid integration, forcing removal of the *sacB* gene, resulting in either revertants to wild-type or knock-outs. KOs were confirmed using colony PCR with primers PsaJ_LPS-KO_Check-F and PsaJ_LPS-KO_Check-R, with PCR products sent for Sanger sequencing with the nested cloning PsaJ_LPS -KO_UP-F and PsaJ_LPS -KO_DN-R primers, as well as using internal gene-specific primers. Selected colonies were also plated on kanamycin-containing medium to confirm loss of the plasmid backbone containing kanamycin resistance and the *sacB* gene.

To clone the knock-in (KI) construct of the *Psa* BV3 LPS TET operon for complementation of the *Psa* BV1 LPS TET operon knockout (*Psa* BV1 ΔLPS), the golden gate cloning system was used (Engler *et al.*, 2009). Firstly, the destination vector was generated by restriction digestion using *Asel* and *NheI* of pK18mobsacB to remove the multiple cloning site (MCS). This was replaced with a golden gate-compatible (*BsaI* restriction enzyme site-flanked) MCS from pICH86988 using In-Fusion cloning according to kit instructions (Takara Bio, USA) to generate a golden gate cloning-compatible version of the pK18mobsacB vector; pK18B-GG. The

528 pK18B-GG vector was transformed into *E. coli* DH5 α , plated on kanamycin LB agar
529 and screened by *Bsa*I restriction digest. Positive transformants were confirmed by
530 Sanger sequencing (Macrogen, South Korea). Entry-level vectors for the LPS-KI
531 construct were generated for the LPS region from *Psa* BV3 (ICMP 18884), divided
532 into four modules (modules 1–4), with overlap-fusions of the upstream (5′) and
533 downstream (3′) regions of the *Psa* BV1 Δ LPS (for modules 1 and 4, respectively,
534 with modules made by overlap PCR) and an additional module 5 for the downstream
535 region of the *Psa* BV1 Δ LPS by PCR-amplification. These were cloned into
536 pICH41021 with the cloning primers (Supplementary Table 2) designed to
537 synonymously mutate internal *Bsa*I, *Eco*53kI, and *Pst*I restriction enzyme sites
538 (Supplementary Figure S2). These five LPS-KI modules in pICH41021 were then
539 single-pot cloned (by golden gate assembly) into destination pK18B-GG to generate
540 the LPS-KI vector: pLPSbv3. The pLPSbv3 vector was transformed into *E. coli*
541 DH5 α , plated on kanamycin LB agar and screened by nested colony PCR for correct
542 assembly. Positive transformants were confirmed by Sanger sequencing (Macrogen,
543 South Korea). *Psa* BV1 Δ LPS was transformed with pLPSbv3 and transformants
544 were selected on LB agar with nitrofurantoin, cephalixin and kanamycin. Selected
545 colonies were subsequently streaked onto LB agar containing 10% (w/v) sucrose to
546 counter-select plasmid integration, resulting in revertants to knock-out or LPS-
547 complemented mutants. Complementation knock-in mutants were screened using
548 colony PCR with primers for *Psa*J_LPS-KO_Check-F and *Psa*J_LPS-KO_Check-R,
549 as well as gene-specific primers and sent for Sanger sequencing. Complemented
550 mutants (Δ LPS +LPS_BV3) and non-complemented knock-out revertants (Δ LPS-
551 LPSrev) were further confirmed by plating on kanamycin-containing agar to confirm
552 loss of the *sacB* gene.

553

554 ***In planta* growth of *Psa***

555 *Psa* infection assays were based on those published previously with some
556 modifications (McAtee *et al.*, 2018). *A. chinensis* Planch. var. *chinensis* ‘Zesy002’ or
557 ‘Hort16A’ plantlets, grown from axillary buds on Murashige and Skoog rooting
558 medium without antibiotics in sterile 400-mL plastic tubs (pottles) with three plantlets
559 per pottle, were purchased (Multiflora, Auckland, NZ). Plantlets were kept at 20°C
560 under Gro-Lux fluorescent lights under long-day conditions (16 h light: 8 h dark) and
561 used when the plantlets were between 10–14 weeks old. Overnight liquid cultures of
562 wild-type or mutant strains of *Psa* were pelleted at 6000 *g*, re-suspended in 500 mL
563 of 10 mM MgSO₄ to an A₆₀₀ = 0.05 (~10⁷ CFU/mL, determined by plating). Surfactant
564 Silwet™ L-77 (Lehle Seeds, Round Rock, USA) was added to the inoculum at
565 0.0025% (v/v) to facilitate leaf wetting. Pottles of ‘Zesy002’ or ‘Hort16A’ plantlets
566 were filled with the inoculum, submerging the plantlets for 3 min, drained, sealed,
567 and then incubated under previously described plant growth conditions. To assess *in*
568 *planta* growth, leaf samples of four leaf discs per replicate, removed with a 1-cm
569 diameter cork-borer, were taken at 2 h (day 0), day 6, and day 12 post-inoculation.
570 All four pseudo-biological replicates per treatment were taken from the same pottle.
571 To estimate *Psa* growth inside the plant, the leaf discs were surface sterilized, placed
572 in Eppendorf tubes containing three sterile stainless-steel ball bearings and 350 µL
573 10 mM MgSO₄, and macerated in a Storm 24 Bullet Blender (Next Advance, NY,
574 USA) for two bursts of 1 min each at maximum speed. A 10-fold dilution series of the
575 leaf homogenates was made in sterile 10 mM MgSO₄ until a dilution of 10⁻⁸ and each
576 dilution was plated as 10 µL droplets on LB agar supplemented with nitrofurantoin

and cephalixin. After 2 days of incubation at 20°C, the CFU per cm² of leaf area was ascertained from dilutions.

Hierarchical clustering and dendrogram analysis

Hierarchical cluster analysis was conducted in R (R-CoreTeam, 2018) using the ward.D2 method, as per the clustering analysis in Baltrus et al. (2019). The ‘phylogram’ vignette was used to convert hierarchical clusters into dendrograms (Paradis and Schliep, 2018). The tanglegram function in the ‘dendextend’ package was used to visually compare dendrograms by connecting matching leaf node labels with lines (Galili, 2015). The ladder function was used to untangle dendrograms by rotating the tree branches at their nodes without altering their topology, allowing for clear visualization (Galili, 2015). The cor.dendlist function was used to calculate the cophenetic correlation coefficient for the compared dendrograms (Galili, 2015). Cophenetic correlation values (correlation coefficients) range from -1 to 1, with values close to 0 indicating that the compared dendrograms are not statistically similar. The resulting cophenetic correlation matrix was then visualized using the ‘corrplot’ package (Wei and Simko, 2017).

ACKNOWLEDGEMENTS

This work was funded (including a post-doctoral fellowship to JJ) by the Bio-protection Research Centre (Tertiary Education Commission). LMH would like to thank Zespri International for an MSc scholarship.

We would like to thank Rick Broadhurst (AgResearch, Ruakura, New Zealand) for rabbit immunology. We would also like to thank Jo Bowen (PFR) and Iain Hay (UoA) for critically reading the manuscript.

REFERENCES

- Aziz, R.K., Bartels, D., Best, A.A., DeJongh, M., Disz, T., Edwards, R.A. *et al.* (2008) The RAST server: Rapid annotations using subsystems technology. *BMC Genomics*, **9**, 75.
- Baltrus, D.A., Clark, M., Smith, C. and Hockett, K.L. (2019) Localized recombination drives diversification of killing spectra for phage-derived syringacins. *The ISME Journal*, **13**, 237–249.
- Boratyn, G.M., Camacho, C., Cooper, P.S., Coulouris, G., Fong, A., Ma, N. *et al.* (2013) BLAST: A more efficient report with usability improvements. *Nucleic Acids Research*, **41**, W29-33.
- Butler, M.I., Stockwell, P.A., Black, M.A., Day, R.C., Lamont, I.L. and Poulter, R.T.M. (2013) *Pseudomonas syringae* pv. *actinidiae* from recent outbreaks of kiwifruit bacterial canker belong to different clones that originated in China. *PLoS One*, **8**(2), e57464.
- Cunty, A., Poliakoff, F., Rivoal, C., Cesbron, S., Saux, M.L., Lemaire, C. *et al.* (2015) Characterization of *Pseudomonas syringae* pv. *actinidiae* (Psa) isolated from France and assignment of Psa biovar 4 to a *de novo* pathovar: *Pseudomonas syringae* pv. *actinidifoliorum* pv. Nov. *Plant Pathology*, **64**, 582-596.
- Cury, J., Jové, T., Touchon, M., Néron, B. and Rocha, E.P.C. (2016) Identification and analysis of integrons and cassette arrays in bacterial genomes. *Nucleic Acids Research*, **44**, 4539-4550.
- Cuthbertson, L., Kimber, M.S. and Whitfield, C. (2007) Substrate binding by a bacterial ABC transporter involved in polysaccharide export. *Proceedings of the National Academy of Sciences*, **104**(49), 19529-34.
- DebRoy, C., Fratamico, P.M., Yan, X., Baranzoni, G., Liu, Y., Needleman, D.S. *et al.* (2016) Comparison of O-antigen gene clusters of all O-serogroups of *Escherichia coli* and proposal for adopting a new nomenclature for O-typing. *PLoS ONE*, **11**, e0147434.
- Dillon, M.M., Almeida, R.N.D., Laflamme, B., Martel, A., Weir, B.S., Desveaux, D. and Guttman, D.S. (2019a) Molecular evolution of *Pseudomonas syringae* Type III secreted effector proteins. *Frontiers in Plant Science*, **10**, 418.
- Dillon, M.M., Thakur, S., Almeida, R.N.D., Wang, P.W., Weir, B.S. and Guttman, D.S. (2019b) Recombination of ecologically and evolutionarily significant loci maintains genetic cohesion in the *Pseudomonas syringae* species complex. *Genome Biology*, **20**(1), 3.
- Domingues, S., da Silva, G.J. and Nielsen, K.M. (2012) Integrons: Vehicles and pathways for horizontal dissemination in bacteria. *Mobile Genetic Elements*, **2**, 211-223.
- Engler, C., Gruetzner, R., Kandzia, R. and Marillonnet, S. (2009) Golden gate shuffling: A one-pot DNA shuffling method based on type IIS restriction enzymes. *PLoS ONE*, **4**, e5553.
- Ey, P.L., Prowse, S.J. and Jenkin, C.R. (1978) Isolation of pure IgG1, IgG2a and IgG2b immunoglobulins from mouse serum using protein A-Sepharose. *Immunochemistry*, **15**, 429-436.
- Frampton, R.A., Acedo, E.L., Young, V.L., Chen, D.N., Tong, B., Taylor, C. *et al.* (2015) Genome, proteome and structure of a T7-like bacteriophage of the kiwifruit canker phytopathogen *Pseudomonas syringae* pv. *actinidiae*. *Viruses-Basel*, **7**, 3361-3379.
- Frampton, R.A., Taylor, C., Moreno, A.V.H., Visnovsky, S.B., Petty, N.K., Pitman, A.R. and Fineran, P.C. (2014) Identification of bacteriophages for biocontrol of the kiwifruit canker phytopathogen *Pseudomonas syringae* pv. *actinidiae*. *Applied and Environmental Microbiology*, **80**, 2216-2228.
- Fujikawa, T. and Sawada, H. (2016) Genome analysis of the kiwifruit canker pathogen *Pseudomonas syringae* pv. *actinidiae* biovar 5. *Scientific Reports*, **6**, 21399.
- Galili, T. (2015) Dendextend: An R package for visualizing, adjusting and comparing trees of hierarchical clustering. *Bioinformatics* **31**, 3718-3720.
- Ghequire, M.G.K. and De Mot, R. (2015) The tailocin tale: Peeling off phage tails. *Trends in Microbiology*, **23**, 587-590.
- Gillings, M.R. (2014) Integrons: Past, present, and future. *Microbiology and Molecular Biology Reviews* **78**, 257-277.

- Hockett, K.L., Clark, M., Scott, S. and Baltrus, D.A. (2017) Conditionally redundant bacteriocin targeting by *Pseudomonas syringae*. *BioRxiv*, 167593.
- Hockett, K.L., Renner, T. and Baltrus, D.A. (2015) Independent co-option of a tailed bacteriophage into a killing complex in *Pseudomonas*. *MBio*, **6**, e00452.
- Jann, B., Reske, K. and Jann, K. (1975) Heterogeneity of lipopolysaccharides: Analysis of polysaccharide chain lengths by sodium dodecylsulfate-polyacrylamide gel electrophoresis. *European Journal of Biochemistry*, **60**, 239-246.
- Kandel, P., Baltrus, D.A. and Hockett, K.L. (2019) Evidence of tailocin persistence and resistance in *Pseudomonas*. *BioRxiv*, 719799.
- King, J.D., Kocincova, D., Westman, E.L. and Lam, J.S. (2009) Review: Lipopolysaccharide biosynthesis in *Pseudomonas aeruginosa*. *Innate immunity*, **15**, 261-312.
- Koh, Y., Cha, B., Chung, H. and Lee, D. (1994) Outbreak and spread of bacterial canker in kiwifruit. *Korean Journal of Plant Pathology*, **10**, 68-72.
- Kutschera, A., Schombel, U., Wröbel, M., Gisch, N. and Ranf, S. (2019) Loss of wbpI disrupts O-polysaccharide synthesis and impairs virulence of plant-associated *Pseudomonas* strains. *Molecular Plant Pathology*, **20**(11), 1535–1549.
- Kvitko, B.H. and Collmer, A. (2011) Construction of *Pseudomonas syringae* pv. *tomato* DC3000 mutant and polymutant strains. In *Plant Immunity: Methods and Protocols* (McDowell, J.M., ed), pp. 109-128. Totowa, NJ: Humana Press.
- Lam, J., Taylor, V., Islam, S., Hao, Y. and Kocincová, D. (2011) Genetic and functional diversity of *Pseudomonas aeruginosa* lipopolysaccharide. *Frontiers in Microbiology*, **2**, 118.
- Lerouge, I. and Vanderleyden, J. (2002) O-antigen structural variation: Mechanisms and possible roles in animal/plant–microbe interactions. *FEMS Microbiology Reviews*, **26**, 17-47.
- Liston, S.D., Clarke, B.R., Greenfield, L.K., Richards, M.R., Lowary, T.L. and Whitfield, C. (2015) Domain interactions control complex formation and polymerase specificity in the biosynthesis of the *Escherichia coli* O9a antigen. *Journal of Biological Chemistry*, **290**, 1075-1085.
- Maier, R.J. and Brill, W.J. (1978) Involvement of *Rhizobium japonicum* O antigen in soybean nodulation. *Journal of Bacteriology*, **133**, 1295-1299.
- Maldonado, R.F., Sá-Correia, I. and Valvano, M.A. (2016) Lipopolysaccharide modification in gram-negative bacteria during chronic infection. *FEMS Microbiology Reviews*, **40**, 480-493.
- Mann, E., Kelly, S.D., Al-Abdul-Wahid, M.S., Clarke, B.R., Ovchinnikova, O.G., Liu, B. and Whitfield, C. (2019) Substrate recognition by a carbohydrate-binding module in the prototypical ABC transporter for lipopolysaccharide O antigen from *Escherichia coli* O9a. *Journal of Biological Chemistry*, **294**, 14978-14990.
- Marcelletti, S., Ferrante, P., Petriccione, M., Firrao, G. and Scortichini, M. (2011) *Pseudomonas syringae* pv. *actinidiae* draft genomes comparison reveal strain-specific features involved in adaptation and virulence to *Actinidia* species. *PLoS ONE*, **6**, 17.
- Mazzaglia, A., Studholme, D.J., Taratufolo, M.C., Cai, R.M., Almeida, N.F., Goodman, T. et al. (2012) *Pseudomonas syringae* pv. *actinidiae* (Psa) isolates from recent bacterial canker of kiwifruit outbreaks belong to the same genetic lineage. *PLoS ONE*, **7**, 11.
- McAtee P.A., Brian L., Curran B., van der Linden O., Nieuwenhuizen N.J., Chen X. et al. (2018) Re-programming of *Pseudomonas syringae* pv. *actinidiae* gene expression during early stages of infection of kiwifruit. *BMC Genomics*, **19**(1), 822.
- McCann, H.C., Li, L., Liu, Y., Li, D., Hui, P., Zhong, C. et al. (2017) The origin and evolution of a pandemic lineage of the kiwifruit pathogen *Pseudomonas syringae* pv. *actinidiae*. *Genome Biology and Evolution*, **9**, 932-944.
- McCann, H.C., Rikkerink, E.H.A., Bertels, F., Fiers, M., Lu, A., Rees-George, J. et al. (2013) Genomic analysis of the kiwifruit pathogen *Pseudomonas syringae* pv. *actinidiae* provides insight into the origins of an emergent plant disease. *PLoS Pathogens*, **9**, e1003503.

- Mesarich, C.H., Rees-George, J., Gardner, P.P., Ghomi, F.A., Gerth, M.L., Andersen, M.T. *et al.* (2017) Transposon insertion libraries for the characterization of mutants from the kiwifruit pathogen *Pseudomonas syringae* pv. *actinidiae*. *PLoS ONE*, **12**, e0172790.
- Ovod, V., Rudolph, K. and Krohn, K. (1997a) Demonstration of the immunological diversity of O-chains of lipopolysaccharide of *Pseudomonas syringae* and inferring of the serogroup- and serotype-specific epitopes with monoclonal antibodies. In *Pseudomonas syringae pathovars and related pathogens* (Rudolph K., Burr T.J., Mansfield J.W., Stead D., Vivian A. and J., v.K., eds), pp. 532-537. Dordrecht: Springer.
- Ovod, V., Rudolph, K. and Krohn, K. (1997b) Serological classification of *Pseudomonas syringae* pathovars based on monoclonal antibodies towards the lipopolysaccharide-O-chains. In *Pseudomonas syringae pathovars and related pathogens* (Rudolph K., Burr T.J., Mansfield J.W., Stead D., Vivian A. and J., v.K., eds), pp. 526-531. Dordrecht: Springer.
- Paradis, E. and Schliep, K. (2018) Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, **35**, 526-528.
- Pilkington, S.M., Crowhurst, R.N., Hilario, E., Nardoza, S., Fraser, L., Peng, Y. *et al.* (2018) A manually annotated *Actinidia chinensis* var. *chinensis* (kiwifruit) genome highlights the challenges associated with draft genomes and gene prediction in plants. *BMC Genomics*, **19**, 257.
- Pinheiro, L.A.M., Pereira, C., Frazão, C., Balcão, V.M. and Almeida, A. (2019) Efficiency of phage $\phi 6$ for biocontrol of *Pseudomonas syringae* pv. *syringae*: An *in vitro* preliminary study. *Microorganisms*, **7**, 286.
- R-CoreTeam (2018) R: A language and environment for statistical computing. R foundation for statistical computing. <https://www.R-project.org>, Vienna.
- Raetz, C.R. and Whitfield, C. (2002) Lipopolysaccharide endotoxins. *Annual Review of Biochemistry*, **71**, 635-700.
- Riley, M.A. and Wertz, J.E. (2002) Bacteriocins: Evolution, ecology, and application. *Annual Reviews of Microbiology*, **56**, 117-137.
- Rooney, W.M., Grinter, R.W., Correia, A., Parkhill, J., Walker, D.C. and Milner, J.J. (2019) Engineering bacteriocin-mediated resistance against the plant pathogen *Pseudomonas syringae*. *Plant Biotechnology Journal*, 1-11.
- Sambrook, J. and Russell, D.W. (2001) *Molecular cloning: A laboratory manual*, 3rd edition edn Cold Spring Harbour.
- Sawada, H., Kondo, K. and Nakaune, R. (2016) Novel biovar (biovar 6) of *Pseudomonas syringae* pv. *actinidiae* causing bacterial canker of kiwifruit (*Actinidia deliciosa*) in Japan. *Japanese Journal of Phytopathology*, **82**, 101-115.
- Schafer, A., Tauch, A., Jager, W., Kalinowski, J., Thierbach, G. and Puhler, A. (1994) Small mobilizable multipurpose cloning vectors derived from the *Escherichia coli* plasmids pk18 and pk19 - selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene*, **145**, 69-73.
- Scortichini, M., Marcelletti, S., Ferrante, P., Petriccione, M. and Firrao, G. (2012) *Pseudomonas syringae* pv. *actinidiae*: A re-emerging, multi-faceted, pandemic pathogen. *Molecular Plant Pathology*, **13**, 631-640.
- Serizawa, S., Ichikawa, T., Takikawa, Y., Tsuyumu, S. and Goto, M. (1989) Occurrence of bacterial canker of kiwifruit in Japan description of symptoms, isolation of the pathogen and screening of bactericides. *Japanese Journal of Phytopathology*, **55**, 427-436.
- Takikawa, Y., Serizawa, S., Ichikawa, T., Tsuyumu, S. and Goto, M. (1989) *Pseudomonas syringae* pv. *actinidiae* pv. Nov the causal bacterium of canker of kiwifruit in Japan. *Japanese Journal of Phytopathology*, **55**, 437-444.
- Vanneste, J.L., Yu, J., Cornish, D.A., Tanner, D.J., Windner, R., Chapman, J.R. *et al.* (2013) Identification, virulence, and distribution of two biovars of *Pseudomonas syringae* pv. *actinidiae* in New Zealand. *Plant Disease*, **97**, 708-719.

Wei, T. and Simko, V. (2017) R package "corrplot": Visualization of a correlation matrix (version 0.84) <https://github.com/taiyun/corrplot>.

Westphal, O. and Jann, K. (1965) Bacterial lipopolysaccharide: Extraction with phenol-water and further application of the procedure. *Methods in Carbohydrate Chemistry*, **5**, 83–91.

Wyres, K.L., Gorrie, C., Edwards, D.J., Wertheim, H.F.L., Hsu, L.Y., Van Kinh, N. et al. (2015) Extensive capsule locus variation and large-scale genomic recombination within the *Klebsiella pneumoniae* clonal group 258. *Genome Biology and Evolution*, **7**, 1267-1279.

Zdorovenko, E.L., Ovod, V.V., Zatonsky, G.V., Shashkov, A.S., Kocharova, N.A. and Knirel, Y.A. (2001) Location of the O-methyl groups in the O-polysaccharide of *Pseudomonas syringae* pv. *phaseolicola*. *Carbohydrate Research*, **330**, 505-510.

Zdorovenko, G.M. and Zdorovenko, E.L. (2010) *Pseudomonas syringae* lipopolysaccharides: Immunochemical characteristics and structure as a basis for strain classification. *Microbiology*, **79**, 47-57.

FIGURE LEGENDS

Figure 1. The common polysaccharide antigen (CPA) locus from *Pseudomonas syringae* pv. *actinidiae* (Psa) and *P. syringae* pv. *actinidifoliorum*. **(A)** The genetic structure of the CPA locus in Psa ICMP18884 (BV3). The four operons annotated are: the D-Rha biosynthesis operon (dark blue), CPA transport, extension, and termination (TET) operon (green), synthesis of unknown sugar operon (red), L-Rha synthesis operon (light blue), neighbouring syntenic genes that are not believed to be part of this functional cluster are shaded grey or yellow. **(B)** Comparative analysis of the conserved D-Rha biosynthesis and polymorphic TET operons of Psa and Pfm. The CPA locus beginning at the syntenic conserved tRNA-Met through *wbdA* of the TET locus for representatives of the Psa biovars (BV1, ICMP 9617; BV2, ICMP 19071; BV3, ICMP 18884; BV5, MAFF 212063; and BV6, MAFF 212141) and Pfm lineages (L1, ICMP 18803; L3, ICMP 18807) aligned in Geneious v10 with consensus identity displayed above the aligned sequences. Flanking DnaJ and MeCP genes are annotated in grey, D-Rha biosynthesis operon and its genes are annotated in dark blue, and TET operon and its genes annotated in green. **(C)** UPGMA tree of the CPA locus from 5' region up to and including *wzm* in (A). **(D)** UPGMA tree of the CPA locus from 3' region from *wzt* to *wbdA* in (B).

Figure 2. LPS profiles of *Pseudomonas syringae* pv. *actinidiae* (Psa) and *P. syringae* pv. *actinidifoliorum* (Pfm). **(A)** Pro-Q™ Emerald 300 staining of lipopolysaccharide (LPS) preparations on a NuPAGE 4–12% gradient bis-tricine SDS gel from representatives of Pfm lineages (L3 ICMP 18807, lane 2; L1 ICMP 18803, lane 8), Psa biovars (BV1 ICMP 9617, lane 3; BV2 ICMP 19071, lane 4; BV3 ICMP 18884, lane 5; BV5 MAFF212063, lane 6; BV6 MAFF 212141, lane 7), with *Escherichia coli* standards (Std; lanes 1 and 9). A total of 0.5–15 µL LPS extract for each sample (normalized by trial runs) was applied per lane. **(B)** Western blot of LPS extracts from representatives of Psa biovars and Pfm lineages run on a NuPAGE 4–12% gradient bis-tricine SDS gel, obtained by probing with polyclonal antibodies to Psa BV3 ICMP 18884 **(Bi)**, Psa BV1 ICMP 9617 **(Bii)**, or Pfm L1 ICMP 18803 **(Biii)** in a 5000:1 ratio.

Figure 3. Effects of CPA TET operon swap on LPS banding and immunological recognition in *Pseudomonas syringae* pv. *actinidiae* (Psa). **(A)** Genetic polymorphism of the TET operon for Psa BV1 (ICMP 9617) versus BV3 (ICMP 18884). Genetic sequences aligned in Geneious v10. Genome sequences are indicated by light grey bars with associated translated regions as black bars, operons are annotated in light blue, genes in light green with gene names indicated, coding sequences in red with predicted biochemical function indicated, the original TET operon from BV1 indicated in dark green with upstream (KO-5') and downstream (KO-3') regions for generating the LPS TET operon knock-out in grey, and the TET operon swap region by knock-in from BV3 indicated in dark blue. **(B)** LPS profiles of Psa BV3 (lane 1 and lane 8), BV1 (lane 2), BV1 TET operon knockout (Δ LPS; lane 3), four BV1-to-BV3 TET operon complemented isolates (Δ LPS+LPS_BV3 1-2, 1-4, 2-1, and 2-5; lanes 4–7), and one BV1 revertant to knock-out isolates (Δ LPS-LPSrev 1-3; lane 9). Pro-Q™ Emerald 300 staining of LPS preparations on a 4–12% gradient Tris-glycine SDS gel. LPS extract (15 µL) for each sample was applied per lane. **(C)**

Immuno-blot of LPS extracts from wild-type, knock-out (KO), KO-complemented, and KO-revertant strains run on a gradient SDS-PAGE gel as for (A), probed with polyclonal antibodies to *Psa* BV3 in a 5000:1 ratio.

Figure 4. The CPA TET operon is required for full pathogenicity of *Pseudomonas syringae* pv. *actinidiae* (*Psa*) in host plants. **(A)** *Psa* BV3 or BV3 R-LPS mutants (Mesarich *et al.*, 2017) were flood-inoculated at $\sim 10^6$ CFU/mL on *Actinidia chinensis* var. *chinensis* 'Zesy002', and bacterial growth was determined at 0, 6, and 12 days post-inoculation. Error bars represent standard error of the mean from four pseudo-biological replicates. Asterisks indicate results of Student's *t*-test between selected sample and wild-type; **($P < 0.01$), ***($P < 0.001$). The experiment was conducted three times with similar results. **(B)** *Psa* BV1, BV1 LPS knock-out mutant (Δ LPS), two independent BV1 knock-outs complemented LPS from BV3 (+LPS_BV3), and two independent BV1 revertant-to-knock-out LPS (-LPSrev) were flood-inoculated at $\sim 10^6$ CFU/mL on *A. chinensis* var. *chinensis* 'Hort16A', and bacterial growth was determined at 0, 6, and 12 days post-inoculation. Error bars represent standard error of the mean from four replicates. Asterisks indicate results of Student's *t*-test between selected sample and wildtype (*Psa* V13); ***($P < 0.001$). The experiment was conducted twice with similar results.

Figure 5. (A) Phylogenetic comparison of TET operons from various *Pseudomonas syringae* strains. Amino acid sequences from three proteins of the CPA locus (Wzt, WbdD and WbdA) were aligned in Geneious v10 and neighbour-joining phylogenetic trees built for their TET operons. Strains were grouped into numbered clades as indicated. Strains are coloured according to their core genome phylogenies as previously described (Dillon *et al.*, 2019b): phylogroup 1 blue; phylogroup 2 red; phylogroup 3 green; phylogroup 4 orange; phylogroup 5 purple; phylogroup 6 yellow; phylogroup 7 pink; phylogroup 9 grey; phylogroup 10 mauve; phylogroup 11 chocolate; phylogroup 13 claret. **(B)** LPS profiles of *Pseudomonas syringae* representing phylogenetic clades for the TET operon. Pro-QTM Emerald 300 staining of lipopolysaccharide (LPS) preparations on a 4–12% gradient Tris-glycine SDS gel from representatives of *P. syringae* strains marked with asterisks from Supplementary Table 1 (top to bottom, lanes 1–11), with *Escherichia coli* control (Std, lane 12). Banding patterns were grouped into clades on their designations from Figure 7. A total of 2–15 μ L LPS extract for each sample (normalized by trial runs) was applied per lane.

Figure 6. Tanglegram comparison of tailocin sensitivity, TET operon, and core gene MLST phylogenies. **(A)** Tanglegram of sensitivity matrix phylogeny (left) and TET operon phylogeny (right). **(B)** Tanglegram of sensitivity matrix phylogeny (left) and core gene MLST phylogeny (right). **(C)** Tanglegram of TET operon phylogeny (left) and core gene MLST phylogeny (right). Hierarchical clustering analysis using the ward.D2 method was used to build a tailocin sensitivity phylogeny from the sensitivity matrix provided in Baltrus *et al.* (2019), which is transposed from a syringacin killing matrix. The TET operon phylogeny was produced by using the TET operon genome sequences of selected strains, assembled into a phylogeny using the UPGMA clustering method, as was the MLST phylogeny. The co-phenetic entanglement coefficients for each comparison are indicated below each panel title. Axes in each

panel represent Euclidean distance. Colours represent broad clade groupings, with the cut-offs being Euclidean distances of 10, 0.5, and 0.02 for the sensitivity, TET, and MLST phylogenies, respectively. This Euclidean cut-off for the MLST branches allows representative separation by phylogroup for the strains used.

Supplementary Figure S1. MLST genome phylogenetic tree for *Pseudomonas syringae* and *Pseudomonas viridiflava* strains. Five genes (*gyrB*, *ropD*, *gapA*, *pgi* and *acnB*) from selected representatives of *P. syringae* phylogroups were concatenated and aligned in Geneious v11 and neighbour-joining phylogenetic trees built. Strains grouped as expected into their phylogroups and are coloured according to their core genome phylogenies as previously described (Dillon *et al.*, 2019b): phylogroup 1 blue; phylogroup 2 red; phylogroup 3 green; phylogroup 4 orange; phylogroup 5 purple; phylogroup 6 yellow; phylogroup 7 pink; phylogroup 9 grey; phylogroup 10 mauve; phylogroup 11 chocolate; phylogroup 13 claret.

Supplementary Figure S2. Schematic for knock-out and knock-in swap of the TET locus in *Pseudomonas syringae* pv. *actinidiae* (Psa) BV1. One-kilobase flanking regions upstream (UP) and downstream (DN) of the polymorphic region of the TET operon in wild-type (WT) Psa BV1 (green; encoding Wzt, WbdD, and WbdA proteins in red) were PCR-cloned and fused at a primer-introduced internal *Xba*I site and used in pK18mobsacB vector-mediated knock-out of the BV1 TET locus to generate BV1_LPS-KO. To generate the BV1-to-BV3 LPS swap (knock-in complementation), a pK18mobsacB vector-mediated method was also used. First, *Eco*53kI, *Bsa*I and *Pst*I sites were removed during PCR (primer-driven) to facilitate blunt-end cloning of modules, golden gate (GG) cloning, and evasion of the BV1-encoded restriction enzyme, respectively. The knock-in pK18mobsacB vector was built (GG assembly) from five modules with sizes indicated. Modules 2, 3, and 5 were PCR-amplified and blunt-end cloned into pICH41021 shuttle vector directly. Modules 1 and 4 were first cloned as two parts (A and B) from their respective Psa BVs and then fused by overlap PCR mediated by primer sequences designed to introduce overlap regions during amplification of part A and B. Following overlap PCR, modules 1 and 4 were blunt-end cloned into the pICH41021 shuttle vector. The modules were then assembled into the final pK18mobsacB knock-in construct. The BV1_LPS-KO was then subsequently complemented with TET operon from BV3 (blue).

Supplementary Figure S3. Bacterial cell sedimentation for *Pseudomonas syringae* pv. *actinidiae* (Psa) in inoculation buffer. (A) Psa BV3 and two R-LPS mutants, $\Delta wbpL$ and Δgmd (Mesarich *et al.*, 2017) were grown overnight in LB liquid culture, pelleted by centrifugation, resuspended in 10 mM MgSO₄ buffer, and normalized to OD₆₀₀ of 0.4. (B) Psa BV3 and BV1 (wild-type; WT), Psa BV1 LPS knock-out (ΔLPS), four BV1-to-BV3 TET operon-complemented isolates ($\Delta LPS + LPS_{BV3}$: #1-2, #1-4, #2-1, and #2-5; green), and four BV1 revertant to knock-out isolates ($\Delta LPS - LPS_{rev}$: #1-3, #1-6, #2-4, and #2-8; red) were grown overnight in LB liquid culture, pelleted by centrifugation, resuspended in 10 mM MgSO₄ buffer, and normalized to OD₆₀₀ of 0.5. Photographs of cell suspensions in both (A) and (B) were taken immediately after resuspension in buffer (T0) and after an hour standing at room temperature (T1).

Supplementary Figure S4. Colony morphologies for *Pseudomonas syringae* pv. *actinidiae* (Psa) BV1 mutant and complemented strains. Psa BV3 mutants ($\Delta wbpL$ or Δgmd) (Mesarich *et al.*, 2017) and BV1 (wild-type; WT), Psa BV1 LPS knock-out (ΔLPS), four BV1-to-BV3 TET operon complemented isolates ($\Delta LPS+LPS_{BV3}$: #1-2, #1-4, #2-1, and #2-5; green), and four BV1 revertant to knock-out isolates ($\Delta LPS-LPS_{rev}$: #1-3, #1-6, #2-4, and #2-8; red) were grown on LB agar plates, for 3 days and photographs of colonies were taken under a stereomicroscope.

Supplementary Figure S5. The CPA TET operon is required for symptom development of *Pseudomonas syringae* pv. *actinidiae* (Psa) in host plants. **(A)** Psa BV3 or BV3 R-LPS mutants ((Mesarich *et al.*, 2017)) were flood-inoculated at $\sim 10^6$ CFU/mL on *Actinidia chinensis* var. *chinensis* 'Zesy002', and photographs of symptom development in pottles and representative leaves taken at 50 days post-inoculation. **(B)** Psa BV1, BV1 LPS knock-out mutant (ΔLPS), or two independent BV1 knock-outs complemented LPS from BV3 ($+LPS_{BV3}$) mutants were flood-inoculated at $\sim 10^6$ CFU/mL on *A. chinensis* var. *chinensis* 'Hort16A', and photographs of symptom development in pottles and representative leaves taken at 50 days post-inoculation.

Supplementary Figure S6. Cophenetic entanglement coefficients for phylogenetic trees for the TET operon and MLST analysis generated using different tree-building methods (UPGMA, Neighbor joining (NJ), or RAxML) compared to each other and to the R-syringacin (tailocin) sensitivity tree for data from Baltrus *et al.* (2019). The cophenetic correlation matrix between the trees have values ranging between -1 to 1, with near 0 values indicating that the two trees are not statistically similar. Values between -0.25 and 0.25 (indicated by the red box on the scale) are poorly correlated and outlined in the table and associated scale. UPGMA was selected over RAxML and NJ for tree-building in Fig. 6 due to the nature of comparing the sensitivity phenotype with genetic similarity treated as a 'phenotype' (thus creating phenetic trees rather than phylogenetic trees). This allowed for pairwise comparisons instead of inferring an evolutionary relationship that NJ and RAxML trees are better at conveying.

Figure 1




Figure 2






Figure 3

A



B



C





Figure 4

A

Psa BV3 in planta growth assay



B

Psa BV1 in planta growth assay




Figure 5A




Figure 5B






Figure 6





Supplementary Figure S1




Supplementary Figure S2



BV1_LPS (WT):




BV1_LPS-
KI-BV3
(GG):




Supplementary Figure S3



Supplementary Figure S4



Supplementary Figure S5



Supplementary Figure S6




Table 1. Loci involved in lipopolysaccharide (LPS) biosynthesis in *Pseudomonas syringae* pv. *actinidiae* BV3.

LPS component	Psa BV3 gene locus-tag	<i>P. aeruginosa</i> PAO1 orthologs	Amino Acid identity	Gene name	Psa R-LPS phenotype ¹
Lipid A	IYO_000255	PA0011	80%	<i>lpxL</i>	N/A
Lipid A	IYO_007660-75	PA3643-6	55-63%	<i>lpxA-D</i>	N/A
Lipid A	IYO_016185-225	PA3552-59	87-94%	<i>arnA-F, arnT</i>	N/A
Core	IYO_025525-640	PA5001-12, PA4988,	66-86%		Yes
CPA	IYO_005330	PA5448	37%	<i>wbpY</i>	Yes
CPA	IYO_008990-9000	PA3141, PA3145-6	40-74%	<i>wbpK-M</i>	Yes
CPA	IYO_022945-023030	PA5447-59, PA5322	55-80% ²		Yes

¹ R-LPS, Rough LPS.

² D- and L-Rha biosynthetic genes only.

Table 2. Gene annotations of *Pseudomonas syringae* pv. *actinidiae* BV3 common LPS locus.

Gene annotation	<i>P. aeruginosa</i>	<i>E. coli</i>	Function	Gene ID	Protein_ID	R-LPS ¹
dTDP-glucose 4,6-dehydratase		<i>rmIB</i>	L-Rha biosynthesis	IYO_022945	AKT32330.1	
dTDP-4-dehydrorhamnose reductase		<i>rmID</i>	L-Rha biosynthesis	IYO_022950	AKT32331.1	
glucose-1-phosphate thymidyltransferase		<i>rmIA</i>	L-Rha biosynthesis	IYO_022955	AKT32332.1	
hypothetical protein			unknown	IYO_022960	AKT32333.1	
hypothetical protein			unknown	IYO_022965	AKT32334.1	
amine oxidase			unknown	IYO_022970	AKT32335.1	
membrane protein			unknown	IYO_022975	AKT32336.1	
glucose-1-phosphate cytidyltransferase		<i>rfbF</i>	synthesis of unknown sugar	IYO_022980	AKT32337.1	
glycosyl transferase			synthesis of unknown sugar	IYO_022985	AKT32338.1	
NAD-dependent epimerase/dehydratase		<i>galE</i>	synthesis of unknown sugar	IYO_022990	AKT32339.1	
CDP-glucose 4,6-dehydratase		<i>rfbG</i>	synthesis of unknown sugar	IYO_022995	AKT32340.1	
glycosyl transferase family 1	<i>wbpX</i>	<i>wbdA</i>	chain elongation	IYO_023000	AKT32341.1	
SAM-dependent methyltransferase		<i>wbdD</i>	chain termination	IYO_023005	AKT32342.1	Yes
sugar ABC transporter ATP-binding protein	<i>wzt</i>	<i>wzt</i>	transport	IYO_023010	AKT32343.1	Yes
ABC transporter	<i>wzm</i>	<i>wzm</i>	transport	IYO_023015	AKT32344.1	Yes

GDP-D-mannose dehydratase	<i>gmd</i>	<i>gmd</i>	D-Rha biosynthesis	IYO_023020	AKT32345.1	Yes
GDP-6-deoxy-D-lyxo-4-hexulose reductase	<i>rmd</i>	<i>rmd</i>	D-Rha biosynthesis	IYO_023025	AKT32346.1	
glycosyl transferase family 1	<i>wbpZ</i>		D-Rha biosynthesis	IYO_023030	AKT32347.1	Yes

¹ R-LPS, Rough LPS.

Supplementary Table 1. Metadata for isolates used in this study

<i>P. syringae</i> isolate	Abbreviation	Strain	Host	Country of isolation	Date	GenBank accession
<i>P. marginalis</i> pv. <i>marginalis</i>	<i>Pmar</i>	ICMP 3553PT	<i>Cichorium intybus</i>	New Zealand	1949	RBPW01
<i>P. syringae</i>	<i>Ps</i>	CC1417	Epilithon	USA		AVEO02
<i>P. syringae</i>	<i>Ps</i>	CC1524	Stream water	France		AVEK02
<i>P. syringae</i>	<i>Ps</i>	CC1583	Epilithon	France		AVEF02
<i>P. syringae</i>	<i>Ps</i>	G33C	<i>Actinidia</i> sp.	New Zealand		
<i>P. syringae</i>	<i>Ps</i>	ICMP 11296	<i>Actinidia deliciosa</i>	New Zealand	1991	RBRK01
<i>P. syringae</i>	<i>Ps</i>	ICMP 19498	<i>Actinidia chinensis</i>	New Zealand	2010	LKCH01
<i>P. syringae</i>	<i>Ps</i>	ICMP 3272	<i>Actinidia deliciosa</i>	New Zealand	1971	LKEK01, RBQZ01
<i>P. syringae</i>	<i>Ps</i>	UB246	Environmental	France		AVEQ01
<i>P. syringae</i>	<i>Ps</i>	CC1416	Epilithon	USA		AVEP02
<i>P. syringae</i>	<i>Ps</i>	CC1458	02	USA		AVEN02
<i>P. syringae</i>	<i>Ps</i>	CC1466	<i>Dodecatheon pulchellum</i>	USA		AVEM02
<i>P. syringae</i>	<i>Ps</i>	CC1513	<i>Pritzelago alpina</i>	France		AVEL02
<i>P. syringae</i>	<i>Ps</i>	CC1544	Lake water	France		AVEI02
<i>P. syringae</i>	<i>Ps</i>	CC1557	Environmental (snow)	France	ND	CP007014-5
<i>P. syringae</i>	<i>Ps</i>	CC1559	Environmental (snow)	France		AVEG02
<i>P. syringae</i>	<i>Ps</i>	CC1630	<i>Onobrychis viciifolia</i>	USA		AVED02
<i>P. syringae</i>	<i>Ps</i>	CC440	Cantaloupe	France		AVEC02
<i>P. syringae</i>	<i>Ps</i>	CC457	Cantaloupe	France		AVEB02
<i>P. syringae</i>	<i>Ps</i>	CC94	Cantaloupe	France		AVEA02
<i>P. syringae</i>	<i>Ps</i>	ICMP 11168	<i>Actinidia deliciosa</i>	New Zealand	1991	LKGV01

<i>P. syringae</i>	<i>Ps</i>	ICMP 11292	<i>Actinidia deliciosa</i>	New Zealand	1991	LKGU01
<i>P. syringae</i>	<i>Ps</i>	ICMP 11293	<i>Actinidia deliciosa</i>	New Zealand	1991	LKEP01
<i>P. syringae</i>	<i>Ps</i>	ICMP 13102	<i>Actinidia deliciosa</i>	France	1985	LKEO01
<i>P. syringae</i>	<i>Ps</i>	ICMP 19198	<i>Actinidia</i> sp.	New Zealand	2011	RBRZ01
<i>P. syringae</i>	<i>Ps</i>	ICMP 19499	<i>Actinidia</i> sp.	New Zealand	2010	LKCI01
<i>P. syringae</i>	<i>Ps</i>	ICMP 3690	<i>Prunus persica</i>	United Kingdom		LKBV01
<i>P. syringae</i>	<i>Ps</i>	ICMP 4122	<i>Prunus armeniaca</i>	New Zealand	1974	LKBX01
<i>P. syringae</i>	<i>Ps</i>	TLP2	Potato	USA		JGI 2507262033
<i>P. syringae</i>	<i>Ps</i>	UB303	Lake water	France		AVDZ
<i>P. syringae</i>	<i>Ps</i>	USA007	Stream water	USA		AVDY02
<i>P. syringae</i>	<i>Ps</i>	USA011	Stream water	USA		CP045799-01
<i>P. syringae</i>	<i>Ps</i>	CC1543	Lake water	France		AVEJ02
<i>P. syringae</i> pv. <i>actinidiae</i> BV1	<i>Psa</i> BV1	ICMP 9853	<i>Actinidia deliciosa</i>	Japan	1984	CP018202-4
<i>P. syringae</i> pv. <i>actinidiae</i> BV2	<i>Psa</i> BV2	ICMP 19071	<i>Actinidia chinensis</i>	Korea	1997	AOJS01, RBSG00
<i>P. syringae</i> pv. <i>actinidiae</i> BV3	<i>Psa</i> BV3	ICMP 18884	<i>Actinidia deliciosa</i>	New Zealand	2010	CP011972-3
<i>P. syringae</i> pv. <i>actinidiae</i> BV5	<i>Psa</i> BV5	MAFF 212063	<i>Actinidia chinensis</i>	Japan	2012	CP024712-4
<i>P. syringae</i> pv. <i>actinidiae</i> BV6	<i>Psa</i> BV6	MAFF 212141	<i>Actinidia deliciosa</i>	Japan	2015	MSBX01
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i> L1	ICMP 18802	<i>Actinidia chinensis</i>	New Zealand	2010	MUKM01
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i> L1	ICMP 18803	<i>Actinidia chinensis</i>	New Zealand	2010	AOKK01
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i> L3	ICMP 18807	<i>Actinidia deliciosa</i>	New Zealand	2010	ANJL01, AOKG01
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i>	ICMP 19497	<i>Actinidia chinensis</i>	New Zealand	ND	LKBQ01

<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i>	ICMP 19486	<i>Actinidia chinensis</i>	Australia	1990	
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i> L4	CFBP 8039	<i>Actinidia deliciosa</i>	France		LJJM01
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i> L2	CFBP 8043	<i>Actinidia deliciosa</i>	France		LJFM01
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i> L4	CFBP 8160	<i>Actinidia deliciosa</i>	France		LJJL01
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i> L1	CFBP 8161	<i>Actinidia deliciosa</i>	France		LJFL01
<i>P. syringae</i> pv. <i>actinidifoliorum</i>	<i>Pfm</i> L1	CFBP 8180	<i>Actinidia deliciosa</i>	France		LJFN01
<i>P. syringae</i> pv. <i>aesculi</i>	<i>Pae</i>	0893_23	Horse chestnut	United States		AEAD01
<i>P. syringae</i> pv. <i>amylgdali</i>	<i>Pam</i>	ICMP 3918	<i>Prunus dulcis</i>	Greece	1967	LJPQ01
<i>P. syringae</i> pv. <i>antirrhini</i>	<i>Pat</i>	ICMP 4959	<i>Antirrhinum majus</i>	United Kingdom	1965	RBQK01
<i>P. syringae</i> pv. <i>aptata</i>	<i>Ptt</i>	G733	<i>Oryza sativa</i>	ND	1976	RBOI01
<i>P. syringae</i> pv. <i>aptata</i>	<i>Ptt</i>	DSM50252				AEAN01
<i>P. syringae</i> pv. <i>atrofaciens</i>	<i>Paf</i>	DSM50255	Wheat			AWUI01
<i>P. syringae</i> pv. <i>avellanae</i>	<i>Pav</i>	BPIC 631	<i>Corylus avellana</i>	Greece	1976	ATDK01
<i>P. syringae</i> pv. <i>avellanae</i>	<i>Pav</i>	ISPave037	<i>Corylus avellana</i>			AKCK01
<i>P. syringae</i> pv. <i>berberidis</i>	<i>Pbe</i>	ICMP 13291	<i>Berberidis</i> sp.	New Zealand	1995	RBQP01
<i>P. syringae</i> pv. <i>cannibina</i>	<i>Pcb</i>	ICMP 2823	<i>Cannabis sativa</i>	Hungary	1957	LJPX01, FNKU01
<i>P. syringae</i> pv. <i>cerasicola</i>	<i>Pce</i>	ICMP 13929	<i>Prunus x yedoensis</i>	Japan	1996	LKCB01
<i>P. syringae</i> pv. <i>cerasicola</i>	<i>Pce</i>	ICMP 17525	<i>Prunus x yedoensis</i>	Japan		LKCC01, RBTI01
<i>P. syringae</i> pv. <i>cichorii</i>	<i>Pci</i>	ICMP 6917	<i>Carthamus tinctorius</i>	New Zealand	1980	RBRY01
<i>P. syringae</i> pv. <i>delphinii</i>	<i>Pde</i>	ICMP 529	<i>Delphinium</i> sp.	New Zealand	1957	LJQH01
<i>P. syringae</i> pv. <i>glycinea</i>	<i>Pgy</i>	ICMP 2236	<i>Glycine max</i>	New Zealand		RBRO01
<i>P. syringae</i> pv. <i>glycinea</i>	<i>Pgy</i>	2411	<i>Glycine max</i>	New Zealand	1971	QJTT01

<i>P. syringae</i> pv. <i>glycinea</i>	<i>Pgy</i>	ICMP 2189	<i>Glycine max</i>	New Zealand	1968	LJQL01
<i>P. syringae</i> pv. <i>glycinea</i> Race 4	<i>Pgy</i>	A29-2				ADWY01
<i>P. syringae</i> pv. <i>lachrymans</i>	<i>Pla</i>	NM002	Cucumber	China	1984	CP020351.1
<i>P. syringae</i> pv. <i>lachrymans</i>	<i>Pla</i>	98A-744	<i>Cucumis sativus</i>			LCWT01
<i>P. syringae</i> pv. <i>lachrymans</i>	<i>Pla</i>	MAFF 302278				AEAM01
<i>P. syringae</i> pv. <i>lachrymans</i>	<i>Pla</i>	MAFF370057			1979	JGI 2505313044
<i>P. syringae</i> pv. <i>lachrymans</i>	<i>Pla</i>	MAFF301315	Cucumber			CP031225-8
<i>P. syringae</i> pv. <i>lapsa</i>	<i>Plp</i>	ATCC 10859	<i>Triticum aestivum</i>	ND	2014	CP013183
<i>P. syringae</i> pv. <i>lapsa</i>	<i>Plp</i>	ICMP 3947	<i>Zea sp.</i>	ND	ND	LJQQ01
<i>P. syringae</i> pv. <i>maculicola</i>	<i>Pma</i>	90-32	<i>Brassica oleracea</i>	United States	1990	LGLH01
<i>P. syringae</i> pv. <i>maculicola</i>	<i>Pma</i>	ICMP 11281	<i>Brassica rapa</i>	China	ND	RBUQ01
<i>P. syringae</i> pv. <i>mori</i>	<i>Pmo</i>	ICMP 4468	<i>Morus alba</i>	New Zealand	1957	RBRW01
<i>P. syringae</i> pv. <i>mori</i>	<i>Pmo</i>	MAFF 301020	Mulberry	Japan	1966	AEAG01
<i>P. syringae</i> pv. <i>morsprunorum</i>	<i>Pmp</i>	NZIPFRPS-6		New Zealand	2010	LKCD01
<i>P. syringae</i> pv. <i>morsprunorum</i>	<i>Pmp</i>	MAFF302280				AEAE01
<i>P. syringae</i> pv. <i>oryzae</i>	<i>Por</i>	ICMP 9093	<i>Oryza sativa</i>	Japan	1983	RBSZ01
<i>P. syringae</i> pv. <i>oryzae</i>	<i>Por</i>	1_6	<i>Oryza sativa</i>		1991	ABZR01
<i>P. syringae</i> pv. <i>persicae</i>	<i>Ppe</i>	ICMP 3706	<i>Prunus cerasifera</i>	New Zealand	1966	RBQE01
<i>P. syringae</i> pv. <i>phaseolicola</i>	<i>Pph</i>	1448A	<i>Phaseolus vulgaris</i>	Ethiopia	1985	CP000058-60
<i>P. syringae</i> pv. <i>phaseolicola</i>	<i>Pph</i>	1644	<i>Vigna radiata</i>			AGAS01
<i>P. syringae</i> pv. <i>pisi</i>	<i>Ppi</i>	1704B				AEAI0
<i>P. syringae</i> pv. <i>pisi</i>	<i>Ppi</i>	PP1	<i>Pisum sativum</i>	Japan	1978	CP034078-81
<i>P. syringae</i> pv. <i>savastoni</i>	<i>Psv</i>	NCPPB 3335	<i>Olea europaea</i>	France	ND	CP008742.1
<i>P. syringae</i> pv. <i>syringae</i>	<i>Psy</i>	B48	<i>Prunus persica</i>	United States		LGKT01
<i>P. syringae</i> pv. <i>syringae</i>	<i>Psy</i>	5264	<i>Prunus avium</i>	United Kingdom	2017	NBAQ01

<i>P. syringae</i> pv. <i>syringae</i>	<i>Psy</i>	B728a	<i>Phaseolus vulgaris</i>	United States	1987	CP000075.1
<i>P. syringae</i> pv. <i>syringae</i>	<i>Psy</i>	PD 2766	<i>Actinidia</i> sp.	United States		LKEM01
<i>P. syringae</i> pv. <i>syringae</i>	<i>Psy</i>	PD 2774	<i>Actinidia</i> sp.	United States		LKEL01
<i>P. syringae</i> pv. <i>syringae</i>	<i>Psy</i>	B301D	<i>Pyrus communis</i>	United Kingdom	1959	CP005969
<i>P. syringae</i> pv. <i>tabaci</i>	<i>Pta</i>	6605	Tobacco	Japan		AJXI01
<i>P. syringae</i> pv. <i>tabaci</i>	<i>Pta</i>	ICMP 2835	<i>Nicotiana tabacum</i>	Hungary	1959	LJRL01
<i>P. syringae</i> pv. <i>tagetis</i>	<i>Ptg</i>	ICMP 4092	<i>Tagetes erecta</i>	United kingdom	1970	RBQC01
<i>P. syringae</i> pv. <i>tagetis</i>	<i>Ptg</i>	ICMP 6369	<i>Tagetes erecta</i>	Australia	1976	RBVF01
<i>P. syringae</i> pv. <i>thea</i>	<i>Pth</i>	NCPPB 2598	<i>Thea sinensis</i>	Japan	1974	AGNN01
<i>P. syringae</i> pv. <i>tomato</i>	<i>Pto</i>	DC3000	<i>Solanum lycopersicum</i>	United Kingdom	1960	AE016853-5
<i>P. syringae</i> pv. <i>tomato</i>	<i>Pto</i>	ICMP 2844	<i>Solanum lycopersicum</i>	United Kingdom	1960	LJRN01
<i>P. syringae</i> pv. <i>tomato</i>	<i>Pto</i>	T1		Canada	ND	ABSM01
<i>P. syringae</i> pv. <i>tremae</i>	<i>Ptr</i>	CC1629	Oats	USA		AVEE02
<i>P. viridiflava</i>	<i>Pvir</i>	CC1582	Epilithon	France		AVDW01
<i>P. viridiflava</i>	<i>Pvir</i>	ICMP 2848	<i>Phaseolus vulgaris</i>	Switzerland	1927	LJRS01, LKEH01
<i>P. viridiflava</i>	<i>Pvir</i>	ICMP 8820	<i>Prunus persica</i>	USA	1984	LKCA01
<i>P. viridiflava</i>	<i>Pvir</i>	TA043	<i>Primula officinalis</i>	France	ND	AVDV01
<i>P. viridiflava</i>	<i>Pvir</i>	CDRTc14	<i>Lepidium draba</i>	Austria	2013	MBPF01
<i>P. viridiflava</i>	<i>Pvir</i>	ICMP 11289	<i>Actinidia deliciosa</i>	New Zealand	1991	LKGX01
<i>Pseudomonas</i> spp.	<i>P. spp.</i>	ICMP 10191	<i>Actinidia</i> sp.	China	1981	LKGW01

Supplementary Table 2. The sequence of primers used in this study. All primers were synthesized by Macrogen, South Korea.

Primer name	Sequence (5'-3')
PsaJ_LPS-KO_UP-F	CGATTCAAAGACTAAACGGTGACT
PsaJ_LPS-KO_UP-R	GGTCTCTCTAGAAGTACATCCACCATTTCGG
PsaJ_LPS-KO_DN-F	GGTCTCTCTAGAGTTTGAATTGTGCGTCGCTCA
PsaJ_LPS-KO_DN-R	CTGCTTGAGTTCCTGAGGCC
PsaJ_LPS-KO_Check-F	ATGAATAATTCGTCTATAGGGTGCCG
PsaJ_LPS-KO_Check-R	CAGTTCGAGAACATCTCTGACCG
GG-lacZ-InFusion-F	GCGTTGGCCGATTCATTACAATTATCGATACAATGGGAGAC
GG-lacZ-InFusion-R	TTGCGGCAGCGTGAAGCATATCTCATTAAAGCAGGACAAGC
Bv3LPS_Bv1-OL1-F	GAAATGGTGGATGTACTATGAGCCTGTTGAGTGTC
Bv3LPS_Bv1-OL1-R	GAACTCAACAGGCTCATAGTACATCCACCATTTC
Bv3LPS_Bv1-OL2-F	GAAAAGTATGCGGTCTGAGAATTGTGCGTCGCTCAA
Bv3LPS_Bv1-OL2-R	TTGAGCGACGCACAATTCTCAGACCGCATACTTTTC
Bv3LPS-mod1-F	GGTCTCGAATGAATAATTCGTCTATAGGGTGCCG
Bv3LPS-mod1-R	GGTCTCACAAGCCGGATAAAGCTCTTGGA
Bv3LPS-mod2-F	GGTCTCGCTTGCAGAGCCCTTGATCTTGG
Bv3LPS-mod2-R	GGTCTCAGCACGGTGTGGAAACGCGAA
Bv3LPS-mod3-F	GGTCTCGGTGCAATTGCTACAGGAAAACACGC
Bv3LPS-mod3-R	GGTCTCAACTGCTCCGCGCGTTGCAGA
Bv3LPS-mod4-F	GGTCTCGCAGTTCTCATGGGATCGCACC
Bv3LPS-mod4-R	GGTCTCAGGGTGTGGTTTCAGGCGC
Bv3LPS-mod5-F	GGTCTCGACCCTCTCTTTTTGAAGAGCTGG
Bv3LPS-mod5-R	GGTCTCAAAGCCAGTTCGAGAACATCTCTGACCG