

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**BLOOD PRESSURE AND THE EFFECT OF CALCIUM
ENRICHED MILK IN HUMANS WITH NORMAL OR
MILDLY ELEVATED BLOOD PRESSURE:
METHODOLOGICAL CONSIDERATIONS**

A thesis presented in partial fulfilment of
the requirements for the degree of
MASTER OF SCIENCE
in Physiology
at Massey University

BRETT MICHAEL GUTHRIE

2001

ACKNOWLEDGEMENTS

Thanks must go, firstly, to the participants, without whom there would have been no trial. They persevered while we took numerous measurements.

Thank you to Dr Hilary Green who allowed me to attach myself to the trial. Thanks also for her patience while I tried to make sense of the data that we did obtain and for reading many attempts at a “thesis” that were often changed before she had a chance to reply.

Thanks to the General Practitioner whose original idea prompted this study, and who became a subject as well. The cold medicine was also appreciated.

Dr Gordon Reynolds deserves mention for turning a blind eye while I worked on this thesis, when, perhaps, I should have been doing other things.

Thanks to Jill Richards for allowing me on her “turf”, and to Chris Booth for data entry which saved me valuable time.

Finally, many, many thanks to Sarah who ensured that I really did finish, and without whom I would not have got this far.

ABSTRACT

The aim of the present study was to assess the influence of three different milk products on blood pressure in patients with mild hypertension who were not under medication or had ceased medication for the trial period under medical supervision. In addition, the effect of calcium on blood pressure was to be explored.

The initial objectives of the trial were not met because the trial was stopped due to an adverse coronary event in one volunteer. However, there was sufficient data to reinforce the validity of the methodology, especially the ambulatory blood pressure recording, and to ensure that all other aspects of the trial were achievable.

Subjects with elevated blood pressure could be recruited and undergo a series of tests to determine their physical parameters, have blood pressure taken by a variety of methods and on several occasions, and meaningful data obtained. In addition, a small, potentially beneficial, modification could be made to their diet, and the effect of this dietary change monitored by both blood pressure and blood profile changes. These changes were made with minimal disruption to their daily routine, and were generally well-received.

The present study confirmed that ambulatory blood pressure monitoring could be conducted with little intrusion into the lives of the subjects. The data obtained from a variety of methods was able to identify those subjects who presented with some of the cluster of factors which characterise Syndrome X.

These results, while only from a small sample group, strongly support the use this research methodology to provide an accurate representation of a population subgroup, such as those with elevated blood pressure. In addition, the effect of a dietary intervention on blood pressure and blood lipid profiles can be monitored in free-living subjects.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS.....	I
ABSTRACT.....	II
LIST OF TABLES.....	XI
LIST OF FIGURES.....	XIII
CHAPTER ONE: INTRODUCTION.....	1
CHAPTER TWO: LITERATURE REVIEW	3
2.1. BLOOD PRESSURE	3
2.1.1. Blood pressure regulation.....	3
2.1.2. Pulse pressure	3
2.1.3. Diurnal variation.....	4
2.1.4. Morning surge.....	4
2.2. HYPERTENSION	5
2.2.1. Types of Hypertension.....	6
2.2.1.1. Primary hypertension.....	6
2.2.1.2. Secondary hypertension.....	7
2.2.1.3. Pregnancy-induced hypertension	7
2.2.1.4. Malignant hypertension	7
2.2.1.5. White coat hypertension	8
2.2.3. Effects of hypertension	9
2.2.4. Treatment of hypertension	9
2.2.4.1. Lifestyle changes	10
2.2.4.2. Nutrition in the prevention of hypertension.....	11
2.2.4.3. Pharmaceutical therapy.....	12
2.3. MINERALS AND BLOOD PRESSURE.....	12
2.3.1. Calcium.....	13
2.3.1.1. Plasma calcium	13
2.3.1.2. Calcium regulation.....	13
2.3.1.3. Parathyroid hormone	14
2.3.1.4. Calcitonin	15
2.3.2. Calcium and blood pressure	15

2.3.2.1. Dietary calcium.....	16
2.3.2.2. Calcium supplementation	17
2.3.2.3. Calcium absorption.....	18
2.3.2.4. Bioavailability.....	20
2.3.2.5. Abnormal calcium metabolism	20
2.4. OTHER MINERALS.....	21
2.4.1. Sodium	21
2.4.2. Potassium.....	22
2.4.3. Phosphorous.....	22
2.5. DAIRY PRODUCTS	23
2.5.1. Introduction.....	23
2.5.2. Milk.....	23
2.5.3. Another Dairy Factor?.....	24
2.5.4. Conclusion.....	24
2.6. BLOOD LIPIDS.....	25
2.7. SYNDROME X.....	28
CHAPTER 3: METHODOLOGY	30
3.1. BLOOD PRESSURE MONITORING.....	30
3.1.1. Ambulatory blood pressure monitoring.....	30
3.1.1.1. Inconvenience	32
3.2. BODY COMPOSITION.....	33
3.2.1. Anthropometry.....	33
3.2.1.1. Weight for height.....	36
3.2.1.2. BMI Criteria.....	37
3.2.1.3. Circumferences	37
3.2.1.3.1. Waist to hip ratio.....	37
3.2.1.3.2. Mid-arm circumference.....	38
3.2.1.3.3. Mid-arm muscle circumference.....	39
3.2.1.3.4. Arm muscle area.....	39
3.2.1.4. Skinfolds.....	40
3.2.1.4.1. Measurement	41
3.2.1.4.2. Derivations	42
3.2.2. Bioelectrical Impedance Analysis.....	42
3.2.2.1. Total body water	42
3.2.2.2. Assumptions.....	43

3.2.2.3. Validation	43
3.2.2.4. Hydration constant	44
3.3. FITNESS TESTING	44
3.3.1. <i>Strength test</i>	45
CHAPTER FOUR: PRELIMINARY TRIAL	47
4.1. INTRODUCTION	47
4.2. METHODS	47
4.2.1. <i>Statistical Analysis</i>	48
4.3. RESULTS	48
4.3.1. <i>24-hour ABP and HR (individual)</i>	48
4.3.2. <i>Day ABP and HR (individual)</i>	49
4.3.3. <i>Night ABP and HR (individual)</i>	49
4.3.4. <i>Dipping (individual)</i>	50
4.3.5. <i>Males and females</i>	51
4.3.6. <i>24-hour ABP (pooled)</i>	51
4.3.7. <i>Day ABP and HR (pooled)</i>	53
4.3.8. <i>Night ABP and HR (pooled)</i>	53
4.3.9. <i>Dipping (pooled)</i>	53
4.3.10. <i>Smoking</i>	54
4.4. DISCUSSION	55
4.4.1. <i>Subjects</i>	55
4.4.2. <i>Sleep disruption</i>	55
4.4.3. <i>Night</i>	56
4.4.4. <i>Dipping</i>	56
4.4.5. <i>Males and females</i>	57
4.4.6. <i>Over-reading?</i>	57
4.5. CONCLUSION	57
CHAPTER FIVE: MAIN TRIAL	59
5.1. INTRODUCTION	59
5.2. METHODS	59
5.2.1. <i>Subject recruitment</i>	59
5.2.2. <i>Physical data</i>	60
5.2.2.1. <i>Weight</i>	60

5.2.2.2. Height	60
5.2.2.3. Triceps skinfolds	61
5.2.2.4. Circumferences	61
5.2.2.5. Body composition	61
5.2.2.6. Fitness test	61
5.2.2.7. Strength test	62
5.2.3. <i>Blood lipids</i>	62
5.2.4. <i>24-hour food recall</i>	63
5.2.5. <i>Calcium intake</i>	64
5.2.6. <i>Energy expenditure</i>	64
5.2.7. <i>Urine collection</i>	64
5.2.8. <i>Estimation of cardiovascular risk</i>	65
5.2.9. <i>Blood pressure</i>	65
5.2.9.1. Clinic and home blood pressure	65
5.2.9.2. On-site blood pressure	66
5.2.10. <i>Analysis by case study</i>	66
5.2.11. <i>Statistical analysis</i>	67
5.3. SUBJECT DEMOGRAPHICS	67
5.4. SUBJECT ONE	68
5.4.1. <i>Physical data</i>	68
5.4.2. <i>Blood pressure</i>	68
5.4.2.1. Blood pressure differences	70
5.4.3. <i>Blood profile</i>	70
5.4.4. <i>Estimation of cardiovascular risk</i>	71
5.4.5. <i>Reported food intake</i>	71
5.4.6. <i>Basal metabolic rate</i>	72
5.4.7. <i>Activity</i>	72
5.4.8. <i>Summary</i>	72
5.5. SUBJECT TWO	72
5.5.1. <i>Physical data</i>	72
5.5.2. <i>Blood pressure</i>	73
5.5.2.1. Blood pressure differences	74
5.5.3. <i>Blood profile</i>	75
5.5.4. <i>Estimation of cardiovascular risk</i>	75
5.5.5. <i>Reported food intake</i>	75

5.5.6. Basal metabolic rate.....	76
5.5.7. Activity.....	76
5.5.8. Summary.....	77
5.6. SUBJECT THREE.....	77
5.6.1. Physical data.....	77
5.6.2. Blood pressure.....	78
5.6.3. Blood profile.....	79
5.6.4. Estimation of cardiovascular risk.....	79
5.6.5. Reported food intake.....	79
5.6.6. Basal metabolic rate.....	80
5.6.7. Activity.....	80
5.6.8. Summary.....	80
5.7. SUBJECT FOUR.....	81
5.7.1. Physical data.....	81
5.7.2. Blood pressure.....	81
5.7.3. Blood profile.....	82
5.7.4. Estimation of cardiovascular risk.....	83
5.7.5. Reported food intake.....	83
5.7.6. Basal metabolic rate.....	84
5.7.7. Activity.....	84
5.7.8. Summary.....	84
5.8. SUBJECT FIVE.....	84
5.8.1. Physical data.....	84
5.8.2. Blood pressure.....	85
5.8.2.1. Blood pressure differences.....	86
5.8.3. Blood profile.....	87
5.8.4. Estimation of cardiovascular risk.....	87
5.8.5. Reported food intake.....	87
5.8.6. Basal metabolic rate.....	88
5.8.7. Summary.....	88
5.9. SUBJECT SIX.....	89
5.9.1. Physical data.....	89
5.9.2. Blood pressure.....	89
5.9.3. Blood profile.....	90

5.9.4. Estimation of cardiovascular risk.....	91
5.9.5. Reported food intake	91
5.9.6. Basal metabolic rate.....	91
5.9.7. Summary	92
5.10. SUBJECT SEVEN	92
5.10.1. Physical data	92
5.10.2. Blood pressure.....	93
5.10.3. Blood profile.....	93
5.10.4. Estimation of cardiovascular risk.....	94
5.10.5. Reported food intake	94
5.10.6. Basal metabolic rate	95
5.10.7. Summary	95
5.11. SUBJECT EIGHT	95
5.11.1. Physical data	95
5.11.2. Blood pressure.....	96
5.11.3. Blood profile.....	97
5.11.4. Estimation of cardiovascular risk.....	98
5.11.5. Reported food intake	98
5.11.6. Basal metabolic rate	98
5.11.7. Summary	99
5.12. SUBJECT NINE	99
5.12.1. Physical data	99
5.12.2. Blood pressure.....	100
5.12.3. Blood profile.....	100
5.12.4. Estimated cardiovascular risk.....	101
5.12.5. Reported food intake	101
5.12.6. Basal metabolic rate	102
5.12.7. Summary	102
5.13. SUBJECT TEN.....	102
5.13.1. Physical data	102
5.13.2. Blood pressure.....	103
5.13.3. Blood profile.....	104
5.13.4. Estimation of cardiovascular risk.....	104
5.13.5. Reported food intake	104

5.13.6. Basal metabolic rate	105
5.13.7. Summary	105
5.14. SUBJECT ELEVEN	106
5.14.1. Physical data	106
5.14.2. Blood pressure.....	106
5.14.3. Blood profile.....	107
5.14.4. Estimation of cardiovascular risk.....	107
5.14.5. Reported food intake	108
5.14.6. Basal metabolic rate	108
5.14.7. Summary.....	109
5.15. DISCUSSION.....	109
5.15.1. Subjects.....	109
5.15.2. Physical.....	109
5.15.3. Fitness.....	110
5.15.4. Basal metabolic rate	110
5.15.5. Blood pressure.....	110
5.15.6. Intervention.....	110
5.15.7. Blood profile.....	110
5.15.8. 24-hour food intake recall	111
5.15.9. Mineral deficiencies	111
5.15.9.1. Calcium	111
5.15.9.2. Magnesium.....	111
5.15.9.3. Potassium	111
5.15.9.4. Phosphorous.....	112
5.15.10. Cardiovascular risk.....	112
5.15.11. Conclusion.....	112
CHAPTER SIX: GENERAL DISCUSSION & CONCLUSION	113
6.1. BLOOD PRESSURE.....	113
6.1.1. Blood pressure recording methods.....	113
6.1.2. Use of the ambulatory apparatus	113
6.1.3. White coat hypertension.....	114
6.1.4. Activity records.....	115
6.2. BODY COMPOSITION	115
6.2.1. Anthropometry.....	115

6.2.2. <i>Bioelectrical impedance analysis</i>	116
6.3. HYPERTENSION	116
6.4. INTERVENTION	117
6.5. MINERAL DEFICIENCIES	117
6.6. CONCLUSION	118
REFERENCES	119
APPENDIX 1	139
APPENDIX 2	140

LIST OF TABLES

Table 2.1. Blood pressure and stages of hypertension _____	5
Table 2.2. Blood lipid concentrations _____	26
Table 2.3. Relative Risk of Coronary Heart Disease for people 50-80 years of age	27
Table 3.1. Maximal oxygen uptake (ml/kg/min) _____	45
Table 4.1. Individual 24-hour ABP and HR means \pm SEM _____	48
Table 4.2. Individual day ABP and HR means \pm SEM _____	49
Table 4.3. Individual night ABP and HR means \pm SEM _____	50
Table 4.4. Percentage dipping calculated from night MBP change from the day MBP. _____	50
Table 5.4.1. Summary of physical data _____	68
Table 5.4.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline (1); mid-intervention (2) and post-intervention (3) _____	69
Table 5.4.3. Comparison of blood pressure (mm Hg) changes \pm SD from baseline mean _____	70
with post-intervention mean _____	70
Table 5.4.4. Blood lipid concentrations _____	70
Table 5.4.6. Reported food intake percentage of RDI _____	71
Table 5.4.7. BMR by different methods (kcal/day) _____	72
Table 5.5.1. Summary of the physical data _____	73
Table 5.5.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline (1); mid-intervention (2) and post-intervention (3) _____	74
Table 5.5.3. Comparison of blood pressure (mm Hg) changes \pm SD from baseline mean with post-intervention mean _____	74
Table 5.5.4. Blood lipid concentrations _____	75
Table 5.5.5. Reported food intake percentage of RDI _____	76
Table 5.5.6. BMR by different methods (kcal/day) _____	76
Table 5.6.1. Summary of physical data _____	77

Table 5.6.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline (1) and mid-intervention (2)	78
Table 5.6.3. Blood lipid concentrations	79
Table 5.6.4. Reported food intake percentage of RDI.	80
Table 5.6.5. BMR by different methods (kcal/day)	80
Table 5.7.1. Summary of the physical data	81
Table 5.7.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline (1) and mid-intervention (2)	82
Table 5.7.3. Blood lipid concentrations	82
Table 5.7.4. Reported food intake percentage of RDI	83
Table 5.7.5. BMR by different methods (kcal/day)	84
Table 5.8.1. Summary of physical data	85
Table 5.8.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline (1); mid-intervention (2) and post-intervention (3)	86
Table 5.8.3. Comparison of blood pressure changes (mm Hg) \pm SD from baseline mean with post-intervention mean	86
Table 5.8.4. Blood lipid concentrations	87
Table 5.8.5. Reported food intake percentage of RDI	88
Table 5.8.6. BMR by different methods (kcal/day)	88
Table 5.9.1. Summary of physical data	89
Table 5.9.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline (1) and mid-intervention (2)	90
Table 5.9.3. Blood lipid concentrations	90
Table 5.9.4. Reported food intake percentage of RDI	91
Table 5.9.5. BMR by different methods (kcal/day)	92
Table 5.10.1. Summary of physical data	92
Table 5.10.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline	93

Table 5.10.3. Blood lipid concentrations	94
Table 5.10.4. Reported food intake percentage of RDI	94
Table 5.10.5. BMR by different methods (kcal/day)	95
Table 5.11.1. Summary of physical data	96
Table 5.11.2. Baseline means for blood pressure (mm Hg) and HR (bpm) \pm SD	97
Table 5.11.3. Blood lipid concentrations	97
Table 5.11.4. Reported food intake percentage of RDI	98
Table 5.11.5. BMR by different methods (kcal/day)	99
Table 5.12.1. Summary of physical data	99
Table 5.12.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline	100
Table 5.12.3. Blood lipid concentrations	101
Table 5.12.4. Reported food intake percentage of RDI	101
Table 5.12.5. BMR by different methods (kcal/day)	102
Table 5.13.1. Summary of physical data	103
Table 5.13.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline	103
Table 5.13.3. Blood lipid concentrations	104
Table 5.13.4. Reported food intake percentage of RDI	105
Table 5.13.5. BMR by different methods (kcal/day)	105
Table 5.14.1. Summary of physical data	106
Table 5.14.2. Summary of means for blood pressure (mm Hg) and HR (bpm) \pm SD for baseline	107
Table 5.14.3. Blood lipid concentrations	107
Table 5.14.4. Reported food intake percentage of RDI	108
Table 5.14.5. BMR by different methods (kcal/day)	108

LIST OF FIGURES

Figure 2.1. Hormonal control of plasma Ca^{2+}	14
Figure 4.1. Individual recordings comparing males and females.	51
Figure 4.2. Pooled hourly mean \pm SEM SBP/DBP/HR with time lines for ceasing work (17.00h), bedtime (22.00h) and awake (6.00h)	52
Figure 4.3. Pooled mean \pm SEM SBP/DBP with BP 120/80 (mm Hg)	53
Figure 4.4. Subject Five MBP and HR compared to times of smoking for each recording day. Raw data shown.	54

CHAPTER ONE: INTRODUCTION

Elevated blood pressure is an increasing problem in industrialised nations, especially as the population ages (Levy, D., 1999) and becomes more obese (Whelton *et al.*, 1998: New Zealand Guidelines, 2000).

It has been demonstrated by numerous studies that adequate calcium lowers blood pressure (Buonopane *et al.*, 1992; Osbourne *et al.*, 1996; McCarron, 1998; Barr *et al.*, 2000). Sufficient calcium in the diet may have a greater effect on people who have elevated blood pressure. Known as hypertensives, these people usually submit to a drug regime to control their blood pressure. Simply drinking a daily glass of calcium-reinforced milk provides an attractive alternative. Calcium-enriched fat-free milk is currently available on the New Zealand market, as the benefits of sufficient calcium in the diet are well-recognised (Karanja *et al.*, 1994). These benefits are for both osteoporosis and blood pressure reduction (McCarron, 1998). The former relationship is well established, however the latter is somewhat controversial (Seely, 1991; Buonopane *et al.*, 1992; Osbourne *et al.*, 1996; McCarron, 1998; Barr *et al.*, 2000).

Fat offers enhanced flavour to most pallets, therefore there is market resistance to fat-free products. This observation prompted the milk industry to explore the effect of a low-fat calcium-enriched milk that is currently unavailable on the market. The low fat content may offer improved flavour and therefore increased sales, but the health benefits of the calcium may be offset by the increased potential of raised blood cholesterol levels. This effect is especially important for those with elevated blood pressure, who may further compromise their health by consumption of such a higher fat product.

Control of blood lipids is a means of reducing the risk of cardiovascular disease (Karanja *et al.*, 1994). Dairy food consumption has been linked to the development of arteriosclerosis (McNamara, 1992), thus people with an increased risk of cardiovascular disease are usually encouraged to reduce their intake of these foods. However, the strong link between calcium and blood pressure begs the investigation whether the potential benefit of consuming calcium enriched milk outweighs any potential harm of dairy fat.

A nine-week trial was conducted to determine if drinking modified milk powder lowered blood pressure. The purpose of this study was to assess the influence of three different milk products on blood pressure in patients with mild hypertension who were not under medication or had ceased medication for the trial period under medical supervision. The three milk products were normal skim milk; normal skim milk with added calcium; and low fat milk (1.5% fat) with added calcium. These were formulated by the New Zealand Dairy Research Institute. In addition, the effect of the higher fat milk on blood lipids was also to be explored.

The 9-week trial was stopped prematurely after only three subjects had completed the measurements and, therefore, the number of subjects is limited. The purpose of the thesis was revised to assess in more detail the suitability of the various methodologies for this kind of intervention trial.