

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

STUDIES ON THE EFFECT OF SOME NITROGENOUS  
MATERIALS ON THE FOOD INTAKE OF SHEEP

A thesis presented in partial fulfilment of the  
requirements for the degree of  
Doctor of Philosophy  
at Massey University

Arnold Max Bryant

May, 1968

MASSEY UNIVERSITY LIBRARY



1061963846

MASSEY UNIVERSITY  
LIBRARY  
PALLISER BUILDING  
MASSEY PARK, PALMERSTON NORTH

## ACKNOWLEDGEMENTS

Grateful acknowledgement is made to my chief supervisor, Professor D.A. Titchen of the Department of Physiology and Anatomy, Massey University for his interest and guidance. He is also thanked for the surgical preparation of the majority of animals used during the course of this work.

Dr C.S.W. Reid of Plant Chemistry Division, D.S.I.R., is especially thanked for the help and encouragement extended to me both as a supervisor and a friend.

I also wish to thank

1. The Department of Agriculture for granting me study leave on full salary.
2. D.S.I.R. and Massey University for the provision of facilities.
3. The late A.C. Glenday for advice on statistical analysis.
4. The professional and technical staff of D.S.I.R. for their help.
5. L. Douglas and D.H.B. MacQueen of Ruakura Agricultural Research Centre for the preparation of the figures.
6. Mrs Henrickson for typing this thesis.

## CONTENTS

	PREFACE	page 1
Chapter 1	REVIEW OF LITERATURE	
	Introduction	4
	A. The regulation of food intake in non-ruminants	5
	1. Role of the central nervous system	5
	2. Nature of the stimuli and location of the receptors involved in the regulation of food intake by non-ruminants	7
	(a) Role of gastrointestinal tract	7
	(i) Hunger contractions	7
	(ii) Oropharyngeal metering	8
	(iii) Gastric distension	8
	(iv) Gastric emptying	9
	(b) Metabolic and chemostatic hypotheses	14
	(i) Glucostatic	14
	(ii) Thermostatic	16
	(iii) Lipostatic	17
	(iv) The effect of protein on food intake	19
	3. Relationship between food and water intake	22
	B. Regulation of food intake in ruminants	27
	1. Role of the central nervous system	28
	2. Role of the gastrointestinal tract	29
	(a) Oropharyngeal regulation	29
	(b) Gastric distension	30
	(c) Rate of disappearance of digesta from the reticulorumen	32
	(d) Rate of breakdown of digesta in the reticulorumen	32
	(e) Flow of digesta from the reticulorumen	35
	(f) Influence of the gut beyond the reticulorumen	36
	3. Food intake and water metabolism	40
	(a) Relationship between food and water intake	40
	(b) The consistency of rumen contents and food intake	41
	(c) Osmotic control of food intake	42

Chapter		page
	4. Chemostatic regulation of food intake	43
	5. Protein and food intake in ruminants	47
2	SOME EFFECTS OF INTRADUODENAL INFUSIONS OF CASEIN ON FOOD INTAKE	
	Introduction	50
	Materials and methods	51
	Experimental and results	55
	Discussion	73
3	OBSERVATIONS ON THE NATURE AND LOCATION OF THE MECHANISMS INVOLVED IN THE EFFECT OF PROTEINACEOUS INFUSIONS ON FOOD INTAKE	
	Introduction	77
	Materials and methods	79
	Results	83
	Discussion	96
4	THE EFFECT OF INTRADUODENAL INFUSIONS OF CASEIN ON THE URINARY EXCRETION OF SODIUM AND POTASSIUM	
	Introduction	104
	Materials and methods	107
	Results	110
	Discussion	112
5	THE EFFECTS OF SOME NITROGENOUS COMPOUNDS ON RETICULORUMEN MOTILITY	
	Introduction	116
	Materials and methods	117
	Results	119
	Discussion	128
6	GENERAL DISCUSSION	140
	BIBLIOGRAPHY	152

LIST OF FIGURES

Fig. No.		Facing page
1	The food intake of the three sheep in Expt.1	55
2	A comparison of the effects on food intake when casein was infused into the duodenum in three different vehicles.	58
3	Effect of intraduodenal infusions of 750ml. Blaxter's Saline alone and containing 50g casein on (a) food intake and (b) voluntary water intake.	60
4	Effect of intraduodenal infusions of 750ml. Blaxter's Saline containing 0, 50 and 75g casein on (a) plasma amino acid concentration, (b) packed cell volume and (c) blood CO <sub>2</sub> content.	62
5	The mean effects of the treatments of Expt. 8 on food intake compared with those on faecal dry matter output.	69
6	A comparison of the effects of intraduodenal infusions of casein and casein hydrolysate on some blood parameters.	88
7	The effects of intraduodenal infusions of (a) an amino acid mixture and (b) Salt Solution on the circadian pattern of eating and ruminating of one sheep.	90
8	The effects of (a) intraduodenal infusion of casein and (b) the intravenous infusion of casein hydrolysate on the circadian pattern of eating and ruminating.	93
9	Effect of intraduodenal infusions of Blaxter's Saline alone and containing casein and the intravenous infusion of 0.9% NaCl and casein hydrolysate on some blood parameters.	95
10	Changes in the composition of urine and venous blood when intraduodenal infusions of Blaxter's Saline alone and containing 40g casein were made to four sheep.	110
11	The effect of injecting 100ml. water containing (a) 3g and (b) 6g urea into the rumen on the frequency of reticulum contraction, pH and ammonia levels of rumen liquor, CO <sub>2</sub> and urea content of venous blood.	120
12	The effect on frequency of reticulum contraction of a continuous infusion of urea into the rumen.	121
13	Changes in ammonia content and pH of rumen liquor, CO <sub>2</sub> and urea content of venous blood during the experiments shown in Fig.12.	121
14	Inhibition of rumen and reticulum contractions on injection of 200ml. of isotonic NaHCO <sub>3</sub> into the jugular vein.	122

Fig.No.		Facing page
15	Effect on food intake and faecal dry matter output of intraruminal infusions of 750ml. water containing urea equivalent to 0, 7.5 and 10g nitrogen.	123
16	The effect on food intake of infusions of 750ml. water containing urea equivalent to 0, 2.5, 7.5 and 10g nitrogen into the rumen of one sheep.	123
17	The frequency of reticulum contraction and the time spent each hour in eating, ruminating and idling during the four infusion days of the experiment shown in Fig.16.	123
18	Changes in ammonia content and pH levels of rumen liquor, urea, CO <sub>2</sub> and ammonia content of venous blood of the experiment shown in Fig.16 and 17.	125
19	The frequency of reticulum contraction and the circadian pattern of eating and ruminating on the day before, the day of and the day after the intraduodenal infusion of an amino acid mixture to two sheep.	127
20	Relationship between the mean frequency of reticulum contraction and food intake for each of three sheep.	128