

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

A COMPARATIVE STUDY OF THE PHOSPHORUS
CHARACTERISTICS OF OIL PALM VOLCANIC SOILS IN
PAPUA NEW GUINEA AND NEW ZEALAND VOLCANIC SOILS

A thesis presented in partial fulfillment of the requirements for the
Degree of Master of Applied Science in Soil Science at
Massey University, Palmerston North, New Zealand

Murom Banabas

1998

ABSTRACT

Oil Palm (*Elaeis guineensis* Jacq.) grown on volcanic ash soils in Papua New Guinea (PNG) generally respond well to N fertilisers but shows a lack of consistent response to inorganic phosphorus (P) fertilisers. This is true even on soils with high phosphate retention (PR) and where Olsen P values highlighted in the preliminary survey of PNGOPRA field trial data are very low (<10 mg/kg). A notable exception occurs at Bialla (Trial 201) where significant responses (yield and growth parameters) to P fertilisers have been found on soils with very low Olsen P (< 4 mg/kg) and very high PR values (>90% to at least 60 cm depth).

This study was done to characterise the PNG oil palm growing volcanic soils in relation to P responsiveness, to identify P fractions and their relative amounts, to determine the fate of applied P fertilisers and to compare chemical and mineralogical characteristics of PNG soils with some New Zealand (NZ) equivalent soils.

Mineralogical analysis indicates that the PNG soils used in this study are relatively young as evidenced by the presence of very high amounts of readily-weatherable volcanic glass in the sand, silt and clay fractions. Soils at Hoskins, Kapiura and Bialla, all in West New Britain (WNB) Province, contain similar amounts and types of primary and secondary minerals. Soils at Bialla are probably older than those at Hoskins and Kapiura and contain large amounts of secondary amorphous minerals (alophane and ferrihydrite) in the clay fraction. Soils at Popondetta are different from those in WNB with high amounts of hornblende and no augite or hypersthene in the heavy mineral fraction. Allophane levels in the clay fraction are high to very high in soil surface layers at Hoskins and Kapiura and at all depths in Bialla soils. At Popondetta, allophane content is very low at all depths

PR in all soils and at all depths was highly correlated with acid oxalate extractable Al (Al_o) ($r = 0.84^*$) and iron (Fe_o) ($r = 0.89^*$). The sources of these 2 extracts (allophane and ferrihydrite) are largely responsible for the high PR in the soils studied. High allophane and ferrihydrite levels at all depths in Bialla soils correspond well with very

high PR values (>90%) to at least 2 m depth. Low levels of these 2 minerals in Popondetta soils correspond well with low PR values (30%). Intermediate PR values (60 - 70%) for Hoskins and Kapiura surface soils correlates well with the occurrence of intermediate levels of allophane and ferrihydrite.

In all PNG soils, a P fractionation scheme showed that the major P fractions are organic. At Hoskins, NaOH-Po accounts for 38 to 48% of total P. For Kapiura NaOH-Po accounts for approximately 50% of total P, and Bicarb.-Po accounts for 59% of total bicarbonate-extractable P. For Bialla soils, NaOH-Po and Bicarb.-Po comprise between 74 and 76%, on average, of their respective total extracted P for all depths. At Popondetta, NaOH-Po comprises 62% and Bicarb.-Po 63% of their respective total extractable P contents.

P fertiliser accumulation in Hoskins and Kapiura soils occurs mostly in organic forms and within the top 10 cm of soil. At Hoskins, 83% of total added P accumulated in the top 10 cm (53% being NaOH-Po) while 17% was found in the next 10 cm depth (31% being NaOH-Po). At Kapiura, 74% of total accumulated P was found in the top 10 cm of soil (61% being NaOH-Po) and 26% within the 20 - 30 cm layer (81% being NaOH-Po).

The presence of amorphous minerals explains much of the behaviour of P in trial soils, with the major P source/sink in PNG soils being as organic forms.

In relation to soil mineralogical and chemical characteristics, PNG soils were classified into one of the major 3 groups in terms of responsiveness to P fertilisers; (a) soils with very high PR (>90%) and Olsen P values of less than 4 mg/kg which are considered most likely to respond to inorganic P fertilisers e.g. Bialla soil, (b) soils with medium to high PR (60 - 70%) will likely show inconsistent responses to P fertilisers and P responses are most likely to be secondary to N e.g. Hoskins and Kapiura soils and (c) soils with low PR (30 - 40%) which are unlikely to respond to P fertilisers at least in the foreseeable future e.g. Popondetta soils.

This study highlights a future need for further study of the dynamics of P nutrient cycling, specifically the mineralisation rates of organic matter and the release of Pi for plant uptake in PNG oil palm growing soils. Also there is a need to re-establish the leaf critical concentration because in PNG soils though leaf levels are generally less than 0.150% DM, palms do not always respond to P fertilisers. This suggests that the "critical" P concentrations under PNG conditions is probably less than the international standard at 0.150% DM.

Mineralogical and P sorption characteristics of young volcanic ash soils in NZ are sufficiently similar to those in PNG to provide useful information about the general behaviour of P fertilisers and P reaction products in oil palm production systems.

ACKNOWLEDGMENTS

I express my gratitude to the following people for their help in doing this thesis.

Firstly to my 2 supervisors, Dr. Max Turner and Professor Russ Tillman for their encouragement, supervision and critical suggestions.

To the New Zealand Landcare staff; Dr. Roger Parfitt for helpful suggestions and analysis for allophane and ferrihydrite, Mr. Joe Whitton for helping with mineralogical analysis and Ms. Linda Hill for carbon and nitrogen analysis.

To Mr. Bob Toes for assistance with Olsen P tests.

To Soil Science Department technicians for assistance in use of various laboratory equipment.

To Mr. Ian Orrell (DOR-PNGOPRA) for help in making this study possible for me.

To Mr. Graham King and Mr. Allan Oliver for providing literature review materials and Joe Yambun and Doreen for helping in soil sampling and handling. Also to all PNGOPRA workers who actually helped in sampling the soils.

A special thanks to my wife, Nambua for looking after our 3 children; Enzeng, Serah and Yandisao during my absence. Also a special thanks to my parents at home for their never ending support.

Finally, this piece of work though may not be the best, I would like to dedicate it to the 20 people in my village (Mataya, Wantoat Sub-District, Morobe Province, PNG) who were covered by landslide while gardening in 1993.

TABLE OF CONTENTS

ABSTRACT.....	ii
ACKNOWLEDGMENTS.....	v
TABLE OF CONTENTS.....	vi
LIST OF TABLES.....	xii
LIST OF FIGURES.....	xiv

CHAPTER 1

INTRODUCTION.....	1
--------------------------	---

CHAPTER 2 LITERATURE REVIEW

2.1 OIL PALM GROWING SOILS.....	3
2.1.1 Introduction.....	3
2.1.2 Latosols.....	3
2.1.3 Alluvial Soils.....	4
2.1.4 Oil Palm Growing Soils in Papua New Guinea (PNG).....	4
2.1.4.1 Hoskins, Kapiura and Bialla.....	5
2.1.4.2 Popondetta.....	6
2.2 MINERALOGY OF SOIL PARENT MATERIALS.....	7
2.2.1 Primary Mineral Composition of Volcanic Ash Materials	8
2.2.2 Secondary Mineral Composition of Volcanic Ash Soils.....	8
2.2.2.1 Allophane.....	8
2.2.2.2 Imogolite.....	9
2.2.2.3 Halloysite.....	9
2.2.2.4 Ferrihydrite.....	10
2.2.3 Soil Weathering and Formation.....	10
2.2.3.1 Temperature.....	11

2.2.3.2 Rainfall, Leaching and Drainage.....	11
2.2.3.3 Tephra Thickness and Depth of Burial Effect.....	12
2.2.3.4 Organic Cycle Effects.....	12
2.3 REACTIONS OF PHOSPHORUS IN SOILS.....	13
2.3.1 Permanent and Variable Charge Clay Minerals.....	13
2.3.1.1 Permanent Charge.....	13
2.3.1.2 Variable Charge.....	14
2.3.2 Phosphate Adsorption.....	15
2.3.2.1 Adsorption Mechanisms of Phosphates on the Variable Charge Surface Clay Minerals.....	15
2.3.2.1.1 Allophane.....	16
2.3.2.1.2 Iron Oxides.....	17
2.3.3 Phosphorus Desorption.....	17
2.3.4 Phosphate Retention.....	17
2.4 PHOSPHORUS FORMS IN SOILS.....	17
2.4.1 Soil Inorganic P Fractions.....	19
2.4.1.1 Reactions of P Fertilisers in Soils.....	20
2.4.2 Soil Organic P Forms.....	20
2.4.2.1 Factors Influencing Organic P Contents	21
2.4.2.2 Nature of Soil Organic P Compounds.....	21
2.4.2.2.1 Inositol	21
2.4.2.2.2 Phospholipids.....	22
2.4.2.2.3 Nucleic Acids.....	22
2.4.2.3 Organic P Turnover.....	22
2.4.2.3.1 Immobilisation / Mineralisation.....	23
2.4.2.4 Organic-P as P Source for Crops.....	24
2.5 PHOSPHORUS NUTRITION OF OIL PALM.....	25
2.5.1 Oil Palm Responses to Inorganic P Fertilisers.....	25
2.5.1.1 P responses in PNG	25
2.5.1.1.1 Hoskins Trials.....	25
2.5.1.1.2 Kapiura	27
2.5.1.1.3 Bialla	27

2.5.1.1.4 Popondetta.....	27
2.5.1.2 Oil Palm Responses to P Fertiliser Applications in Other Parts of the World.....	28
2.5.2 Physiological Roles of Phosphorus in Oil Palm.....	28
2.5.2.1 Energy Transfer.....	29
2.5.2.2 Genetic Functions.....	29
2.5.2.3 Cell Structural Components.....	29
2.5.2.4 Storage in Seeds.....	30
2.5.3 Effects of Phosphorus Deficiency.....	30
2.5.4 P Concentration in Oil Palm Leaves.....	31
2.5.5 General Description and Functions of Oil Palm Root System	32
2.5.6 The Oil Palm Phosphorus Uptake.....	33
2.5.7 Phosphorus in Soil Solution and Root Rhizosphere	34
2.5.8 Processes Involved in the Movement of Phosphorus to Root/Soil Interface..	35
2.5.9 Absorption of Phosphorus by Plant Roots.....	36
2.5.10 Factors Affecting P Uptake.....	36
2.5.10.1 Changes in P Absorption with Age.....	36
2.5.10.2 Changes with Soil Depth	37
2.5.10.3 Morphological Properties of Roots.....	37
2.5.11 Rhizosphere Microorganisms and Phosphorus Uptake.....	37
2.5.12 Summary.....	38

CHAPTER 3

METHODS AND MATERIALS

3.1 SAMPLING SITES AND SAMPLING PROCEDURES.....	40
3.1.1 Site Descriptions.....	40
3.1.1.1 Hoskins Trial 107.....	40
3.1.1.2 Kapiura Trial 402.....	40
3.1.1.3 Bialla Trials 201 and 205.....	41
3.1.1.4 Popondetta Trial 323.....	41
3.1.2 Sampling Procedure.....	41

3.2 SOIL MINERAL ANALYSIS.....	43
3.2.1 Determination of Non-crystalline Secondary Minerals	
(Allophane and Ferrihydrite)	48
3.2.1.1 Acid Oxalate Extract.....	48
3.2.1.2 Sodium Pyrophosphate Extraction.....	48
3.3 SOIL CHEMICAL ANALYSIS.....	49
3.3.1 pH in Water.....	49
3.3.2 pH in 1M KCl.....	49
3.3.3 pH in 1M NaF Solution.....	49
3.3.4 Phosphate Retention.....	50
3.4 PHOSPHORUS FRACTIONATION.....	50
3.4.1 Sequential Extraction Procedure.....	50
3.4.2 Bicarbonate P Fractionation.....	53
3.4.2.1 Bicarbonate Pi.....	53
3.4.2.2 Bicarbonate Po	53
3.5 TOTAL CARBON AND NITROGEN DETERMINATION.....	54
3.6 OVEN DRY WEIGHT.....	54

CHAPTER 4

RESULTS AND DISCUSSION

4.1 SOIL MINERALS.....	55
4.1.1 Sand and Silt Fractions.....	55
4.1.2 Clay Fraction.....	58
4.1.3 Non-crystalline Secondary Soil Minerals.....	61
4.1.3.1 Hoskins Soils (107).....	62
4.1.3.2 Kapiura Soils (402).....	63
4.1.3.3 Bialla Soils (205 and 201)	63
4.1.3.4 Popondetta Soils (323).....	65
4.1.2.5 New Zealand Soils	65
4.2 SOIL CHEMICAL PROPERTIES.....	66
4.2.1 pH in Water and 1M KCl.....	66

4.2.2 pH in 1M NaF and Phosphate Retention.....	69
4.2.2.1 pH in 1M NaF.....	69
4.2.2.2 Phosphate Retention.....	70
4.2.2.2.1 Hoskins and Kapiura Trials 107 and 402.....	70
4.2.2.2.2 Bialla Trials 205 and 201.....	70
4.2.2.2.3 Popondetta Trial 323.....	71
4.2.3 Relationships; Phosphate Retention and Oxalate-Extractable Al and Fe.....	72
4.3 PHOSPHORUS FRACTIONATION.....	74
4.3.1 Introduction - P Fractions.....	74
4.3.1.1 Readily Available P Forms.....	74
4.3.1.2 Less Readily-Available P Forms.....	75
4.3.1.2.1 Organic P Forms.....	75
4.3.1.2.2 Inorganic P Forms.....	75
4.3.1.3 Unavailable P Forms.....	76
4.3.2 Hoskins Soils.....	76
4.3.3 Kapiura Soils.....	82
4.3.4 Bialla Soils (Trials 205 and 201).....	85
4.3.5 Popondetta Soils (Trial 323).....	87
4.3.6 New Zealand Soils.....	87
4.3.7 Comparison Amongst PNG Soils	88
4.3.8 Comparison Between PNG and New Zealand Soils.....	89
4.4 PHOSPHORUS ACCUMULATION FROM FERTILISER APPLICATIONS.....	89
4.5 TOTAL CARBON AND NITROGEN ANALYSIS.....	98
4.6 GENERAL DISCUSSION.....	102
4.6.1 Review of PNG Oil Palm Field Trials.....	102
4.6.2 Soil Characteristics (Chemical and Mineralogical) and Phosphorus Status.103	103
4.6.2.1 Mineralogical Properties and Phosphate Retention.....	103
4.6.2.1.1 Hoskins and Kapiura.....	103
4.6.2.1.2 Bialla.....	104
4.6.2.1.3 Popondetta.....	105
4.6.3 Comparison of NZ and PNG Soils.....	105

4.6.4 Fate of P Fertilisers in PNG Soils.....	106
4.6.5 Categorising Soils for P Responsiveness.....	107
 CHAPTER 5	
5.0 CONCLUSION AND IMPLICATIONS.....	109
REFERENCES.....	113
APPENDICES.....	128

LIST OF TABLES

Table 2.1 Total SiO ₂ % Composition of Volcanic Ash Materials.....	7
Table 2.2 Examples of Common Primary P Minerals.....	18
Table 2.3 PNG Field P Fertiliser Trial Results, January 1988 to December 1990.....	26
Table 2.4 Critical Nutrient Concentration in Oil palm Leaf (IRHO Trials).....	31
Table 2.5 Critical Nutrient Concentration in Oil palm Leaf (Malaysian Trials)	31
Table 2.6 Nutrient Removal, Immobilisation and Turnover in Adult Oil Palms (kg/ha/yr).....	34
Table 4.1 Sand and Silt Fractions Minerals in a Range of Volcanic Soils from PNG and NZ	57
Table 4.2 Clay Fraction Minerals.....	60
Table 4.3 Acid Oxalate Al, Fe and Si Extracts.....	64
Table 4.4 Soil pH's and Phosphate Retention at 0 - 30 cm Depths.....	68
Table 4.5 Correlation of Phosphate Retention with Al _o and Fe _o for all Soils and Depths.....	72
Table 4.6 Correlation of Al _o and Fe _o with Phosphate Retention at Individual Depths..	72
Table 4.7 Regression Equations for Relationship Between Phosphate Retention and Allophane and Ferrihydrite Contents Over All Soils and Depths.....	73

Table 4.8 P Fractions in mg/kg (% of Total Extractable P is shown in Parenthesis).....	78
Table 4.9 Average Organic P % for all Soil Depths for PNG Soils.....	88
Table 4.10 Total Soil P Contents of Fertilised and Nonfertilised Soils at Hoskins (107) and Kapiura (402)	92
Table 4.11 P Accumulation of P Fractions in Hoskins Soils at 0 - 10 cm Depth.....	94
Table 4.12 P Accumulation of P Fractions in Hoskins Soils at 10 - 20 cm Depth.....	95
Table 4.13 P Accumulation of P Fractions in Kapiura Soils at 0 - 10 cm Depth.....	95
Table 4.14 P Accumulation of P Fraction in Kapiura Soils at 20 - 30 cm Depth.....	96
Table 4.15 Total Soil C, N and C:N Ratios for PNG and NZ Soils.....	100

LIST OF FIGURES

Figure 3.1 Flow Sheet for Soil Mineralogical Analysis.....	44
Figure 3.2 Outline of Hedley's Sequential P Fractionation Procedure	51
Figure 4.1 P Fractionation Scheme.....	74
Figure 4.2 (a) Relative Proportions of P Fractions - P Fractions, Hoskins Soils Trial 107 HP.....	81
Figure 4.2 (b) Relative Proportions of P Fractions - Bicarbonate Extractable P, Hoskins Soils Trial 107 HP.....	81
Figure 4.3 (a) Relative Proportions of P Fractions - P Fractions, Kapiura Soils Trial 402 HP.....	84
Figure 4.3 (b) Relative Proportions of P Fractions - Bicarbonate Extractable P, Kapiura Soils Trial 402 HP.....	84
Figure 4.4 (a) Relative Proportions of P Fractions - P Fractions, Bialla Soils Trial 201	86
Figure 4.4 (b) Relative Proportions of P Fractions - Bicarbonate Extractable P, Bialla Soils Trial 201.....	86
Figure 4.5 Outline of P Fertilised Area per Oil Palm Tree.....	91