

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Characterisation of ERK Distribution and Activity in Rat Pheochromocytoma Cells

A thesis presented in partial fulfilment of the requirements for the degree of
Master of Science in Molecular Biology at Massey University

Matthew Edgar MacCormick
Institute of Molecular Biosciences
August 1998

ABSTRACT

Nerve growth factor (NGF) binds to the NGF receptor, TrkA, at the tips of nerve cell axons, sending a signal that prevents programmed cell death and causes survival, growth, and differentiation of the nerve cell. Both NGF and TrkA have been demonstrated to be retrogradely transported from axon tips to nerve cell bodies, however the mechanism of this transport, and its function, is strongly debated. Using a recently developed cell fractionation protocol in conjunction with *in vitro* reactions using an ATP regenerating system, our lab has isolated small vesicles containing NGF bound to activated TrkA. These vesicles may provide a vehicle for retrograde transport of the NGF signal and initiation of signal transduction in the cell body.

ERK1 is a serine/threonine kinase that is activated by NGF-activated TrkA. Prolonged ERK1 activity is characteristic of cells stimulated by NGF. The purpose of the experiments in this thesis was to characterise the intracellular distribution and activity of ERK1 before and after NGF stimulation, in rat pheochromocytoma (PC12) cells, which are a good model for nerve cells. We have found that ERK1 activity is redistributed between cell compartments after NGF stimulation of PC12 cells. ERK1 activity increased in sedimentable fractions that emerged from mechanically permeabilised cells after NGF treatment and *in vitro* reactions with ATP. Importantly, the results from glycerol velocity gradient experiments showed that ERK1 was not associated with membranes. Instead ERK1 was found in a rapidly sedimenting particle whose sedimentation was not affected by detergent solubilisation. These results suggest that ERK1 is recruited into a protein complex, after activation, which may be an important step in signal transduction. Formation of this complex is likely to be downstream of signalling vesicles containing NGF bound TrkA.

LIST OF ABBREVIATIONS

AMP	Adenosine monophosphate
APS	Ammonium persulphate
ATP	Adenosine triphosphate
BB	Bud buffer
BDNF	Brain-Derived Neurotrophic Factor
BSA	Bovine serum albumin
ChAT	Choline acetyltransferase
CNS	Central nervous system
COS	Transformed African Green Monkey kidney cell line
CRE	Cyclic AMP response element
CREB	Cyclic AMP response element binding protein
ddH ₂ O	Double-distilled water
DMSO	Dimethyl sulphoxide
DNA	Deoxyribonucleic acid
DTT	Dithiothreitol
ECL	Enhanced chemi-luminescence
EDTA	Ethylenediamine tetraacetic acid
EGFR	Epidermal growth factor receptor
EGTA	Ethylene glycol-bis(β -aminoethyl ether) N,N,N',N'-tetraacetic acid
ERK	Extracellular regulated kinase
FGF	Fibroblast growth factor
GAP	GTPase activating protein
GDP	Guanosine diphosphate
GST	Glutathione S-transferase
GTP	Guanosine triphosphate
Grb2	Growth factor receptor binding protein 2
IEG	Immediate early gene
IP.	Immunoprecipitation
kDa	Kilo Dalton
LRG	Late response gene
MAPK	Mitogen activated protein kinase

MEK	MAPK or ERK Kinase
MOPS	3-(N-Morpholino)propanesulphonic acid
mRNA	messenger ribonucleic acid
m-Sos	Mammalian Son-of-Sevenless
NGF	Nerve growth factor
NIH 3T3	Contact inhibited NIH Swiss mouse embryo cell line
NT	Neurotrophin
p75 ^{NTR}	p75 neurotrophin receptor
PBS	Phosphate buffered saline
PC12	Adrenal Pheochromocytoma rat cell line
PCD	Programmed cell death
PEE	PBS with EDTA and EGTA
PGB	PBS with glucose and BSA
PI-3 kinase	Phosphatidylinositol-3' kinase
PKC	Protein kinase C
PLC- γ 1	Phospholipase C gamma 1
PMSF	Phenylmethanesulphonyl fluoride
PNS	Peripheral nervous system
PtdIns	Phosphatidylinositol
SDS-PAGE	Sodium dodecyl sulphate polyacrylamide gel electrophoresis
SH	Src homology
SRF	Serum response factor
SY5Y	SK-N-SH-SY5Y human neuroblastoma cell line
TCA	Trichloroacetic acid
TBS	Tris buffer saline
TEMED	N,N,N',N' tetramethyl ethylene diamine
TH	Tyrosine hydroxylase
Tris	Tris(hydroxymethyl)aminoethane
Triton X-100	t-Octylphenoxypolyethoxyethanol
Trk	Tyrosine receptor kinase
Tween 20	Polyoxyethylenesorbitan monolaurate
Tween 40	Polyoxyethylenesorbitan monopalmitate
v/v	Volume/volume
w/v	Weight/volume

TABLE OF CONTENTS

ABSTRACT.....	i
LIST OF ABBREVIATIONS.....	ii
TABLE OF CONTENTS.....	iv
LIST OF FIGURES AND TABLES.....	vi
ACKNOWLEDGEMENTS.....	vii
CHAPTER 1: INTRODUCTION	1-1
1.1 OVERVIEW OF PROGRAMMED CELL DEATH.....	1-1
1.2 NGF IS IMPORTANT FOR NEURON SURVIVAL.....	1-2
1.3 NGF RECEPTORS.....	1-3
1.4 THE ROLE OF P75 ^{NTR}	1-4
1.5 <i>TrkA</i> GENE EXPRESSION AND REGULATION IN THE NERVOUS SYSTEM.....	1-6
1.6 PC12 CELLS.....	1-8
1.7 ACTIVATION OF <i>TrkA</i> – STIMULATION OF MULTIPLE DOWNSTREAM PATHWAYS	1-9
1.8 PI-3 KINASE AND AKT	1-10
1.9 PLC- γ 1.....	1-12
1.10 THE MAPK PATHWAY.....	1-13
1.10.1 SH2/SH3/PH Domains	1-13
1.10.2 Shc, Grb2, and Sos Bind First.....	1-14
1.10.3 Ras.....	1-16
1.10.4 Ras Association with Raf Leads to Raf Activation.....	1-17
1.10.5 Raf Activates MEK	1-18
1.10.6 MEK Activates ERK	1-19
1.10.7 Prolonged Activation of ERK Leads to Differentiation.....	1-20
1.11 GENE ACTIVATION BY NGF.....	1-21
1.12 RETROGRADE TRANSPORT OF NGF.....	1-24
1.13 THESIS OBJECTIVES.....	1-28
CHAPTER 2: MATERIALS AND METHODS	2-30
2.1 CHEMICALS AND SOLUTIONS.....	2-30
2.2 CELL CULTURE.....	2-30
2.3 CELL FRACTIONATION PROTOCOL.....	2-31
2.4 CELL CRACKING.....	2-33
2.5 SDS-PAGE.....	2-34
2.6 WESTERN BLOTTING WITH ERK.....	2-34

2.7 ECL METHOD	2-35
2.8 MYELIN BASIC PROTEIN IN-GEL KINASE ASSAY	2-36
2.9 IMMUNOPRECIPITATION	2-36
2.10 NON-RADIOACTIVE MAPK ASSAY	2-37
2.11 GLYCEROL GRADIENTS	2-38
2.12 STRIP AND REPROBE WITH SYNAPTOPHYSIN.....	2-38
2.13 NIH IMAGE.....	2-39
2.14 CALCULATIONS.....	2-39
CHAPTER 3: RESULTS	3-40
3.1 ERK DISTRIBUTION IS UNEQUAL BETWEEN CELL FRACTIONS.....	3-40
3.2 ERK ACTIVITY CAN BE MEASURED BY IN GEL KINASE ASSAYS	3-46
3.3 ERK ACTIVITY DOES NOT MATCH ERK DISTRIBUTION	3-50
3.4 ERK ACTIVITY IS ASSOCIATED WITH P2 AND P3 FRACTIONS.....	3-53
3.5 ERK IS FOUND IN HEAVY AND LIGHT P3 GRADIENT FRACTIONS	3-59
3.6 EVIDENCE FOR PHOSPHORYLATED ERK	3-63
CHAPTER 4: DISCUSSION	4-65
4.1 DISCUSSION AND CONCLUSIONS	4-65
4.2 FUTURE WORK	4-70
REFERENCES.....	72

LIST OF FIGURES AND TABLES

FIGURES

FIGURE 1: TYROSINE PHOSPHORYLATION SITES ON TRKA.....	1-9
FIGURE 2: MULTIPLE DOWNSTREAM PATHWAYS STIMULATED BY TRKA ACTIVATION....	1-11
FIGURE 3: MODELS FOR THE RETROGRADE TRANSPORT OF NGF ACTIVATED TRKA.....	1-26
FIGURE 4: OVERVIEW OF CELL FRACTIONATION PROTOCOL.....	2-33
FIGURE 5: CELL CRACKER	2-34
FIGURE 6: IDENTIFICATION OF ERK1 AND 2 IN PC12 FRACTIONS.....	3-42
FIGURE 7: AVERAGE VALUES FOR ERK DISTRIBUTION BETWEEN CELL FRACTIONS.....	3-43
FIGURE 8: IDENTIFICATION OF ERK1 AND 2 IN PC12 CELL FRACTIONS WITH <i>IN VITRO</i> REACTIONS.....	3-45
FIGURE 9: MBP IN-GEL KINASE ACTIVITY ASSAY WITH ERK IP	3-47
FIGURE 10: ERK TOTAL ACTIVITY USING MBP IN-GEL KINASE ASSAYS.....	3-49
FIGURE 11: AVERAGED VALUES FOR ERK ACTIVITY BETWEEN CELL FRACTIONS.....	3-51
FIGURE 12: ERK1 SPECIFIC ACTIVITY	3-53
FIGURE 13: MBP IN-GEL KINASE ACTIVITY ASSAY	3-54
FIGURE 14: NON-RADIOACTIVE MAPK ASSAYS.....	3-56,57
FIGURE 15: WESTERN BLOT WITH PHOSPHOMAPK	3-58
FIGURE 16: WESTERN BLOT OF GLYCEROL VELOCITY GRADIENTS.....	3-60
FIGURE 17: DETERGENT SOLUBILISED SAMPLES SUBJECTED TO VELOCITY GRADIENTS..	3-61
FIGURE 18: BLOTS REPROBED WITH ANTI-SYNAPTOPHYSIN.....	3-62

TABLES

TABLE 1: GENES ACTIVATED IN NGF STIMULATED CELLS	1-22
--	------

ACKNOWLEDGEMENTS

Firstly thanks go to my supervisor Dr. Mark Grimes who has given a lot of freedom and financial support for my project, so I can only hope I have produced something worthwhile! He has also patiently helped me with my less gifted areas, which has no doubt cost him a few cups of coffee. I would also like to thank my friend and fellow sufferer Scott Walker, who gave me a lot of help, especially when I was starting my project. I have also enjoyed working with, and received help and advice from Gretchen, Fleur, Treena, Louise, Lisa, and Annika. Special thanks also go to Treena and Louise, who gave me lots of advice in producing the finished product, and any mistakes in this thesis were no doubt inserted after they checked it! Lastly, and most importantly, I would like to thank Mum and Dad for their love and support.