

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

A STUDY OF SEED PRODUCTION
IN
'GRASSLANDS RUANUI' PERENNIAL RYEGRASS (LOLIUM PERENNE L.)
'GRASSLANDS KAHU' TIMOTHY (PHLEUM PRATENSE L.)
AND PRAIRIE GRASS (BROMUS UNIOLOIDES H.B.K.)

A Thesis Presented in Partial Fulfilment
of the Requirements for the Degree of
Doctor of Philosophy at
Massey University
New Zealand

by

Murray John Hill
November 1971

CONTENTS

	<u>Page</u>
<u>INTRODUCTION</u>	1
<u>STUDY OUTLINE</u>	3
<u>DESCRIPTION OF THE GRASSES STUDIED</u>	4
<u>MATERIALS AND METHODS</u>	11
<u>FIELD STUDY ON RYEGRASS, TIMOTHY AND PRAIRIE GRASS</u>	21
Tiller Production	21
Tiller Longevity and Survival	36
Floral Initiation and Development	43
Ear Emergence	57
Seedhead number	60
Fertile tiller production	61
Head contribution in emergence groups	64
Seedhead component analysis	69
Anthesis	77
Field experiment	80
Anthesis in timothy	82
Seed Development	86
Moisture content	90
Seed weight	91
Seed colour changes	94
Endosperm consistency	98
Germination	99
Seed Ripening, Harvesting and Yield	105
Seed yield	114
Seed quality	117
Assessment of optimum harvest time	120
Discussion of Field Study in Relation to Seedcrop Management	126
<u>ENVIRONMENTAL STUDY ON RYEGRASS AND PRAIRIE GRASS</u>	141
Effect of Environment on the Pattern of Daily Anthesis	147
Pollination, Fertilisation and Seed 'Set'.	167
Seed Development	179
Conclusion	188
<u>BIBLIOGRAPHY</u>	
<u>APPENDICES</u> (presented in separate volume)	

LIST OF TABLES

	<u>Page</u>
1. Fate of tillers formed each month as determined at time of seed harvest. (First year 1966)	Following 40
2. Fate of tillers formed each month as determined at time of seed harvest (second year 1967).	Following 40
3. Mean apex length and distance of internodal elongation at ear emergence (1966) (Extracted from Appendix 5).	54
4. Total average seedheads per row foot at harvest.	60
5. Percentage fertile tiller production.	61
6. Percentage contribution of seedheads in each emergence group.	65
7. Percentage of total seedhead population contributed by tillers of different ages.	67
8. Head length variation between treatments and emergence groups.	70
9. Spikelet number variation between heads in different treatments and emergence groups.	71
10. Floret number variation between heads in different treatments and emergence groups.	72
11. Culm length variation between heads in different treatments and emergence groups.	74
12. Progress of daily anthesis in selected timothy heads.	83
13. Seed numbers produced on timothy heads following self and cross pollination.	84
14. Summary of anthesis and seed development data in different regions of timothy heads.	84
15. Seed yields 1966 (means in lb/ac. at 15% seed moisture content).	115
16. Seed yields 1967 (means in lb/ac at 15% seed moisture content).	116
17. Results of analysis of treatment data for components of daily anthesis pattern in ryegrass (time of onset, time of peak, total florets open at peak anthesis).	153
18. Results of analysis of treatment data for components of daily anthesis pattern in ryegrass (number florets open at peak anthesis, length of time florets open, anthesis duration).	155
19. Results of analysis of treatment data for components of daily anthesis pattern in prairie grass (time of onset, time of peak, total florets open at peak anthesis).	157

	<u>Page</u>
20. Results of analysis of treatment data for components of daily anthesis pattern in prairie grass (number florets open at peak anthesis, length of time florets open, anthesis duration).	158
21. Mean time intervals required for completion of various stages of pollination, fertilisation and seed 'set' in ryegrass and prairie grass.	175
22. Statistical analysis of percentage of seeds formed at varying times from date of anthesis.	185

LIST OF PLATES

	<u>Page</u> Following
1. Experimental layout of field trial area	11
2. Prairie grass sample removed for tiller density counts.	28
3. Vegetative apex showing leaf primordia - perennial ryegrass.	48
4. Reproductive apex, double ridge stage - perennial ryegrass.	49
5. Sheltered treatment used in environmental study.	143
6. Shaded treatment used in environmental study.	144
7. Ovary prior to fertilisation - perennial ryegrass.	173
8. Pollen tube tip pushing between the two synergidae immediately prior to discharge of male nuclei - perennial ryegrass.	173
9. First division of egg mother cell - perennial ryegrass	173
10. Ryegrass seed seven days after pollination showing ovule collapse following ineffective fertilisation.	176
11. Developing prairie grass seed four days after pollination.	177
12. Ovule classification groups following floret dissection.	182
13. Ovule showing inverted T ovary shape (Group B)	183
14. Developing prairie grass seed 4 days after pollination.	187
15. Developing prairie grass seed 7 days after pollination.	187
16. Developing prairie grass seed 14 days after pollination.	187
17. Developing prairie grass seed 21 days after pollination.	187

LIST OF FIGURES

Following Page

1.	Field trial layout.	11
2.	Apparatus for artificially drying seed.	16
3.	Mean tiller densities per row foot, perennial ryegrass.	29
4.	Mean tiller densities per row foot, timothy.	30
5.	Mean tiller densities per row foot, prairie grass.	31
6.	Stages of reproductive development, perennial ryegrass.	46
7.	Stages of reproductive development, timothy.	46
8.	Stages of reproductive development, prairie grass.	46
9.	Percentage of reproductive tillers of different months of origin following apical dissection.	51
10.	Mean apex length and distance of internodal elongation at different stages of reproductive development (-G-N).	53
11.	Annual pattern of anthesis.	80
12.	Changes in seed moisture content.	90
13.	Changes in seed weight components.	91
14.	Seed weight differences between heads from different emergence groups.	92
15.	Percentage of heads in different colour groups during seed development.	95
16.	Percentage of seeds in different colour groups during development.	95
17.	Pigment changes during seed development.	96
18.	Germination capacity of seed harvested at different intervals after anthesis.	99
19.	Germination capacity of seeds from heads in different emergence groups.	102
20.	Changes in seed components and yield during development, perennial ryegrass.	117
21.	Changes in seed components and yield during development, timothy.	118
22.	Changes in seed components and yield during development, prairie grass.	119
23.	Seed drying curves, perennial ryegrass.	120
24.	Seed drying curves, timothy.	120
25.	Seed drying curves, prairie grass.	120

Following Page

26.	Trial layout for environmental study of anthesis and seed development.	142
27.	Daily anthesis pattern in relation to environmental conditions, perennial ryegrass.	152
28.	Daily anthesis pattern in relation to environmental conditions, prairie grass.	156
29.	Diagrammatic representation of pollination and the path of pollen tube development in perennial ryegrass.	172
30.	Percentage of seeds formed at different intervals after anthesis.	181

LIST OF APPENDICES (presented in separate volume)

1. Dates of grazing, nitrogen application and general management of field trial.
2. Mean tiller densities per row foot.
3. Stages of reproductive development - ryegrass and prairie grass.
4. Percentage of reproductive tillers observed following apical dissection.
5. Mean measurement of apex length and distance of internodal elongation.
6. Percentage distribution of heads produced in each emergence group according to month of tiller origin (1966, 1967).
7. Data on selected components of seedheads in emergence groups (1966, 1967).
8. Results of analysis of variance between means of treatment and emergence groups (1966, 1967).
9. Distribution of timothy heads in arbitrary head length categories (1966, 1967).
10. Percentage of seedheads with anthers exerted at each observation date (1966, 1967).
11. Anthesis and seed development data in different regions of timothy heads.
12. Technique for determination of seed chlorophyll and anthocyanin content.
13. Variation in seed moisture content during seed development.
14. Changes in individual weight components during seed development.
15. Mean weight of 1000 seeds developed from heads in different emergence groups.
16. Percentage of heads in different colour categories at successive harvest dates.
17. Percentage of seeds in different colour categories at successive harvest dates.
18. Changes in chlorophyll and anthocyanin content during seed development.
19. Data on onset of germination capacity - perennial ryegrass, timothy and prairie grass.
20. Germination capacity of seeds from heads in different emergence groups.

21. Seed yield - perennial ryegrass, timothy and prairie grass (1966).
22. Seed yield - perennial ryegrass, timothy and prairie grass (1967).
23. Relationship between seed yield and germination capacity at different harvest periods following anthesis depending on harvesting method (1966).
24. Relationship between seed yield and germination capacity at different harvest periods following anthesis depending on harvesting method (1967).
25. Daily anthesis data - perennial ryegrass (days 1 to 12).
26. Correlation matrix - perennial ryegrass.
27. Daily anthesis data - prairie grass (days 1 to 6).
28. Correlation matrix - prairie grass.
29. Average length of time individual florets remained open - prairie grass.
30. Percentage of seeds developed from florets open on different days and harvested at intervals up to 30 days after anthesis.
31. Percentage germination of seed developed from florets open on different days and harvested at intervals up to 30 days after anthesis.