

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

NITROGEN RELATIONSHIPS IN GRAZED AND CUT

GRASS-CLOVER SYSTEMS

A thesis presented in partial fulfilment for the degree of Doctor of Philosophy in Agronomy at Massey University.

Philip Roger Ball

1979

| TABLE OF CONTENTS | PAGE |
|--|------|
| <u>SECTION I</u> - GENERAL INTRODUCTION | 2 |
| 1 Productivity of New Zealand Pastures | 2 |
| 2 Concepts of Nitrogen Relationships Based on Research in New Zealand | 4 |
| 3 Recent Developments in Farming and Research | 7 |
| (a) Farming patterns | 7 |
| (b) Fertiliser nitrogen | 8 |
| (c) Research findings and concepts | 8 |
| 4 Experimental Objectives | 12 |
| <u>SECTION II</u> - DRY HERBAGE YIELDS AND SWARD CHARACTERISTICS | 13 |
| A INTRODUCTION | 13 |
| B MATERIALS AND METHODS | 17 |
| 1 Site and Climate | 17 |
| 2 Treatments | 18 |
| (a) Method of defoliation | 18 |
| (b) Nitrogen treatments | 19 |
| 3 Experimental Procedure | 20 |
| 4 Design, Statistical Procedure and Format of Results | 22 |
| C RESULTS AND DISCUSSION | 23 |
| 1 Total and Species ⁰ Herbage DM Yields | 23 |
| (a) Method of defoliation | 24 |

| | | |
|-----|---|----|
| (b) | Responses to fertiliser nitrogen | 25 |
| (c) | Efficiency of nitrogen responses | 28 |
| (d) | Performance of pure grass swards | 29 |
| (e) | Defoliation method X nitrogen treatment interactions | 30 |
| 2 | Plant Populations and Sward Structure | 31 |
| (a) | Method of defoliation | 31 |
| (b) | Fertiliser nitrogen treatments | 32 |
| (c) | Pure grass swards | 33 |
| (d) | Sward characteristics at March 1974 | 33 |
| D | GENERAL DISCUSSION | 34 |
| 1 | Method of Defoliation | 35 |
| 2 | Fertiliser Nitrogen Responses | 39 |
| 3 | Pure Grass Swards | 45 |
| E | CONCLUSIONS | 46 |

SECTION III - HERBAGE NITROGEN, AND MINERAL

| | | |
|-----|--|----|
| | CHARACTERISTICS OF HERBAGE AND SOIL | 48 |
| A | INTRODUCTION | 48 |
| B | MATERIALS AND METHODS | 54 |
| 1 | Site and Climate | 54 |
| 2 | Treatments | 55 |
| (a) | Method of defoliation | 55 |
| (b) | Nitrogen treatments | 56 |
| 3 | Experimental Procedure | 56 |
| 4 | Design, Statistical Procedure and Format of Results | 59 |

| | | |
|---|---|----|
| C | RESULTS | 60 |
| 1 | Herbage Total and Nitrate Nitrogen Contents | 60 |
| 2 | Herbage Nitrogen Yields | 62 |
| | (a) Defoliation method X nitrogen treatment interactions | 63 |
| | (b) Method of defoliation | 63 |
| | (c) Responses to fertiliser nitrogen | 64 |
| | (d) Pure grass swards | 66 |
| 3 | Mineral Composition of Herbage | 67 |
| | (a) Herbage phosphorus content | 67 |
| | (b) Herbage potassium content | 68 |
| | (c) Herbage magnesium and calcium contents | 69 |
| 4 | Soil 'Quicktest' Measurements | 70 |
| 5 | Apparent DM Intake and Digestibility of Herbage | 71 |
| D | DISCUSSION | 71 |
| 1 | Nitrogen Yields, Total and Nitrate Levels in Herbage | 71 |
| | (a) Defoliation treatments | 73 |
| | (b) Fertiliser nitrogen | 75 |
| | (c) Pure grass swards | 77 |
| | (d) Seasonal influences | 80 |
| | (e) Total nitrogen levels in herbage | 82 |
| 2 | Mineral Characteristics of Herbage and Soil | 83 |
| | (a) Phosphorus, potassium, magnesium and calcium levels in herbage | 84 |
| | (b) Soil 'Quicktest' values | 86 |

| | | |
|---|--|---------|
| (c) | Assessment of basal fertiliser policy | 87 |
| 3 | A Brief Analysis of Fertiliser Nitrogen Responses, and Their Possible Modification by Potassium Deficiency | 90 |
| 4 | Herbage Intake and Digestibility | 96 |
| E | CONCLUSIONS | 99 |
| <u>SECTION IV - SOIL NITROGEN AND NITROGEN BALANCES</u> | | 101 |
| A | INTRODUCTION | 101 |
| B | MATERIALS AND METHODS | 103 |
| 1 | Site and Climate | 103 |
| 2 | Treatments | 106 |
| (a) | Method of defoliation | 106 |
| (b) | Nitrogen treatments | 106 |
| 3 | Experimental Procedure | 107 |
| (a) | Field sampling procedure | 107 |
| (b) | Analytical procedure | 110 |
| 4 | Design and Statistical Procedure | 112 |
| C | RESULTS AND DISCUSSION | 112 |
| 1 | Soil Total Nitrogen | 112 |
| (a) | Preliminary error studies | 113 |
| (b) | Sampling schedule | 118 |
| (c) | Changes over time | 119 |
| (d) | Treatment effects | 122 |
| (e) | Conclusions | 124 |
| 2 | Nitrogen Fixation in Grazed Swards | 125 |

| | | |
|-----|--|-----|
| (a) | Results | 125 |
| (b) | Discussion | 127 |
| (c) | Conclusions | 131 |
| 3 | Mineral Nitrogen Pools | 131 |
| (a) | Results | 131 |
| (b) | Discussion | 132 |
| (c) | Conclusions | 135 |
| 4 | Nitrogen Balances in Grazed and Cut Treatments | 136 |
| (a) | Results | 136 |
| (b) | Discussion | 137 |
| (c) | Conclusions | 143 |
| 5 | Nitrogen Mineralisation as Indicated by Herbage Nitrogen Yields from a Cut, Cloverless Sward | 143 |
| D | GENERAL DISCUSSION | 144 |
| 1 | Changes in Soil Total Nitrogen Pools | 144 |
| 2 | Influence of Grazing and Cutting on Changes in Soil Total Nitrogen | 147 |
| 3 | Influence of Nitrogen Treatments on Changes in Soil Total Nitrogen | 149 |
| 4 | Symbiotic Fixation and Fixation Efficiency in Grazed and Cut Swards | 150 |
| 5 | Nitrogen Losses from Grazed Systems | 151 |
| E | CONCLUSIONS | 152 |

| | |
|---|-----|
| <u>SECTION V - PERSPECTIVE COMMENTS AND MAJOR</u> | |
| CONCLUSIONS | 154 |
| A PERSPECTIVE COMMENTS | 154 |
| 1 Soil Organic Matter Quality and Level | 157 |
| 2 Grazing <u>vs</u> Cutting | 162 |
| 3 Mineralisation of Soil Nitrogen | 165 |
| 4 Nitrogen Fixation by White Clover, and Fixation Efficiency | 168 |
| 5 Nitrogen Transfer from Clover to Grasses | 178 |
| 6 Nutrient 'Recycling' by Grazing Animals | 183 |
| 7 An Alternative Hypothesis for the Role of Grazing Animals | 188 |
| 8 Fertiliser Nitrogen | 190 |
| B MAJOR CONCLUSIONS | 192 |
| | |
| ACKNOWLEDGEMENTS | 198 |
| | |
| ABSTRACT | 199 |
| | |
| REFERENCES | 202 |

LIST OF TABLES

PAGE

SECTION II

| | |
|--|-----|
| TABLE 1 - Climatic data (from Grasslands Division's recording station) and soil moisture levels over the experimental period | 18a |
| TABLE 2 - Annual and seasonal dry herbage yields (t DM/ha); adjusted means for defoliation treatments | 24a |
| TABLE 3 - Annual and seasonal dry herbage yields (t DM/ha); adjusted means for nitrogen treatments | 24b |
| TABLE 4 - Average total dry herbage increments by grass-clover swards, and efficiencies of response, for seasons and year | 28a |
| TABLE 5 - Plant populations in swards (growing points/dm ²); adjusted means for main treatment effects | 31a |
| TABLE 6 - Structure of swards; adjusted means for main treatment effects | 31c |
| TABLE 7 - Average annual dry herbage yields and clover contents of grazed and mown swards, with and without animal return, in previous studies at Palmerston North | 34a |

SECTION III

| | |
|--|-----|
| TABLE 8 - Herbage nitrogen contents (g N/100 g dry herbage); adjusted means for main treatment effects | 60a |
|--|-----|

| | |
|--|-----|
| TABLE 9 - Herbage nitrate levels (ppm NO ₃ -N); adjusted means for main treatment effects | 60c |
| TABLE 10 - Annual herbage nitrogen yields (kg N/ha); adjusted means for main treatment effects | 62a |
| TABLE 11 - Seasonal herbage nitrogen yields (kg N/ha); adjusted means for main treatment effects | 62b |
| TABLE 12 - Herbage phosphorus contents (g P/100 g dry herbage); adjusted means for main treatment effects | 67a |
| TABLE 13 - Herbage potassium contents (g K/100 g dry herbage); adjusted means for main treatment effects | 68a |
| TABLE 14 - Herbage magnesium contents (g Mg/100 g dry herbage); adjusted means for main treatment effects | 69a |
| TABLE 15 - Herbage calcium contents (g Ca/100 g dry herbage); adjusted means for main treatment effects | 69c |
| TABLE 16 - Soil 'Quicktest' values; means for treatments NO and N448, sampled on commencement and conclusion of the experiment | 70a |
| TABLE 17 - Apparent dry herbage intakes (kg DM/sheep-day) over the three years of the experiment, and dry matter digestibility (%) estimates for mixed herbage at 5 harvests over mid spring-summer in the final year | 71a |

SECTION IV

- TABLE 18 - Variability in measured values for soil total nitrogen content:
 (a) among repeated analyses of reference samples; (b) among individual cores drawn from within small paddocks and small plots 114a
- TABLE 19 - Changes in total soil nitrogen pools (\bar{d} kg N/ha) throughout the sampled profile, and at various sampling depths, between successive measurements 122a
- TABLE 20 - Seasonal and annual estimates for $N_2(C_2H_2)$ fixation (kg N/ha), and fixation efficiency (kg N fixed/t clover DM measured) in grazed swards 125b
- TABLE 21 - Nitrogen balances in grazed and cut systems; all data are annual averages (kg N/ha) 136a

| LIST OF FIGURES | PAGE |
|--|------|
| <u>SECTION II</u> | |
| FIGURE 1 - General views across the experimental area | 18b |
| FIGURE 2 - Herbage DM responses to fertiliser nitrogen by grass-clover swards; averages of annual and seasonal yields | 25a |
| FIGURE 3 - Comparison of grazed and cut swards under drought conditions; mid summer 1974 | 33a |
| FIGURE 4 - Annual mean clover and grass DM yields and sward characteristics measured in autumn, expressed as the ratios Grazed:Cut and N448:NO, throughout the experiment | 37a |
| <u>SECTION III</u> | |
| FIGURE 5 - Herbage nitrogen yield responses to fertiliser nitrogen by grass-clover swards; averages for annual and seasonal yields | 64a |
| FIGURE 6 - Annual mean herbage nitrogen yields and weighted herbage nitrogen contents, expressed as the ratios Grazed:Cut and N448:NO (or NO:Grass), throughout the experiment | 73a |
| FIGURE 7 - Seasonal averages for herbage nitrogen yields and herbage nitrogen contents, expressed as the ratios Grazed:Cut and N448:NO | 80a |

FIGURE 8 - Pastures at grazing height; early
spring 1974 95a

SECTION IV

FIGURE 9 - General trends in soil total
nitrogen pools and C/N ratio over
the course of the experiment 119a

FIGURE 10 - General trends in soil total
nitrogen content and bulk density
over the course of the experiment 119b

FIGURE 11 - Daily $N_2(C_2H_2)$ fixation rates
(kg N/ha) in grazed treatments
during the final year 125a

FIGURE 12 - Soil mineral nitrogen pools in
the sampled profile (kg N/ha-
45.7 cm) below grazed pastures
over the final 15 months; also
indicated, the water balance
between consecutive samplings 131a