

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

A STUDY OF KEMP VARIATION IN THE FLEECES  
OF CHEVIOT EWES

D.F.G. ORWIN

October 1959

A thesis presented at Massey Agricultural College  
in partial fulfilment of the requirements for the  
Degree of Master of Agricultural Science in the  
University of New Zealand.

## ACKNOWLEDGMENTS

The author wishes to make grateful acknowledgment to his supervisor, Professor A.L. Rae, for his continued interest, assistance and guidance in this project.

For assistance in various problems, appreciation is expressed to members of the Sheep Husbandry Department, particularly Dr. F.R.M. Cockrem and Mr. G.A. Wickham.

The author is indebted to other staff members and students who helped in this project and especially to: Mr. T.C. Crotty and the farm staff at Tuapaka, Mr. J.P. Kerr, Mr. H. Nevill and Miss H. Riseborough for assistance when sampling; Miss M.G. Campbell and her staff for co-operation in the use of library facilities, and Dr. R.M. Dolby, Mr. D.C. Collis, Mr. G. Reece and Miss M.E. Burton for help in photographic work.

Especial acknowledgment is due to Mrs. Puckey for typing this thesis.

## TABLE OF CONTENTS

	Page
I INTRODUCTION . . . . .	1
II REVIEW OF LITERATURE	
A. General	
1. Definition of kemp . . . . .	3
2. The morphology of kemps. . . . .	5
3. The importance of kemps. . . . .	6
B. The Methods of Measurement of Kempiness. . . . .	9
C. Physiological Aspects of Kemp Development, Growth and Shedding	
1. The development of kempiness in the fleece	12
2. The shedding of kemps . . . . .	24
D. Variation in Kempiness	
1. Regional variation . . . . .	34
2. Variation between sheep. . . . .	35
3. Breed variation . . . . .	36
4. Seasonal variation . . . . .	38
E. The Inheritance of Kempiness . . . . .	43
III MATERIALS AND METHODS	
A. Experimental Design . . . . .	45
B. Collection of Data . . . . .	45
C. Treatment and Analysis of Samples. . . . .	48
IV RESULTS	
A. Methods of Measurement	
1. Selection of measures . . . . .	52

	Page
2. Some errors involved in obtaining data for the number of kemps per unit area	62
3. The recut of previously sampled areas	68
4. The bilateral symmetry of kemp density in position A . . . . .	74
B. The Variation of Kempiness over the Body of the Cheviot Ewe	
1. The variation of kempiness within sampled areas . . . . .	75
2. The variation of kempiness in the areas comprising position A. . . . .	80
3. The variation in kempiness between positions A, B, C and D . . . . .	85
C. The Relationship between Positions A, B, C and D	
1. The relationship between total kempiness in the fleece and positions A and D . .	90
2. The relationship between position A and positions B, C and D . . . . .	100
3. The relationship of position D with positions A, B and C . . . . .	105
D. The Variation of Kempiness between Individual Sheep . . . . .	111
E. The Variation of Kempiness between Age Groups	116
F. The Variation of Kempiness at Different Sampling Times . . . . .	122
G. The Relationship between Percentage Medullation and Kempiness. . . . .	137
V DISCUSSION . . . . .	140
VI SUMMARY AND CONCLUSIONS . . . . .	163
VII BIBLIOGRAPHY . . . . .	166
APPENDIX . . . . .	170

## LIST OF TABLES

Table		Page
1	Normality test of logarithm transformed data	55
2	Correlation coefficients of total numbers of kempes per unit area with percentage weight, total numbers of kempes per g of wool and weight of kempes per unit area	57
3	Bartlett's test for homogeneity of variance of transformed data	59
4	The differences in sampled areas calculated as for a rectangle and a non-rectangular figure for 10 randomly selected Cheviot ewes	63
5	Analysis of variances of areas calculated from two sets of measurements taken at different periods of time	65
6	Area and density changes in recuts of previously sampled areas	67
7	Analysis of variance of data from position A samples taken from both sides of the body of 17 three-year-old Cheviot ewes	73
8	Logarithm means of total numbers of kempes per square centimetre of samples from position A on both sides of body	73
9	Analyses of variance of numbers of kempes in repeat subsamples from three-year-old Cheviot ewes (position A)	76
10	Analyses of variance of numbers of kempes in repeat subsamples from three-year-old Cheviot ewes (positions A, B, C, D)	78
11	Analysis of variance testing in 12 sheep of each age group the variation in total kempes per sq. cm. in the four areas of position A	79
12	The log means and standard errors of total kempes per square centimetre for each area in position A over all sampling times and for each age group	81
13	Multiple range test for differences in areas in position A	82

Table		Page
14	Analysis of variance of data from positions A, B, C and D sampled in 10 two-year-old Cheviot ewes	83
15	Log means and standard errors of positions A, B, C and D over all age groups and all sampling times	84
16	Orthogonal comparisons of between positions variation (Table 15)	87
17	The correlation and regression coefficients of total kempiness on kempiness in positions A and D at each sampling time for 10 two-year-old and five three-year-old Cheviot ewes	92
18	The test for differences between sampling times in the regression coefficients of total kempiness on kempiness in positions A or D at different sampling times for different age groups	95
19	The correlation and regression coefficients of kempiness in positions B, C and D on kempiness in position A at each sampling time for 10 two-year-old Cheviot ewes	98
20	The test for differences between sampling times in the regression coefficients of kempiness in positions B, C and D on kempiness in position A for 10 two-year-old Cheviot ewes	99
21	The correlation and regression coefficients of kempiness in positions B, C and D on kempiness in position A at each sampling time for five three-year-old Cheviot ewes	101
22	The test for differences between sampling times in the regression coefficients of kempiness in positions B, C and D on kempiness in position A for five three-year-old Cheviot ewes	102
23	The correlation and regression coefficients of kempiness in positions A, B and C on kempiness in position D at each sampling time for 10 two-year-old Cheviot ewes	104
24	The test for differences between sampling times in the regression coefficients of kempiness on positions A, B and C on kempiness in position D for ten two-year-old Cheviot ewes	106

Table		Page
25	The correlation and regression coefficients of kempiness in positions A, B and C on kempiness in position D at each sampling time for five three-year-old Cheviot ewes	107
26	The test for differences between sampling times in the regression coefficients of kempiness in positions A, B and C on kempiness in position D for five three-year-old Cheviot ewes	109
27	The analysis of variance of data on total kemps per sq. cm. from position A of all sheep	112
28	The analysis of variance of data on shed kemps per sq. cm. from position A of all sheep	113
29	The variances of data for position A for each age group at each sampling time	114
30	The logarithm means, the antilogarithms of the means and the standard errors of the logarithm means for data from position A in each age group	117
31	Multiple range test: Testing the differences in mean kempiness of four age groups at the 5% level	118
32	The logarithm means, the antilogarithms of the means and the standard errors of the logarithm means for data from position A in each age group	119
33	Multiple range test: Testing the differences in mean kempiness of four age groups at the 5% level	120
34	The regressions and standard errors of estimate of weight of kemps per sq. cm. on total kemps per sq. cm. for position A at all sampling times for each age group.	123
35	The test for differences between sampling times of the regression lines in Table 34 for each age group	124
36	The correlations of percentage medullation and number of kemps per sq. cm. over all sampling times and all age groups	138

## LIST OF FIGURES

Figure		Page
1	Kemps from the fleeces of Cheviot ewes. . . . .	7
2	The positions sampled on the Cheviot ewes in this experiment . . . . .	47
3	The frequency distributions of untransformed data from position A . . . . .	53
4	The frequency distributions of transformed data from position A . . . . .	54
5	The log means of kempiness in positions A, B, C and D with time for two and three-year-old ewes	86
6	The log means of kempiness in positions A, B, C and D with time for each of two age groups. .	89
7	The regressions of total kempiness on kempiness in position A for each sampling time . . . . .	93
8	The regressions of total kempiness on kempiness in position D for each sampling time . . . . .	94
9	The regressions of weight of total kemps/sq. cm. on total numbers of kemps/sq.cm. for each sampling time . . . . .	126
10	The regressions of weight of total kemps/sq. cm. on total numbers of kemps/sq. cm. for each sampling time . . . . .	127
11	The change in log means of total kemps/sq. cm. and shed kemps/sq. cm. with time for each age group. . . . .	129
12	The change in the total number of kemps/sq. cm. in position A with time for individual two-year- old sheep. . . . .	131
13	The change in the total number of kemps/sq. cm. in position A with time for individual three- year-old sheep. . . . .	132
14	The change in the total number of kemps/sq. cm. in position A with time for individual four- year-old sheep. . . . .	133
15	The change in the total number of kemps/sq. cm. in position A with time for individual five-year- old sheep. . . . .	134