

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

EFFECTS OF GRAZING ON THE
GROWTH AND DEVELOPMENT OF
WHITE CLOVER (Trifolium repens L.)

A thesis
presented in partial fulfilment of the requirements
for the degree
of
MASTER OF AGRICULTURAL SCIENCE
at
MASSEY UNIVERSITY

William Blamires Pascoe

1973

ACKNOWLEDGEMENTS

I wish to acknowledge the constant, unrestricted help, encouragement and stimulus that I have received from my supervisor, Professor B.R. Watkin.

The time and help, liberally given by Professor R. Townsley in the statistical analysis of the experiment was greatly appreciated.

For help in the field, I am grateful to Mr. R. Battersby and Mr. M. Thompson

I am indebted also to Mrs. Pauline Martis for her understanding and good service in the typing of this thesis.

For her forbearance and encouragement, I am indebted to my wife.

Finally, in sincerity, I wish to acknowledge the encouragement and motivation that I have received from my spiritual Father, God, during the course of this thesis.

CONTENTS

	<u>Page</u>
CHAPTER 1. <u>Review of Literature</u>	
Introduction	1
1.1. The White Clover plant and its components	1
1.2. White Clover cultivars and experimentation	2
1.3. The growth and development of White Clover	2
1.4. Petiole and leaf growth	6
1.5. Root growth	11
1.6. The influence of management on the growth and development of white clover	13
Conclusions	16
CHAPTER 2. <u>Experimental procedures</u>	
Introduction	18
2.A. Experimental site	18
2.B. Experimental treatments	18
2.C. Experimental layout	19
2.D. Experimental procedures and techniques	19
2.E. The split plot experiment	22
2.F. Statistical methods	23
CHAPTER 3. <u>Results</u>	
Introduction	31
3.1. Meteorological data	31
3.2. Herbage D.M. yield measurements	33
3.3. Botanical composition analysis	42
3.4. Weights of clover stolons, petioles and leaves	48
3.5. Length of stolon, number of terminal buds and new stolons	59
3.6. Petiole lengths	69
3.7. Leaf area measurements	69
3.8. Individual stolon measurements	83
3.9. Individual stolon lengths	85
3.10. Clover plant tracings	86
3.11. Root measurements	91
3.12. Vibrarecorders	92
3.13. Soil moistures and soil temperatures	92

	<u>Page</u>
CHAPTER 4. <u>Discussion</u>	
Introduction	
A. Discussion of experimental techniques	96
B. Discussion of results	10
C. General discussion	11
 SUMMARY	 12
 REFERENCES	 12
 <u>APPENDICES</u>	
Appendix 1 Meteorological data for Palmerston North	13
Appendix 2 Treatment interactions	13
Appendix 3 Comparison of core sampling and individual	13
Appendix 4 Explanation of statistical methods	13

LIST OF TABLES

<u>TABLE</u>		<u>Page</u>
<u>CHAPTER 1.</u>		
1.1.	Percentage of roots as measured by dry weight in 10 cm soil layers under several swards.	11
<u>CHAPTER 2.</u>		
2.1.	Experimental treatments.	
<u>CHAPTER 3.</u>		
3.2.1.	Total production for the grazing pressure treatments.	33
3.2.2.	Total production for the 3, 6 and 9 day treatments.	33
3.2.3.	Increase in d.m. in each regrowth period for the grazing pressures.	37
3.2.4.	Increases in d.m. over each regrowth period for the 3, 6 and 9 day treatments.	40
3.2.5.	Decreases in d.m. at each grazing for the grazing pressure treatments.	40
3.2.6.	The decreases in d.m. at each grazing for the 3, 6 and 9 day treatments.	41
3.3.1.	Comparisons between botanical analyses.	45
3.4.1.	The increases in fraction weights for the grazing pressure treatments for the first three regrowth periods.	51
3.4.2.	Increase in fraction weights in the first, second and third regrowth periods for the 3, 6 and 9 day grazing periods.	55
3.4.3.	The decreases in fraction weights at each grazing for the grazing pressure treatments.	56
3.4.4.	The decreases in fraction weights at each grazing for the 3, 6 and 9 day grazing periods.	58
3.5.1.	The increases in the parameters in the first, second and third regrowth periods for the grazing pressures.	62
3.5.2.	The increases in the parameters for the 3, 6 and 9 day grazing periods.	66
3.5.3.	The decreases in parameters at each grazing.	67
3.5.4.	The decreases in parameters at each grazing for the 3, 6 and 9 day grazing periods.	68
3.7.1.	The number of leaves in each size group to equal 1 LAI per 62.2 cm ² .	72
3.7.2.	The increases in leaf area during the first, second and third regrowth periods for the grazing pressure treatments.	75
3.7.3.	The increases in total leaf area of developing, mature and senescing leaf for the grazing periods.	80

	<u>Pa</u>
3.7.4. The decreases in leaf area at each grazing.	8
3.7.5. The decreases in leaf area for the grazing periods.	8
3.8.1. The levels of certain parameters measured on the stolons cited in the quadrats for the high pressure 'grazed' and low pressure treatments after each grazing.	8
3.8.2. Certain parameters measured after the third grazing for the high pressure 'grazed' and 'ungrazed' treatments.	8
3.9.1. The mean stolon lengths after grazing for the highpressure 'grazed', low pressure and high pressure 'ungrazed' treatments.	8
3.11.1. The weights of topgrowth and roots down to 76 cm.	9

CHAPTER 4.

4.4.1. Some clover production parameters for the high pressure 'grazed' and low pressure treatments.	1
4.4.2. Some clover productionparameters of the 3, 6 and 9 day grazing periods for the high pressure 'grazed' treatment.	1
4.4.3. Some clover production of the 3, 6 and 9 day grazing periods for the low pressure treatment.	1

LIST OF FIGURES

<u>FIGURE</u>	<u>Page</u>
<u>CHAPTER 2.</u>	
2.1. Experimental layout.	20
2.2. The four grazings.	19
2.3. Plot sampling.	21
<u>CHAPTER 3.</u>	
3.1.1. Daily rainfall over the experimental period.	32
3.2.1. The effects on the regrowth of white clover swards of the high pressure 'grazed' and low pressure treatments.	35
3.2.2. The effects on the regrowth of white clover swards of the high pressure 'grazed' and high pressure 'ungrazed' treatments.	36
3.2.3. The effects of the 3, 6 and 9 day grazing periods on the regrowth of white clover swards. (Mean of the high pressure 'grazed' and low pressure treatments).	38
3.2.4. The effects of the 3, 6 and 9 day grazing periods on the regrowth of white clover swards. (Mean of the high pressure 'grazed' and high pressure 'ungrazed' treatments).	39
3.3.1. The effects of the high pressure 'grazed' and low pressure treatments on botanical composition.	43
3.3.2. The effects of the high pressure 'grazed' and high pressure 'ungrazed' treatments on botanical composition.	44
3.3.3. The effects of the 3, 6 and 9 day grazing periods on botanical composition. (Mean of the high pressure 'grazed' and low pressure treatments).	46
3.3.4. The effects of the 3, 6 and 9 day grazing periods on botanical composition. (Mean of the high pressure 'grazed' and high pressure 'Ungrazed' treatments).	47
3.4.1. The effects on the weight of stolons, petioles and leaves of the high pressure 'grazed' and low pressure treatments.	49
3.4.2. The effects on the weight of stolons, petioles and leaves of the high pressure 'grazed' and high pressure 'ungrazed' treatments.	50
3.4.3. The effects on the weight of stolons, petioles and leaves of the 3, 6 and 9 day grazing periods. (Mean of the high pressure 'grazed' and low pressure treatments).	53
3.4.4. The effects on the weight of stolons, petioles and leaves of the 3, 6 and 9 day grazing periods. (Mean of the high pressure 'grazed' and high pressure 'ungrazed' treatments).	54
3.5.1. The effects on the length of stolon, numbers of terminal buds and new stolons of the high pressure 'grazed' and low pressure treatments.	60
3.5.2. The effects on the length of stolon, numbers of terminal buds and new stolons of the high pressure 'grazed' and high pressure 'ungrazed' treatments.	61

	<u>Page</u>
3.5.3. The effects on the length of stolon, numbers of terminal buds and new stolons of the 3, 6 and 9 day grazing periods. (Mean of the high pressure 'grazed' and low pressure treatments).	64
3.5.4. The effects on the length of stolon, numbers of terminal buds and new stolons of the 3, 6 and 9 day grazing periods. (Mean of the high pressure 'grazed' and high pressure 'ungrazed' treatments).	65
3.6.1. The effects on mature petiole lengths of the high pressure 'grazed' and low pressure treatments.	70
3.7.1. The effects on the composition of leaf area of the high pressure 'grazed' and low pressure treatments.	71
3.7.2. The effects on the distribution of mature leaf area into leaf size ranges of the high pressure 'grazed' and low pressure treatments.	73
3.7.3. The effects on the composition of leaf area of the high pressure 'grazed' and high pressure 'ungrazed' treatments.	74
3.7.4. The effects on the composition of leaf area of the 3, 6 and 9 day grazing periods. (Mean of the high pressure 'grazed' and low pressure treatments).	76
3.7.5. The effects on the distribution of mature leaf area into leaf size ranges of the 3, 6 and 9 day grazing periods. (Mean of the high pressure 'grazed' and low pressure treatments).	78
3.7.6. The effects on the composition of leaf area of the 3, 6 and 9 day grazing periods. (Mean of the high pressure 'grazed' and high pressure 'ungrazed' treatments).	79
3.10.1. Clover plant tracing.	88
3.10.2. Clover plant tracing.	89
3.10.3. Clover plant tracing.	90
3.13.1. The levels of soil moisture in the top 7.6 cm of soil during the experiment for the high pressure 'grazed' and low pressure treatments.	93
3.13.2. Soil temperatures at 0, 7.6 and 30.5 cm for three different clover canopies.	94

LIST OF PLATES

<u>PLATE</u>		<u>Page</u>
1	This shows a low pressure 3 day plot immediately after the first grazing.	26
2	This shows a high pressure 3 day plot immediately after the first grazing.	26
3	A close picture of a high pressure 3 day plot immediately after the first grazing.	27
4	A close up picture of a high pressure 'grazed' plot immediately after the second grazing.	27
5	A low pressure 3 day plot immediately after the second grazing.	28
6	During regrowth the denuded plants on the high pressure 'grazed' treatment grew again from plant centres.	28
7	54 days of regrowth for a high pressure 'ungrazed' 9 day plot.	29
8	Patch grazing was assessed using a quadrat.	29
9	The feather-marked white clover plants were plainly visible.	30
10	Soil profiles for root sampling.	30