

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**IDENTIFICATION OF DOTHISTROMIN
BIOSYNTHETIC PATHWAY GENES**

A thesis presented in partial fulfilment of
the requirements for the degree of
Masters of Science in Molecular Genetics

at

Massey University, Palmerston North
New Zealand

Carmel Jane Gillman

1996

ABSTRACT

Dothistromin is a polyketide-derived toxic secondary metabolite produced by the filamentous fungus *Dothistroma pini* which causes the disease *Dothistroma* needle blight in *Pinus radiata*. Dothistromin is considered to be an important component in the disease process, although its exact function is yet to be identified. By isolating and identifying genes involved in dothistromin biosynthesis, and subsequently obtaining mutants blocked or altered in the synthesis of dothistromin, the role of this toxin in pathogenicity will be able to be assessed. Dothistromin is structurally related to the mycotoxins, aflatoxin (AF) from *Aspergillus parasiticus* and *A. flavus*, and sterigmatocystin (ST) from *A. nidulans*. Three intermediates in the ST and AF biosynthetic pathways (averantin, averufin, and versicolorin B) are thought to also be intermediates dothistromin biosynthesis. Due to these similarities, cloned AF pathway genes were used as heterologous probes in Southern hybridisation analysis to provide a direct method for identifying dothistromin biosynthetic genes.

A fragment of the *A. parasiticus nor-1* gene, encoding a reductase involved in the conversion of norsolorinic acid (NA) to averantin (AVN) in the AF biosynthetic pathway, was used as a probe to detect a region of sequence similarity to *D. pini* genomic DNA. A *D. pini* genomic library was then constructed and screened, resulting in clone λ CGN2. However, Southern hybridisation analysis suggested that this clone did not contain a homologue of the *nor-1* gene from *A. parasiticus*.

A fragment of the *Aspergillus parasiticus ver-1* gene, encoding a reductase involved in the conversion of versicolorin A (VA) to ST in the AF biosynthetic pathway, was also used as a probe to detect a region of sequence similarity to *D. pini* genomic DNA. The *D. pini* genomic library was then screened. Two clones, λ CGV1 and λ CGV2, were isolated and Southern hybridisation analysis confirmed that these clones contained sequences hybridising to the *A. parasiticus ver-1* gene fragment. Fragments of these clones which hybridised were then sequenced and compared to the GenBank database. The amino acid coding sequence of a 0.8 kb *SalI* region from clone λ CGV1 exhibited a high degree of similarity with the *A. nidulans verA* and *A. parasiticus ver-1* genes, involved in the ST and AF biosynthetic pathways, and the *Magnaporthe grisea ThnR*, and *Colletotrichum lagenarium Thr1* genes, involved in melanin biosynthesis. This data suggested a *ver-1* homologue is present in the *D. pini* genome. Limited sequence analysis of a 2.1 kb region from clone λ CGV2 suggested that a second independent copy of a *ver-1*-like gene may also be present in the genome.

ACKNOWLEDGEMENTS

I feel very overwhelmed and honoured when I think of all the people I would like to thank for their contribution to my project and to helping me through recent really difficult times. Firstly, I would like to thank my supervisor, Rosie Bradshaw. Her never ending faith in me, encouragement, guidance, and kindness kept me going, and enabled me to reach a goal that I wasn't sure I could reach. Thank you so much to Tania and Paul, without whom none of this would have been possible. I couldn't have won the battle without such wonderful, dedicated, patient, and loving friends. Thank you also to my parents for their love, strength, and financial support. Thank you to my brother and sister for their support. Because my project was done in two parts there are lots of people to thank for getting me started, and other people to thank for re-teaching me and building up my confidence again. Thank you to all of you. Thank you to all the people I did fourth year with, for making such a difficult year more bearable. Thank you Dianne and Karyn for getting me started on my lab work. Thank you to Carolyn for her always useful advice, and for saving lots of my sequencing gels. Thanks also to Mike and Rich for helpful tips. Thanks to Austen for his friendship. Thank you to Tash for keeping my project going while I was away. Thank you to Anita for always being really positive, especially for making the lab a fun place to be in. Special thanks must also go to Linda for her friendship, and her patience in helping me belong again in the lab. Thank you to Brendon for reminding me how to run a sequencing gel, and to Paul for teaching me sequencing the first time around. Thanks to Bran for late night company in the lab. Thank you to David for drawing my chemical structures. Thanks again to Tania, this time for helping me draw my thesis figures. Thank you to all the Friday night staff club regulars for keeping me sane and refreshed. Thank you to all the other past and present people in the MGU I haven't mentioned. Thank you to all my friends outside of the department, especially Craig and Maree for their support from afar, and Justin for always being there if I needed him. I would also like to thank my flatmates for keeping me in touch with reality over the last few months, especially Tony "I love birds, eh" Roeven for his reassurance and tolerance. I'm sure I've forgotten heaps of people who also played a really valuable part in getting me where I am today, so even if you're not on this list I would like to thank you. Thanks must also go to Molecular Genetics Unit for providing the facilities and financial assistance that enabled me to undertake this project, especially Barry for getting our lab a freezer. In addition, thank you to John Linz (Michigan State University) for providing the *Asperillus parasiticus nor-1* and *ver-1* gene clones.

TABLE OF CONTENTS

| | | |
|-------------------|--|----|
| ABSTRACT | ii | |
| ACKNOWLEDGEMENTS | iii | |
| TABLE OF CONTENTS | iv | |
| LIST OF TABLES | xi | |
| LIST OF FIGURES | xii | |
| 1.0 | INTRODUCTION | 1 |
| 1.1 | General Features | 1 |
| 1.2 | Infection | 2 |
| 1.3 | Chemical Control | 4 |
| 1.4 | Resistant Strains | 5 |
| 1.5 | Dothistromin Toxin | 6 |
| 1.6 | Inactivation of Dothistromin | 10 |
| 1.7 | Aflatoxin Biosynthesis | 13 |
| 1.8 | Gene Cloning Strategies in <i>Aspergillus</i> Species | 14 |
| 1.9 | Organisation and Arrangement of AF/ST Biosynthetic Pathway Genes | 18 |
| 1.10 | Aims and Objectives | 20 |
| 2.0 | MATERIALS AND METHODS | 22 |
| 2.1 | Bacterial and Fungal Strains, λ Clones and Vectors | 22 |
| 2.2 | Media | 22 |
| 2.2.1 | Fungal Media | 22 |
| 2.2.1.1 | <i>D. pini</i> Media (DM) Broth | 22 |
| 2.2.1.2 | <i>D. pini</i> Media (DM) Agar | 22 |
| 2.2.1.3 | Malt Extract Agar (MEA) | 22 |
| 2.2.1.4 | Malt Yeast Glucose (MYG) Agar | 23 |

| | | |
|----------|------------------------------------|----|
| 2.2.1.5 | Yeast Peptone Glycerol (YPG) Agar | 23 |
| 2.2.1.6 | Potato Dextrose Agar (PDA) | 23 |
| 2.2.1.7 | Nutrient Malt Yeast (NMY) Agar | 23 |
| 2.2.1.8 | Minimal Medium (MM) Agar | 23 |
| 2.2.1.9 | Nutrient Yeast (NY) Broth | 23 |
| 2.2.1.10 | Nutrient Malt Yeast (NMY) Broth | 23 |
| 2.2.2 | Bacterial Media | 26 |
| 2.2.2.1 | Luria-Bertaini (LB) Media | 26 |
| 2.2.2.2 | TB Top Agar | 26 |
| 2.2.2.3 | NZCYM | 26 |
| 2.3 | Growth and Maintenance of Cultures | 26 |
| 2.3.1 | Fungal Cultures | 26 |
| 2.3.2 | Bacterial Cultures | 27 |
| 2.4 | Buffers and Solutions | 27 |
| 2.4.1 | TEG | 27 |
| 2.4.2 | Tris-Equilibrated Phenol | 27 |
| 2.4.3 | TE Buffer | 28 |
| 2.4.4 | DNase free RNase | 28 |
| 2.4.5 | 10 x TAE Buffer | 28 |
| 2.4.6 | 10 x Gel Loading Dye | 28 |
| 2.4.7 | 20 x SSC | 28 |
| 2.4.8 | TES (10/1/100) | 28 |
| 2.4.9 | 50 x Denhardt`s | 29 |
| 2.4.10 | SM Buffer | 29 |
| 2.4.11 | Acrylamide Mix | 29 |
| 2.4.12 | 10 x TBE Sequencing Buffer | 29 |

| | | |
|---------|--|----|
| 2.5 | DNA Preparations | 29 |
| 2.5.1 | Alkaline Lysis <i>E. coli</i> Plasmid Preparation | 29 |
| 2.5.2 | Large Scale <i>D. pini</i> Genomic DNA Preparation | 30 |
| 2.5.3 | Preparation of Genomic DNA for <i>D. pini</i> Library Construction | 31 |
| 2.5.4 | Mini-prep of λ Phage DNA | 32 |
| 2.5.5 | Extraction of DNA from Seaplaque Agarose | 33 |
| 2.5.5.1 | Bio 101 GeneClean kit | 33 |
| 2.5.5.2 | Gibco BRL GlassMax DNA Isolation Spin Cartridge System kit | 33 |
| 2.5.6 | Purification of DNA | 34 |
| 2.6 | DNA Manipulations | 34 |
| 2.6.1 | Restriction Enzyme Digests | 34 |
| 2.6.1.1 | Plasmid Digests | 34 |
| 2.6.1.2 | Genomic Digests | 35 |
| 2.6.2 | Agarose Gel Electrophoresis | 35 |
| 2.6.2.1 | Minigels | 35 |
| 2.6.2.2 | Overnight Gels | 35 |
| 2.6.3 | Determination of Fragment Sizes | 36 |
| 2.6.4 | Determination of DNA Concentration | 36 |
| 2.6.4.1 | Concentration Standards | 36 |
| 2.6.4.2 | Spectrophotometric Method | 36 |
| 2.6.4.3 | Fluorometric Method | 37 |
| 2.7 | Subcloning | 37 |
| 2.7.1 | Preparation of Insert DNA | 38 |
| 2.7.2 | Linearisation and CAP-Treatment of Vector DNA | 38 |
| 2.7.3 | Ligation | 38 |
| 2.8 | Transformation of <i>E. coli</i> | 39 |
| 2.8.1 | Calcium Chloride Transformation | 39 |

| | | |
|----------|--|----|
| 2.8.1.1 | Preparation of CaCl ₂ Competent Cells | 39 |
| 2.8.1.2 | Transformation | 39 |
| 2.8.2 | Transformation of <i>E. coli</i> by Electroporation | 39 |
| 2.8.2.1 | Preparation of Electro-Competent <i>E. coli</i> Cells | 39 |
| 2.8.2.2 | Electroporation | 40 |
| 2.9 | DNA Hybridisations | 40 |
| 2.9.1 | Southern Blotting | 40 |
| 2.9.2 | Random-Primer Labelling of Probes | 41 |
| 2.9.3 | Separation of Unincorporated Nucleotides by Minispin Column Chromatography | 42 |
| 2.9.4 | Hybridisation of Probe DNA to Southern Blots | 42 |
| 2.9.5 | Autoradiography of Southern Blots | 43 |
| 2.9.6 | Stripping Filters | 43 |
| 2.10 | Genomic Library Construction | 43 |
| 2.10.1 | Establishing Conditions for Partial Digestion of High Molecular Weight Genomic DNA | 43 |
| 2.10.2 | Large Scale Preparation of Partially Digested DNA | 44 |
| 2.10.3 | Partial Fill-in Reaction for Genomic DNA | 44 |
| 2.10.4 | Ligation of Insert to Vector Arms | 45 |
| 2.10.4.1 | Klenow/Ligation Controls | 45 |
| 2.10.4.2 | Determination of Optimum Ligation Conditions | 46 |
| 2.10.5 | Packaging of Ligated DNA and Titration of Recombinant Phage | 46 |
| 2.10.5.1 | Packaging of Ligated DNA | 46 |
| 2.10.5.2 | Titration of Packaged Phage on LB Plates | 46 |
| 2.10.6 | Large Scale Packaging of Ligated DNA | 47 |
| 2.10.7 | Amplification of the Library | 48 |
| 2.11 | Library Screening by Plaque Hybridisation | 49 |
| 2.11.1 | Plating Phage λ | 49 |

| | | |
|----------|--|----|
| 2.11.2 | Filter Lifts | 49 |
| 2.11.3 | Hybridisation of Phage λ DNA [α - 32 P]dCTP Labelled DNA | 50 |
| 2.12 | DNA Sequencing | 50 |
| 2.12.1 | Preparation of DNA for Sequencing | 50 |
| 2.12.1.1 | Preparation of Single Stranded M13 Template DNA | 50 |
| 2.12.1.2 | Preparation of Template DNA for AmpliCycle Sequencing | 51 |
| 2.12.2 | Sequenase Version 2.0 Protocol | 51 |
| 2.12.3 | AmpliCycle Sequencing Protocol | 52 |
| 2.12.4 | Polyacrylamide Gel Electrophoresis (PAGE) of Sequencing Reactions | 52 |
| 3.0 | RESULTS | 56 |
| 3.1 | Determining Optimal Mycelial Growth Conditions in Culture | 56 |
| 3.1.1 | Determining Optimal Solid Media for Mycelial Growth | 56 |
| 3.1.1.1 | Temperature Effect on Mycelial Growth | 56 |
| 3.1.1.2 | Media Effect on Mycelial Growth | 56 |
| 3.1.1.3 | Effect of Inoculation Method and Incubation Period on Mycelial Growth | 61 |
| 3.1.2 | Quantitation of Mycelial Growth in Liquid Media | 61 |
| 3.2 | Southern Hybridisations | 64 |
| 3.2.1 | Detection of a <i>D. pini</i> Region Heterologous to the <i>A. parasiticus</i> <i>nor-1</i> Gene | 64 |
| 3.2.2 | Detection of a <i>D. pini</i> Region Heterologous to the <i>A. parasiticus</i> <i>ver-1</i> Gene | 64 |
| 3.3 | Library Construction | 68 |
| 3.4 | Isolation of λ Clones Hybridising to the <i>A. parasiticus</i> <i>ver-1</i> Gene | 74 |
| 3.4.1 | Library Screening | 74 |
| 3.4.2 | Restriction Digestion of λ Clones and Southern Hybridisation | 74 |
| 3.5 | Further Characterisation of Clone λ CGV1 | 80 |

| | | |
|-------|---|-----|
| 3.5.1 | Mapping Clone λ CGV1 Further | 80 |
| 3.5.2 | Subcloning of a λ CGV1 Region Required for Sequencing | 85 |
| 3.5.3 | Sequence Analysis of Clone λ CGV1 | 85 |
| 3.5.4 | Sequence Identification | 85 |
| 3.5.5 | Sequence Comparison | 90 |
| 3.5.6 | Comparison of Intron Positions | 90 |
| 3.5.7 | Comparison of GC Content | 90 |
| 3.5.8 | Comparison of Codon Usage | 93 |
| 3.6 | Further Characterisation of Clone λ CGV2 | 93 |
| 3.6.1 | Further Hybridisation Analysis of Clone λ CGV2 | 93 |
| 3.6.2 | Subcloning of a λ CGV2 Region Required for Sequencing | 93 |
| 3.6.3 | Sequence Analysis of Clone λ CGV2 | 93 |
| 3.6.4 | Sequence Identification | 97 |
| 3.6.5 | Comparison Between <i>D. pini</i> Sequences | 97 |
| 3.6.6 | Comparison of GC Content | 97 |
| 3.7 | Isolation and Characterisation of λ Clone Hybridising to the <i>A. parasiticus nor-1</i> Gene | 97 |
| 3.7.1 | Library Screening | 97 |
| 3.7.2 | Restriction Digestion of Clone λ CGN2 and Southern Hybridisation | 100 |
| 4.0 | DISCUSSION | 104 |
| 4.1 | Identification of Clone λ CGV1 | 104 |
| 4.2 | Role of the <i>ver-1</i> -like Genes | 104 |
| 4.2.1 | The Role of the <i>Aspergillus ver-1</i> and <i>ver-A</i> Genes | 104 |
| 4.2.2 | The Role of the <i>ThnR</i> and <i>Thr1</i> Genes | 106 |
| 4.3 | Duplication of the <i>ver-1</i> Gene | 107 |

| | | |
|-------|---|-----|
| 4.4 | Identification of Clone λ CGV2 | 108 |
| 4.5 | Identification of Clone λ CGN2 | 110 |
| 4.6 | Potential Uses of the <i>D. pini ver-1</i> Gene | 111 |
| 4.6.1 | Cloning of Other Pathway Genes | 111 |
| 4.6.2 | Gene Disruptions | 113 |
| 4.6.3 | Identification of Molecular Mechanisms which Regulate Pathway Genes | 114 |
| 5.0 | SUMMARY AND CONCLUSIONS | 116 |
| | APPENDIX 1.0 | 118 |
| | APPENDIX 2.0 | 122 |
| | REFERENCES | 124 |

LIST OF TABLES

| | | |
|---------|---|-----|
| Table 1 | Fungal and Bacterial strains, λ Clones and Vectors | 24 |
| Table 2 | Primers used in Sequencing Reactions | 53 |
| Table 3 | Optimisation of Media and Method of Inoculation | 62 |
| Table 4 | Quantitation of Mycelial Growth with Malt Extract Concentration | 63 |
| Table 5 | Titres of Small Scale, Large Scale, and Amplified Libraries | 75 |
| Table 6 | Data from Restriction Mapping of Clone λ CGV1 | 81 |
| Table 7 | Data from Southern Hybridisation Analysis of Clone λ CGV2 | 82 |
| Table 8 | Codon Bias Table | 94 |
| Table 9 | Data from Southern Hybridisation Analysis of Clone λ CGN2 | 103 |

LIST OF FIGURES

| | | |
|------------|---|-----|
| Fig. 1 | Structures of sterigmatocystin, aflatoxin B1, and dothistromin | 8 |
| Fig. 2 | Comparison of the aflatoxin biosynthetic pathway with the proposed dothistromin biosynthetic pathway | 15 |
| Fig. 3A-F | Demonstration of media differences on mycelial growth and appearance | 57 |
| Fig. 4A-C | Southern blot of <i>D. pini</i> genomic DNA probed with <i>nor-1</i> | 65 |
| Fig. 5A-C | Southern blot of <i>D. pini</i> genomic DNA probed with <i>ver-1</i> | 69 |
| Fig. 6A-B | Profiles of partial <i>Mbo</i> I digestions of genomic DNA from <i>D. pini</i> | 72 |
| Fig. 7A-B | Mapping the position of <i>ver-1</i> on clone λ CGV1 | 76 |
| Fig. 8A-B | Mapping the position of <i>ver-1</i> on clone λ CGV2 | 78 |
| Fig. 9A-B | Further mapping analysis of clone λ CGV1 | 83 |
| Fig. 10A-B | Restriction map of clone λ CGV1 from a <i>D. pini</i> genomic library that hybridised to <i>Aspergillus parasiticus ver-1</i> | 86 |
| Fig. 11 | Partial sequence of the putative <i>D. pini ver-1</i> gene from clone λ CGV1 | 88 |
| Fig. 12 | Comparison of <i>D. pini</i> sequence to other amino acid sequences | 91 |
| Fig. 13 | Southern blot of λ CGV2 probed with the <i>D. pini ver-1</i> fragment | 95 |
| Fig. 14 | Partial sequence of a <i>ver-1</i> hybridising fragment from clone λ CGV2 | 98 |
| Fig. 15A-B | Southern blot of λ CGN2 probed with <i>nor-1</i> | 101 |