Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Influence of Dietary Fat Inclusion on the Ileal Digestibility of Amino Acids in Broilers

A thesis presented in partial fulfilment of the requirements for the degree

 \mathbf{of}

Master of Science

in

Animal Science



at Massey University, Manawatu,

New Zealand

Thomas Harrison Whitehouse

Acknowledgments

This project would not have been completed if not help from the Massey poultry unit. Special thanks to Ed James, Shaun de Malmanche, Elham Ebrazeh, Nipuna Perera, Asnath Mtei, Kaliwyn Pereka, Fifi Zaefarian, Colin Naftel and Naveed Anwar for help with diet preparation, sample collection and other aid with conducting the experiments. Thank you to my supervisors Ravi Ravindran and Reza Abdollahi for advice and aid with my project.

I would also like to thank the Massey University Nutrition Laboratory and Felicity Jackson for analyzing various digesta samples.

I would also like to thank my parents, Patrice and Thomas Whitehouse for their support during my master's degree studies.

List of Contents

Acknowledgments	!
List of Abbreviations:	iv
List of Tables	iv
Chapter 1: General introduction	1
Chapter 2: Literature review	2
2.1 Introduction	2
2.2 Fats in broiler diets	3
2.3 Endogenous amino acid losses	7
2.3.1 Measurement of basal endogenous losses	7
2.3.1.1 Protein-free diets	8
2.3.1.2 Regression method	9
2.3.1.3 Fasting method	9
2.3.2 Sources of endogenous protein	10
2.3.3 Effects of feed components	11
2.3.3.1 Non-starch polysaccharides	11
2.3.3.2 Phytate	13
2.3.3.3 Fats	14
2.4 Measurement of amino acid digestibility	15
2.5 Influence of fats on amino acid digestibility	15
2.6 Mode of action	20
2.6.1 Increase in gut retention time	20
2.6.2 Effects of fatty acid composition	21
2.6.3 Changes in digestive morphology	23
2.6.4 Bird Age	24
2.6.5 Other factors	24
2.7 Conclusions	27
Chapter 3: Effect of dietary fat on endogenous amino acid losses	28

3.1 Abstract	28
3.2 Introduction	28
3.3 Materials and Methods	30
3.3.1 Birds and housing	30
3.3.2 Diets	30
3.3.3 Digesta collection	30
3.3.4 Chemical analysis	31
3.3.5 Calculations	32
3.3.6 Statistical analysis	32
3.4 Results	33
3.4.1 Fatty acid composition	33
3.4.2 Ileal endogenous losses	34
3.5 Discussion	35
3.6 Conclusion	40
Chapter 4: The interaction between grain type and dietary fat source on nutrient digestibility in broad 41	ilers
4.1 Abstract	41
4.2 Introduction	41
4.3 Materials and methods	43
4.3.1 Diets	43
4.3.2 Birds and housing	44
4.3.3 Digesta collection	45
4.3.4 Chemical analysis	45
4.3.5 Calculations	46
4.3.6 Statistical Analysis	47
4.4 Results	47
4.4.1 Dry matter digestibility	47

4.4.2 Nitrogen digestibility	49
4.4.3 Starch digestibility	50
4.4.4 Fat digestibility	51
4.4.5 Gross energy digestibility	53
4.4.6 Amino acid digestibility at the ileal level	54
4.5 Discussion	57
4.5.1 Dry matter digestibility	57
4.5.2 Nitrogen digestibility	59
4.5.3 Starch digestibility	60
4.5.4 Fat digestibility	60
4.5.5 Gross energy digestibility	61
4.5.6 Amino acid digestibility at the ileal level	63
4.5 Conclusion	64
References	lxvi

List of Abbreviations:

AA: Amino acid(s)

CAD: Coefficient of apparent digestibility

DM: Dry matter

EL: Endogenous Loss(es)

GE: Gross energy

NSP: Non-starch polysaccharide(s)

List of Tables

Table 2.1: Fatty acid composition of common dietary fats3
Table 2.2: Effect of dietary fat sources of amino acid digestibility16
Table 2.3: Effect of differing dietary fat levels of amino acid digestibility in
pigs17
Table 2.4: Effect of differing dietary fat sources on fat and dry matter digestibility in
pigs19
Table 2.5: Effect of soybean oil and yellow grease on Dry Matte and Nitrogen
digestibility
pigs21
Table 2.6: Effect of dietary fat source on bacterial growth26
Table 3.1. Ingredient composition (%, as fed) of the N-free diets32
Table 3.2 Fatty acid composition of soybean oil and tallow (g/kg diet)33
Table 3.3: Influence of fat sources on the ileal endogenous amino acids in broilers (mg/kg dry
matter intake)34
Table 4.1: Experimental diet composition (g/Kg)43
Table 4.2: Influence of grain type and fat source on coefficient of apparent digestibility of dry
matter in different segments of the digestive tract in male broilers at 21 days post-hatch47

Table 4.3: Influence of grain type and fat source on coefficient of apparent digestibility of
nitrogen in different segments of the digestive tract in male broilers at 21 days post-hatch48
Table 4.4: Influence of grain type and fat source on coefficient of apparent digestibility of starch
in different segments of the digestive tract in male broilers at 21 days post-hatch49
Table 4.5: Influence of grain type and fat source on coefficient of apparent digestibility of fat in
different segments of the digestive tract in male broilers at 21 days post-hatch51
Table 4.6: Influence of grain type and fat source on coefficient of apparent digestibility of gross
energy in different segments of the digestive tract in male broilers at 21 days post-hatch53
Table 4.7: Influence of grain type and fat source on coefficient of apparent ileal digestibility of
indispensable amino acids in male broilers at 21 days post-hatch55
Table 4.8: Influence of grain type and fat source on coefficient of apparent ileal digestibility of
dispensable amino acids in male broilers at 21 days post-hatch56