Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

## Disturbance in the North Island of New Zealand: A case study using floodplain cores from the Coromandel to determine anthropogenic disturbance

A thesis presented in partial fulfilment of the requirements for the degree of

Master of Science

in Geography

Massey University, New Zealand

Elizabeth Grace Fox

2016



 ${\it Kuaotunu~Stream~with~the~surrounding~Kuaotunu~floodplain}.$ 

Photographer E.Fox, 25-03-2014.

## **Table of Contents**

List of Figures	4
List of Tables	5
Abstract	6
Acknowledgements	7
Chapter 1: Introduction	8
1.1 General Introduction	8
1.2 Purpose of Study	10
Chapter 2: Disturbance	11
2.1 Measuring Disturbance	11
2.2. Disturbance in New Zealand	17
2.2.1 Overview	17
2.2.2 New Zealand Setting	18
2.2.3 Disturbance prior to Human Occupation	18
2.2.5 Evidence of Anthropogenic Activities	23
Chapter 3: North Island Case Study Review	25
3.1 Aim	25
3.2 Method & Limitations	25
3.3 Results & Discussion	27
3.3.1 Disturbance Studies	37
3.3.2 The influence of Geology on Disturbance	38
3.3.3 Rainfall and Disturbance	39
3.3.4 Proxies used for determining Disturbance	41
3.4 Summary	44
Chapter 4: Case Study of the Coromandel	45
4.1 Environmental History	45
4.1.1 Geology	45
4.1.2 Geomorphology, climate and Vegetation	47
4.2 Anthropogenic History	49
4.3 Gold Mining History	51
4.4 Kuaotunu and Paeroa Mining History	51
4.4.1 Kuaotunu	51
4.4.2 Paeroa	54
Chapter 5: Methodology	57

5.1 Core collection	57
5.2 Geochemical Analysis	64
5.3 Stratigraphy and particle size analysis	64
5.4 Radiocarbon dating and sedimentation rates	66
Chapter 6: Results	67
6.1 Stratigraphy:	67
6.1.1 Maori Road Core:	67
6.1.2 Gerard Road Core:	74
6.1.3 Kuaotunu Floodplain Core:	74
6.1.4 Kuaotunu 3 Core:	74
6.1.5 Kuaotunu 4 Core:	75
6.1.6 Kuaotunu 5 Core:	75
6.2 PSA Results:	75
6.2.1 Maori Road Core:	79
6.2.2 Gerard Core:	79
6.2.3 Kuaotunu Floodplain Core:	80
6.2.4 Kuaotunu 3 Core:	80
6.2.5 Kuaotunu 4 Core:	80
6.3 Geochemistry Results:	87
6.3.1 Maori Road Core:	87
6.3.2 Gerard Core:	87
6.3.3 Kuaotunu Floodplain Core:	87
6.3.4 Kuaotunu 3 Core:	87
6.3.5 Kuaotunu 4 Core:	88
6.3.6 Kuaotunu 5 Core:	88
6.4 Age of cores & Sedimentation rates:	88
Chapter 7: Discussion	90
7.1 Disturbance in the Coromandel	90
7.1.1 Maori Road Core	90
7.1.2 Gerard Road	95
7.1.3 Kuaotunu Floodplain core	96
7.1.4 Kuaotunu 3 Core	99
7.1.5 Kuaotunu 4 Core	101
7.1.6 Kuaotunu 5 Core	102

7.2	Anthropogenic disturbance in New Zealand	103
7.3	Use of Geochemistry and Sedimentology	106
7	7.3.1 Geochemistry and sedimentology in New Zealand studies	106
7	7.3.2 Geochemistry and sedimentology as proxies for disturbance reconstructions	108
Chapt	er 8: Conclusion	111
Chapt	er 9: Future Recommendations	113
-	ences	
	ndices:	
пррсі		1 10
List	of Figures	
		Page
2.1	Changes in New Zealand vegetation cover. A: Pre-Maori, B: Pre-European and C: Present	22
	vegetation cover in 1976.	
3.1	Approximate locations of studies that reconstructed the past environment in the North	32
	Island, New Zealand	
3.2	Sites that showed anthropogenic disturbance and sites that did not.	33
3.3	Sites with the underlying geology	34
3.4	Sites with the Annual Rainfall	34
3.5	Sites colour coded based on the main proxy type used	35
4.1	Location of the Coromandel Peninsula in relation to the rest of New Zealand	46
4.2 A	Present day vegetation across the Coromandel Peninsula. Adapted from Landcare	48
	Research	
4.2 B	Potential Natural vegetation prior to human settlement across the Coromandel Peninsula	48
4.3	Areas of the Coromandel Peninsula that were occupied by Maori and Europeans, based	50
	on archaeological sites identified from first Maori settlement to 1980	
4.4	A) Location of the Kuaotunu and Paeroa areas where mining took place directly or	52
	upstream of the townships (red markers). B) Kuaotunu Catchment with key mining	
	locations (red pins). C ) Ohinemuri Catchment with key mining locations (yellow pins).	
4.5	Gold mining claims in the Kuaotunu area	53
5.1	Location of cores extracted from the Coromandel. A: New Zealand with Coromandel area	58
	in red box. B: Coromandel area with yellow box the Kuaotunu area and the orange box	
	the Paeroa area. C: Cores extracted from Kuaotunu D: Cores extracted from outside of	
	the Paeroa township	
6.1	Maori Road core lithology	68

6.2	Gerard Road core lithology	69
6.3	Kuaotunu Floodplain core lithology	70
6.4	Kuaotunu 3 core lithology	71
6.5	Kuaotunu 4 core lithology	72
6.6	Kuaotunu 5 core lithology	73
6.7	Gerard Road Core stratigraphy, PSA and geochemistry results	81
6.8	Maori Road core stratigraphy, PSA and geochemistry results	82
6.9	Kuaotunu Floodplain core stratigraphy, PSA and geochemistry results	83
6.10	Kuaotunu 3 Core stratigraphy, PSA and geochemistry results	84
6.11	Kuaotunu 4 Core stratigraphy, PSA and geochemistry results	85
6.12	Kuaotunu 5 Core stratigraphy and geochemistry results.	86
Liet	of Tables	
LIST	of Tables	Page
2.1	Summary of main proxies and age controls used in previous palaeo reconstructions.	12
3.1	Palaeo environmental studies conducted in and around the North Island, New Zealand,	28
	summary facts.	
3.2	Broad categorises used to group the environmental reconstructions and should be used	36
	in conjunction with Figure 3.3	
3.3	Evidence of increased sedimentation rate following European occupation	42
5.1	Coordinates of the locations of the 6 cores. Kuaotunu cores 3 & 4 were located less than	63
	2 metres apart therefore no distinct difference in location.	
6.1	Maori Road Core amount of reaction indication organic matter and flocculating of the	75
	samples on a nil to high scale.	
6.2	Kuaotunu Floodplain core amount of reaction indication organic matter and flocculating	76
	of the samples on a nil to high scale.	
6.3	Kuaotunu 3 core amount of reaction indication organic matter and flocculating of the	77
	samples on a nil to high scale.	
6.4	Kuaotunu 4 core amount of reaction indication organic matter and flocculating of the	77
	samples on a nil to high scale.	
6.5	Gerard Road core amount of reaction indication organic matter and flocculating of the	78
	samples on a nil to high scale.	
6.6	Calculated Sedimentation rates before and after mining Kuaotunu and Paeroa	88

## Abstract

It is well documented that following human occupation of a region, the surrounding environment may undergo drastic changes through vegetation pattern alterations, displacement of fauna, alteration of sedimentation and fluvial regimes, and changes to the composition of the underlying material. Many case studies of anthropogenic disturbance have been conducted in New Zealand. One of the main outcomes of this research is to collate, contrast and compare this wealth of case studies to look for any underlying trends in timing, distribution and magnitude of disturbance nationwide. This thesis focusses on late Holocene records from the North Island, and compared the history of disturbance with that from the South Island (as per McWethy et al. 2010). Based on the combination of palynology, sedimentology and geochemistry, this review demonstrates the pace of disturbance observed in the North Island was very rapid following occupation, a trend also established in the South Island.

The other main outcome of this research is to add to the knowledge base of North Island disturbance history, through development of a landscape disturbance history in the Coromandel, using floodplain cores from the Paeroa and Kuaotunu areas. Sediment logging and subsequent XRF-geochemical analysis performed on these cores revealed a 'mining layer' that was used as a baseline for mining disturbance in this environment. This layer is interpreted as when European activities began disturbing the environment. Cores extracted from the Paeroa area indicated that the sedimentation rates in the floodplain had increased more than 15-fold since human occupation. Significant rises in the amount of Arsenic and Lead contained within the sediment were also detected. Cores from the Kuaotunu floodplain also showed changes in geochemistry that coincided with historic mining in the area, but reverted back to near pre-mining levels following the initial disturbance. These results suggest that factors such as catchment characteristics and degree of disturbance in an area affect the extent of impact on a site, which may have implications for future management of post mining sites. XRF analysis is a relatively underutilized proxy in New Zealand. It, in conjunction with Particle Size Analysis, has proved valuable in this study and

are recommended for application in future New Zealand environmental reconstructionfocused research.

## **Acknowledgements**

Dr. I. Fuller, Dr. A. Clement and Dr. K. Holt for their support and advice with identifying sediment layers, geochemistry interpretation and assistance with coring and logging the floodplain cores. A big thank you to Mr. D. Feek for his assistance in the field and Dr. A. Moebis for her help with the processing of the subsamples and the use of the Particle Laser Scanner. A special thank you to Dr. T. Novakova at Aberystwyth University for carrying out the XRF NITON scanning to deduce the geochemistry of the Coromandel Cores and provision of these results.

A special thank you to Mr. B Watson, Mr J. Watson and Mrs M. Rose at Horizons Regional Council for their enthusiasm, support and encouragement in the undertaking of this thesis.

And finally a huge thank you to my family and friends for their continued support and motivation they gave me throughout this project.