Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

# REGULATION OF THE ASPERGILLUS NIDULANS CYTOCHROME C GENE

A thesis presented in partial fulfilment of the requirements for the degree of Master of Science in Molecular Genetics at Massey University, Palmerston North New Zealand

Linda Joy McGlynn

1997

#### **ABSTRACT**

The filamentous fungus Aspergillus nidulans has been genetically and biochemically well-characterised and thus provides an attractive model for studies on the regulation of eukaroytic gene expression. This study was undertaken to investigate the factors affecting respiratory function in A. nidulans. Due to the central role of cytochrome c in oxidative respiration, this study was focused on the cytochrome c gene and primarily upon how oxygen availability affects its expression.

The Aspergillus nidulans cytochrome c gene (cycA) appears to be transcriptionally activated in response to oxygen availability (Raitt et al., 1994). In the yeast S. cerevisiae, oxygen availability activates its cytochrome c genes via a heme-activated protein HAP1, which binds to the promoter region of each gene (Pfiefer et al., 1989). Since heme is only synthesised in the presence of oxygen, activity of the HAP1 protein serves as an intracellular signal of oxygen availability.

In the upstream region of the cycA gene, a sequence with homology to the S. cerevisiae HAP1 binding site was present (Raitt, 1992). To determine the significance of the putative HAP1 binding site and the role of other promoter sequences in the A. nidulans cycA gene, a promoter-reporter vector was constructed. However, upon sequencing of the cycA promoter in the reporter vector, a sequencing error was discovered in the published cycA gene by Raitt et al. (1994) which affected the position of the major translational start site. Further examination of the cycA sequence also revealed a possible undetected intron (Intron I). To determine the number of introns in the cycA gene, RT-PCR was performed on cycA RNA. Sequencing of the RT-PCR amplified products showed that the previously undetected intron (Intron I) was present, and that the cycA gene contains three instead of two introns as published by Raitt et al. (1994). Since the published ATG start site was located within Intron I, a new translational start site was proposed. The major consequences of these changes to the cycA gene was that the putative HAP1 site was now located within the coding region of the gene, and therefore could not be a regulatory element. In addition only 247 bp of cycA promoter sequence remained cloned for analysis. To obtain additional promoter sequence, an A. nidulans genomic library was screened with a BamHI cDNA probe containing 224 bp of the 5' region of the cycA gene. Three positive clones were obtained, of which  $\lambda LM9$  and  $\lambda$ LM5 were identical, and  $\lambda$ LM19 was an overlapping clone with  $\lambda$ LM9 and  $\lambda$ LM5. Restriction enzyme and Southern blott analysis of the two overlapping cycA clones, revealed that 2.1 kb EcoRI fragments from both clones contained the 5' region. The 2.1 kb EcoRI fragment from λLM9 was cloned into pUC18 and sequenced. The completed upstream sequence of the A. nidulans cytochrome c gene was obtained, and putative regulatory signals including the HAP1 binding site were found, and compared with published promoter sequences which regulate the expression of respiratory-encoding genes from yeast.

#### **ACKNOWLEDGEMENTS**

Firstly, I would like to thank my supervisor Rosie Bradshaw, for keeping faith in me, and for her continual enthusiasm and encouragement, even when my thesis was not going according to plan! Her support was much appreciated.

I would also like to thank all past and present members of the Microbiology and Genetics Department. Thanks everyone for your friendship, you guys made it a great place to work in. In particular, I would like to thank our neighbouring lab, Scott base, who we were always borrowing things from, especially Carolyn Young, who somehow found time out of her busy research to give all of us MGU students loads of time saving advice, Mike Collett for answering all my questions so extensively and patiently, and Austen Ganley for being around to listen and to entertain us out at the farm. This also applies to Kate and Mike Collett who we shared many yummy dinners and good times with.

Many thanks goes to everyone from my lab, Dianne-for all her *Aspergillus* related advice, Paul-for providing lots of laughs, Carmel-thanks for being such a great friend, Anita-for her lively personality which helped keep us sane, Branwen-for your support especially during those hard times, Brendon-for all those friendly chats, and Bek-for technical support, I hope you have more luck with this project than I did! Also thanks to Tania who was always popping into our lab with lots of smiles.

A big thank you must also go to my family for their love and support. I would also like to thank my special friend Rowena. Thanks for putting up with me, especially during our four years flatting together, and for all your strength and encouragement, which went along way towards the accomplishment of this thesis.

Lastly, I have Richard to thank for many many things, I hardly know how to adequately express my gratitude and love. Thank you Rich for giving me your support, love, encouragement, guidance, cooking more than I did, and all those sacrifices you made because I was too busy! Also I must mention that your scientific advice was invaluable. I am very lucky to have you for a future husband!

### TABLE OF CONTENTS

ABS'	<u>TRACT</u> i
<u>ACK</u>	NOWLEDGEMENTSiv
TABL	LE OF CONTENTS
LIST	OF TABLES.
LIST	OF FIGURES xi
CHAF	PTER 1. INTRODUCTION
1.1	THE ROLE OF THE CYTOCHROME C PROTEIN
1.2	WHY STUDY THE ASPERGILLUS NIDULANS CYTOCHROME C GENE?
1.3	REGULATION OF THE SACCHAROMYCES CEREVISIAE  CYTOCHROME C GENES
1.3.1	HAP1 Regulation of CYC1 and CYC7 in Response to Oxygen  Availability
1.3.1.	Structure and Function of the HAP1 Protein
1.3.1.2	Multi-sequence Recognition by HAP14
1.3.2	Regulation of CYC1 and CYC7 in Response to Carbon Source by
	the Yeast CCAAT-binding Factor
1.3.2.	The Yeast CCAAT-binding Factor (HAP2/3/4/5 Complex)8
1.3.3	Other factors affecting CYC7 expression
1.4	REGULATION OF THE KLUYVEROMYCES LACTIS  CYTOCHROME C GENE (K1CYC1)
1.5	REGULATION OF THE ASPERGILLUS NIDULANS CYTOCHROME C GENE (cycA)
1.4.1	Regulation of the cycA gene in response to Environmental Parameters12
142	Analysis of the 5' Region of the cycA gene 13

1.4.2.1 Putative HAP1 Binding Site
1.4.2.2 Putative Upstream Opening Reading Frame (uORF)13
1.5 AIMS AND OBJECTIVES
CHAPTER 2. MATERIALS AND METHODS
2.1 FUNGAL AND BACTERIAL STRAINS, LAMBDA CLONES
AND PLASMIDS17
2.2 MEDIA
2.2.1 Luria Broth
2.2.3 MYG Media
2.2.4 SOC Medium
2.2.5 NZCYM
2.2.6 TB Top Agarose
2.2.0 TB Top Agarose
2.3 COMMON BUFFERS AND SOLUTIONS
2.3.1 1 x TAE Buffer
2.3.2 1 x TBE Buffer
2.3.3 TE Buffer (10/1)
2.3.4 TE Buffer (10/0.1). 19
2.3.5 10 x Sequencing TBE Buffer
2.3.6 TES (10/1/100)
2.3.7 STET Buffer
2.3.8 SM Buffer:
2.3.9 Hybridisation Solution 19
2.3.10 50 x Denhardt's Solution
2.3.11 Tris-Equilibrated Phenol
2.3.12 20 x SSC
2.3.13 10 x Gel Loading Buffer
2.3.14 Standard DIG Buffer
2.3.15 DIG Buffer 1
2.3.16 DIG Buffer 2
2.3.17 DIG Buffer 3
2.3.18 Blocking Reagent stock solution
2.3.19 Alkaline Lysis Solution I
2.3.20 Alkaline Lysis Solution II

2.3.21 Alk	aline Lysis Solution III	20
2.4 GROW	TH AND MAINTENANCE OF CULTURES	20
2.5 DNA	PREPARATIONS	20
2.5.1 Rapid	1 Boiling Plasmid Preparation	20
2.5.2 Small	Scale Alkaline Lysis Plasmid Preparation	20
2.5.3 Large	Scale Alkaline Lysis Plasmid Preparation	21
2.5.4 Puri	fication of Plasmid DNA by Cesium Chloride-Ethidium Bromide	
Grad	dient	22
2.5.5 Prepa	aration of DNA from Fungal Cultures	22
2.5.6 Prepa	ration of Lambda Phage DNA by Liquid Lysate	23
2.5.7 Puri	fication of DNA by Phenol-Chloroform Extraction and Ethanol	
Pred	cipitation	23
2.6 STANI	DARD PROCEDURES	24
	striction Enzyme Digests	
	crose-gel Electrophoresis of DNA	
	mination of Molecular Weights	
	mination of DNA Concentration	
2.6.4.1	Quantification of DNA Concentration by Electrophoresis	
2.6.4.2	Spectrophotometric Determination of DNA Concentration and	
	Purity	25
2.7 AMPL	FICATION OF DNA BY THE POLYMERASE CHAIN REACTION	
(PCR)		25
2.8 DNA C	LONING PROCEDURES	26
2.8.1 Gel P	Purification of DNA	28
2.8.2 CA	P Treatment of DNA	28
2.8.3 Ligat	ion Reactions	28
2.8.4 Trans	formation of Escherichia coli	29
2.8.4.1	Preparation of Electroporation Competent E. coli Cells	29
2.8.4.2	Electroporation of E. coli	29
2.8.4.3	Preparation of CaCl2-Competent Cells	30
2.8.4.4	CaCl2 Transformation of E. coli	30
2.9 COLO	NY HYBRIDISATION USING THE DIG DETECTION SYSTEM	30
2.9.1 Colo	ny Lifts	30
2.9.2 Colo	ny Hybridisation	31

2.9.3 Preparation of DIG-Labelled Probe	1
2.9.4 Detection of Hybridised DIG-labelled Probe	2
2.10 DNA SEQUENCING	2
2.10.1 Sequenase Version 2.0 Sequencing Protocol	2
2.10.2 Amplicycle Sequencing Kit Protocol	3
	_
2.11 RNA PROCEDURES	
2.11.1 Precautions Taken for Working With RNA	
2.11.2 Agarose-gel Electrophoresis of RNA	
2.11.3 Determination of Fungal RNA Concentration and Purity	4
2.12 RNA PREPARATION	4
2.12.1 Aspergillus nidulans Spore Suspension Preparation	
2.12.2 Isolation of RNA from Aspergillus nidulans	
2.13 RT-PCR ANALYSIS OF RNA	
2.13.1 Reverse Transcriptase Synthesis of cDNA	
2.13.2 Amplification of cDNA by PCR	6
2.14 SOUTHERN BLOTTING AND HYBRIDISATION	6
2.14.1 Southern Blotting	
2.14.2 Preparation of [a-32P]dCTP-Labelled DNA Probe	
2.14.3 Southern Blot Hybridisation	
2.14.4 Stripping Probes from Southern Blots	
2.15 LIBRARY SCREENING BY PLAQUE HYBRIDISATION	8
2.15.1 Plating Phage $\lambda$	8
2.15.2 Plaque Lifts	9
2.15.3 Hybridisation of $\lambda$ Phage DNA to [a-32P]dCTP-Labelled Probe	9
CHAPTER 3. CONSTRUCTION OF PROMOTER-REPORTER VECTORS	
AND DISCOVERY OF A SEQUENCING ERROR	0
THE DISCOVERY OF THE CONTROL BRITISH	
3.1 INTRODUCTION	0
3.2 CLONING STRATEGY	0
3.2.1 The lacZ Reporter Vector	
3.2.2 PCR Amplification of the <i>cycA</i> Promoter Fragments	
3.2.3 General Preparation of pAN923-42B Vector and 0.68 kb cycA Promoter 43	

3.3 LIGATIONS AND TRANSFORMATIONS OF THE 0.68 kb cycA
PROMOTER46
3.3.1 Ligation of 0.68 kb promoter directly into pAN923-42B
3.3.2 Ligation of 0.68 kb Promoter into T-Tailed pUC18 Vector
3.3.3 Ligation of 0.68 kb Promoter into Commercially CAP-treated pUC1855
3.3.4 Ligation of 0.68 kb promoter into pAN923-42B
3.4 LIGATIONS AND TRANSFORMATIONS OF THE $cycA$ $\Delta$ HAP1
PROMOTER58
3.5 SEQUENCING OF THE 0.68 kb PROMOTER IN PLASMID R122
CHAPTER 3 DISCUSSION 64
3.6 PROBLEMS WITH CLONING INTO pAN923-42B VECTOR
2.7 MANUACATIONS FROM THE DISCOVERY OF THE SEQUENCING
3.7 IMPLICATIONS FROM THE DISCOVERY OF THE SEQUENCING
ERROR IN THE PUBLISHED cycA GENE BY RAITT et al (1994)
CHAPTER 4. RT-PCR ANALYSIS OF cycA RNA 69
4.1 INTRODUCTION
4.2 DEVELOPMENT OF A RT-PCR STRATEGY
4.2 DEVELOPMENT OF A RI-PCR STRATEGI
4.3 RT-PCR RESULTS
4.3.1 Sequencing of RT-PCR Products
CHAPTER 4 DISCUSSION 83
A A THE SEA CENE CONTAINS A THIRD INTRON AND HAS A
4.4 THE cycA GENE CONTAINS A THIRD INTRON AND HAS A
CORRECTED N-TERMINAL REGION
4.5 ALL THREE INTRONS OF THE cycA GENE ARE LOCATED AT
CONSERVED POSITIONS
CHAPTER 5. CLONING AND ANALYSIS OF THE 5' REGION OF THE
cycA GENE
5.1 INTRODUCTION
5.2 CORPENING OF AN A MIDIN ANGLED AND
5.2 SCREENING OF AN A. NIDULANS LIBRARY

5.3 SOUTHERN BLOTTING AND RESTRICTION MAPPING OF POSITIVE	
λ CLONES	. 89
5.4 SEQUENCE ANALYSIS OF CLONED $\lambda$ DNA SHOWN TO HYBRIDISE	
TO THE cycA GENE	. 90
5.4.1 Subcloning and Sequencing of the 2.1 kb Fragment	. 90
5.4.2 Confirmation that the 2.1 kb Fragment Corresponds to Regions of the	
A. nidulans A18 Genome	. 100
5.4.3 Analysis of the cycA promoter sequence	. 100
5.4.3.1 Putative HAP1 UAS	. 100
5.4.3.2 Putative HAP2/3/4/5 (Yeast CCAAT-binding Factor) UAS	. 100
5.4.3.3 Putative CREA Binding Site	. 105
5.4.3.4 Other Regulatory Elements	. 105
CHAPTER 5 DISCUSSION	. 106
<u></u>	
5.5 TATA BOX	. 106
5.6 THE ANCF (A. NIDULANS CCAAT BINDING FACTOR) COMPLEX	
MAY REGULATE THE CYCA GENE	. 106
5.7 THE A. NIDULANS CYCA GENE MAY BY REGULATED BY A HAPI-	
LIKE PROTEIN	108
CHAPTER 6. SUMMARY AND CONCLUSIONS	110
APPENDICES	. 112
Appendix 1.0 Vector Maps	
Appendix 2.0 Sequence Data	114
<u>REFERENCES</u>	110
KLI LKLIVCLO	117

### LIST OF TABLES

Table 2.1. Strains, $\lambda$ clones and plasmids	18
Table 2.2. PCR and Sequencing Primers	27
Table 3.1. Transformations A, B & C (with pAN923-42B vector)	47
Table 3.2. Expected Restriction Digestion Fragment Profiles for Ligation of 0.68	
kb Promoter into pAN923-42B	49
Table 3.3. Transformation D (with pAN923-42B vector)	52
Table 3.4. Expected Restriction Enzyme Profiles for Ligation of 0.68 kb Promoter into Plasmid pUC18	54
Table 3.5. Transformations E & F (with CAP-treated pUC18 vector)	56
Table 3.6. Transformation G (with pAN923-42B vector)	57
Table 4.1. PCR Products Expected From RT-PCR Analysis	74
Table 5.1. Restriction Enzyme Digestions of λ clones	95

### LIST OF FIGURES

Figure 1.1. Projection of HAP1 binding at <i>CYC7</i> and <i>CYC1</i> onto DNA helices (Pfiefer et al., 1987)
Figure 1.2. Comparison of HAP1 binding sites in S. cerevisiae and A. nidulans 14
Figure 3.1. The lacZ reporter vector pAN923
Figure 3.2. PCR primer binding sites within the cycA promoter fragments 44
Figure 3.3. Restriction enzyme analysis of plasmids pAN923, R122 and R120 50
Figure 3.4. Comparison of the <i>cycA</i> nucleotide sequence from plasmid R122 with the published <i>cycA</i> sequence (Raitt, 1992)
Figure 3.5. Autoradiograph of sequence obtained from the <i>cycA</i> gene cloned into plasmid R41
Figure 3.6. The published nucleotide sequence of the <i>cycA</i> gene (Raitt <i>et al.</i> , 1994)
Figure 3.7. How the sequencing error affects the cycA amino-terminal sequence 66
Figure 4.1. Re-estimation of the <i>cycA</i> transcription start point
Figure 4.2. RT-PCR outline showing the relative positions of primers
Figure 4.3. RT-PCR analysis of total RNA from A. nidulans strain A18
Figure 4.4. cDNA sequence of the 596 bp RT-PCR product 79
Figure 4.5. The corrected nucleotide sequence of the <i>cycA</i> gene
Figure 4.6. Alignment of cytochrome c N-terminal protein sequences
Figure 4.7. Alignment of three cytochrome c protein sequences showing intron positions
Figure 5.1a. Restriction digestion profiles of λLM9 and λLM19
Figure 5.1b. Southern Blot of λLM9 and λLM19
Figure 5.2. Restriction maps of λLM9 and λLM19 isolated from an <i>A. nidulans</i> genomic library that hybridised to the <i>cvcA</i> gene

Figure 5.3. Nucleotide sequencing strategy for the 5' region of the cycA gene	96
Figure 5.4. Nucleotide sequence of the 5' region of the cycA gene	98
Figure 5.5a. Restriction digestions of <i>A. nidulans</i> A18 genomic DNA	101
Figure 5.5b. Southern blot of A. nidulans A18 genomic DNA	101
Figure 5.6. Comparison of known yeast and putative A. nidulans HAP1 UASs	103

### CHAPTER 1 INTRODUCTION

#### 1.1 THE ROLE OF THE CYTOCHROME C PROTEIN

Cytochrome c is a highly conserved and well characterised protein which forms a key component of the respiratory pathway. The essential function of this process is oxidative phosphorylation from which energy is derived in the form of ATP. Although this pathway operates in the mitochondria, cytochrome c and some other respiratory components are encoded in the nucleus. Once synthesised these nuclear-encoded components are transported to the inner mitochondrial membrane for assembly of the respiratory chain. Respiratory components are thus encoded by both the nuclear and mitochondrial genomes. Coordinate expression of nuclear and mitochondrial genes is therefore required to achieve a balanced production of protein for assembly of respiratory complexes in the mitochondria. Coordinated regulation of these genes also occurs in response to mitochondrial and environmental conditions of the cell, for example, oxygen availability and respiratory status.

### 1.2 WHY STUDY THE ASPERGILLUS NIDULANS CYTOCHROME C GENE?

Extensive studies of the relationship between respiratory activity and environmental conditions have been carried out with the yeast *Saccharomyces cerevisiae*, a facultative anaerobe. In particular the effect of oxygen availability and carbon source on the expression of the cytochrome *c* genes from *S. cerevisiae* has been well characterised (for a review see Forsburg and Guarente, 1989a; Zitomer and Lowry, 1992). In contrast, limited research in this field has been conducted on the filamentous fungus, *Aspergillus nidulans*, and it is not known whether the cytochrome *c* gene of this obligate aerobe is regulated through similar *cis* and *trans* regulatory signals.

Recent research performed on the aerobic respiratory yeast *Kluyveromyces lactis*, has revealed that its cytochrome c gene (*KlCYC1*), is transcriptionally regulated by oxygen in a similar manner to the S. cerevisiae cytochrome c genes, despite differences in their respiratory metabolism (Freire-Picos et al., 1995). However, it is not clear whether the *KlCYC1* gene is regulated similarly by carbon source (Freire-Picos et al., 1993; Mulder et al., 1994; Freire-Picos et al., 1995; NGuyen et al., 1995; Mulder et al., 1995). Thus it seems likely that control of respiration in A. nidulans will be mediated by common regulatory pathways, such as those established for S. cerevisiae. It will be interesting to determine how regulation of the A. nidulans cytochrome c gene differs to that of the S.

cerevisiae cytochrome c genes, and may provide insight into the obligate aerobic nature of some fungi. A. nidulans is an ideal candidate for this comparison, as it provides a well established model system, where many genes have been studied in detail at the molecular and biochemical level (Arst and Scazzocchio, 1985; Davis and Hynes, 1991).

Environmental factors found to affect the regulation of the cytochrome c genes of S. cerevisiae and A. nidulans (such as carbon source, cell growth rate, and oxygen availability), alter the respiratory state and cellular metabolism of the organism (Forsburg and Guarente, 1989; Pillar and Bradshaw, 1991; Zitomer and Lowry, 1992; Raitt et al., 1994). In some Aspergillus species, the control of the organism's respiratory state is important for the production of useful secondary metabolites. For example, if oxidative respiration is restricted, acetyl CoA accumulates and becomes available as a precursor for the biosynthesis of secondary metabolites. For one particular Aspergillus species, A. parasiticus, a highly toxic metabolite can be produced from acetyl CoA called aflatoxin. Biosynthesis of aflatoxin has been demonstrated to be enhanced under conditions of low oxygen and high glucose levels, which minimise oxidative respiration and enhance glycolysis (Shih & Marth, 1974). In addition, oxygen limitation can be a serious problem in large-scale chemostat cultures of obligately aerobic fungi, (for example; penicillin-producing Penicillium chysogenum), where high energy input is required to sparge and mix air in the vessel.

Therefore, the study of the regulation of the cytochrome c gene of *A. nidulans* should prove to be both interesting and of commercial value.

## 1.3 REGULATION OF THE SACCHAROMYCES CEREVISIAE CYTOCHROME C GENES

The yeast, S. cerevisiae, is unusual in that it has two forms of cytochrome c encoded by the independent genes, iso1-cytochrome c (CYC1, Smith et al., 1979) and iso2-cytochrome c (CYC7, Montgomery et al., 1980). CYC1 and CYC7 are isofunctional with 78% identity at the amino acid level (Montgomery et al., 1980). CYC1 accounts for 95% of the cytochrome c which is expressed in aerobic conditions, with CYC7 making up the remaining 5% (Creusot et al., 1988). The minor isoform of cytochrome c, CYC7, is regulated coordinately with CYC1 in response to various factors. The two principal signals that affect the respiratory ability of yeast are carbon source and oxygen. The response to these environmental parameters occurs through a specific interaction of hemeresponsive regulatory proteins at upstream activation sites (UAS) located in the promoter region of each cytochrome c gene. CYC1 contains two functional UAS sequences,

UAS1 and UAS2 (Guarente et al., 1984), whereas CYC7 contains one UAS (Prezant et al., 1987).

## 1.3.1 HAP1 Regulation of CYC1 and CYC7 in Response to Oxygen Availability

Both CYC1 and CYC7 are transcriptionally activated by oxygen, with CYC1 induced 200-fold (Guarente and Mason, 1983) and CYC7 induced about 20-fold (Prezant et al., 1987). Under anaerobic conditions, CYC7 is weakly repressed by lack of oxygen, whereas CYC1 is strongly repressed. The CYC7 gene product is also believed to have a longer half-life in anaerobic conditions which may poise the cell for a transition from anaerobic to aerobic growth conditions (Pillar & Bradshaw, 1991).

The CYC1 and CYC7 genes (and other genes encoding respiratory functions or oxidative damage repair functions), are not activated directly by molecular oxygen. Instead intracellular levels of heme serve as intermediate signalling molecules to convey information about oxygen levels. Since heme requires oxygen for its biosynthesis, heme levels ultimately reflect oxygen levels (Zitomer & Lowry, 1992). Heme exerts its regulatory role on CYC1 and CYC7 through interaction with a heme-activated protein, HAP1. HAP1 (also called CYP1), binds to UAS1 of CYC1 and to the UAS of CYC7 to activate transcription in the presence of heme (Pfiefer et al., 1987).

#### 1.3.1.1 Structure and Function of the HAP1 Protein

The HAP1 protein is a member of the zinc cluster proteins (Creusot *et al*, 1988; Pfiefer *et al*, 1989), and has a general function as a coordinator of the activity of cytochrome-encoding and related genes involved in oxygen regulation (Lodi and Guiard, 1991). Deletion analysis of HAP1 revealed three functional domains: the DNA binding domain, the activation domain and the heme-responsive domain (Pfiefer *et al*, 1989).

The activation domain is located at the carboxyl end of HAP1 between residues 1308-1483. This region is involved in transcriptional activation and is highly acidic, which is consistent with other activators such as GAL4.

The DNA binding domain lies between residues 1 and 148, and specifies binding to UAS1 of CYCI and CYC7. It contains a zinc finger known as a binuclear zinc cluster, which has extensive homology to the well characterised yeast GAL4 zinc cluster protein. The cluster is involved in non-specific DNA binding, with specific amino acids surrounding the cluster involved with specific recognition of each cytochrome c gene

(Pfeifer et al., 1989). For example, serine-63 at the base of the zinc cluster is essential for binding to the UAS1 of CYC1.

Mutational analysis of the HAP1 DNA binding domain (Turcotte and Guarente, 1992), found that HAP1 uses a cofactor at *CYC7* and potentially a different cofactor at *CYC1*. They proposed that the difference in sequence of both UAS sites could specify which cofactor is bound.

The heme-responsive domain has been proposed to mediate heme control of HAP1 activity and is located between residues 245-445. This domain contains seven adjacent repeat units of a conserved K/R C P I/V D H amino acid motif, resembling a metal or heme-binding site, and is therefore believed to be the site for the direct binding of heme (Creusot *et al.*, 1988; Pfiefer *et al.*, 1989).

In the absence of heme, an unknown cellular factor or factors was found to bind to the heme domain at the K/R C P I/V D H amino acid motif of HAP1 to inhibit DNA binding (Fytlovich *et al.* 1993; Zhang and Guarente, 1994a). This interaction with the cellular repressor(s) may weaken DNA binding by blocking HAP1 dimerisation, the active binding form of the protein (Zhang *et al.*, 1993). Consistent with this result, Zhang and Guarente (1994) found that HAP1, like steroid hormone receptors, was sequestered in a high molecular weight complex by one or more cellular factors in the absence of heme, thereby blocking dimerisation. However, the dissociation of the high molecular weight complex to the dimer form produced only partially active HAP1, and was therefore suggested to be only one of the events leading up to the activation of the HAP1 protein.

Recently, HAP1 mutants were obtained which were independent of heme when assayed for activation (Haldi and Guarente, 1995). These mutants define an additional region of HAP1 which is heme-responsive, located near the activation domain. Another heme-independent mutant has also been discovered, but the mutation is located near the heme-responsive domain (Ushinsky and Keng, 1994). These multiple heme-responsive domains have been suggested to allow a fine-tuned response to oxygen availability.

#### 1.3.1.2 Multi-sequence Recognition by HAP1

The UAS regions of CYC1 and CYC7 bound by HAP1 share no obvious sequence similarity. Both sites compete for binding to HAP1 with comparable affinities (Pfiefer et al., 1987; Ha et al., 1996), but HAP1 activates transcription at CYC1 10 fold more efficiently than at CYC7 (Turcotte & Guarente, 1992). HAP1 binds to these UAS sites as a homodimer (Zhang et al., 1993).


Despite the dissimilarity in sequence between the two HAP1 binding sites, overall HAP1 contacts both sites similarly (Figure 1.1; Pfiefer *et al*, 1987). For both sites, binding of HAP1 occurs primarily to one side of the DNA helix, protecting 23 base pairs. Crucial contacts are made in both the major and minor grooves with a single major groove contact on the reverse face of the helix at one end of each binding site. However, due to the dissimilar nature of the two sites, the actual positions of the major and minor groove contacts in *CYC1* and *CYC7* differ. The best alignment of the HAP1 binding sites from this study was obtained by inverting the *CYC7* sequence. This results in a 7 nucleotide match from the 23 nucleotides protected by HAP1.

Further information on the binding specificity of HAP1 was obtained by selecting random HAP1 binding sites using a random oligonucleotide-PCR selection technique (Zhang and Guarente, 1994b). This study showed that HAP1 unlike the other zinc cluster proteins, including the GAL4 protein, binds to a direct repeat of two CGG triplets rather than a palindromic sequence. In addition, the HAP1 site contains a 6 bp spacer between the CGG triplets, with an asymmetrically positioned TA sequence critical for HAP1 binding. However, the UAS of CYC7 does not fit this consensus as it contains two copies of CGC rather than CGG, and a more specific 6 bp spacer of TATTAT. The spacing between the CGG triplet repeats has been proposed to be important in controlling the binding specificity of protein members of the zinc cluster family. For example, studies on GAL4 (which contains a 11 bp spacer) showed that binding was greatly reduced with a decrease in spacing from 11 to 10 bp (Vashee et al., 1993). Therefore the 6 bp spacing observed for the HAP1 binding site was suggested to be crucial for HAP1 binding.

Recently, random and site-directed mutagenesis of the CYC1 and CYC7 HAP1 binding sites showed that all HAP1 sites, including those from CTT1 (catalase T), CYB2 (cytochrome b<sub>2</sub>) and CYT1 (cytochrome c<sub>1</sub>) genes, are related imperfect direct repeats with the 'optimal' sequence CGG N<sub>3</sub> TA N CGG N<sub>3</sub> TA (Ha et al., 1996). Mutation of either CGG triplet in CYC1 decreased in vivo activity and in vitro binding of HAP1, indicating that the CGG triplets are important for HAP1 binding. However, changing the CGC triplets in CYC7 to CGG, increased HAP1 binding and activation, suggesting that HAP1 can recognise a CGC repeat, but less efficiently. In agreement with the earlier study by Zhang and Guarente (1994b), TA repeats were found to be important for HAP1 binding, but were found to be more essential for the imperfect HAP1 sites, such as the weaker CGC motif of CYC7. Thus, Ha et al. (1996) proposed that the sequence of the HAP1 binding sites influences the conformation of the protein to allow the activity of HAP1 to be altered.

Figure 1.1 Projection of HAP1 binding at CYC7 and CYC1 onto DNA helices (Pfiefer et al., 1987).

The DNA helices are drawn with 10.5 base pairs per helical turn. DNaseI protected sequences are shown (-251 to -229 for CYC7 and -249 to -247 for CYC1). Guanine (major groove) contacts are shown as diamonds; adenine (minor groove) contacts are shown as circles. Open symbols represent contacts on the opposite side of the helix. The orientation of the CYC7 sequence relative to the TATA box has been inverted to give maximal alignment of the binding motif.



Regulatory proteins that coordinate two or more genes, generally recognise a consensus sequence in the promoter region of the genes. The UAS1 of CYC1 and UAS of CYC7 appeared to be obvious exceptions to this, however it now seems that these binding sites are degenerate forms of the consensus sequence CGG N3 TA N CGG N3 TA (Ha et al., 1996).

# 1.3.2 Regulation of CYC1 and CYC7 in Response to Carbon Source by the Yeast CCAAT-binding Factor

CYC1 and CYC7 genes are both regulated by carbon source at the transcriptional level. CYC1 expression is repressed 10-20 fold with glucose as the carbon source (Zitomer & Lowry, 1992), and induced 50-fold when shifted from glucose to the non-fermentable carbon source, lactate (Guarente et al., 1984). CYC7 expression is induced 4- to 8-fold with a shift from glucose to lactate medium (Prezant et al., 1987).

#### 1.3.2.1 The Yeast CCAAT-binding Factor (HAP2/3/4/5 Complex)

A non-fermentable carbon source such as lactate induces transcription of the *CYC1* gene through the binding of a conserved CCAAT-binding factor to a CCAAT box with the consensus TNATTGGT (the CCAAT box is encoded on the other strand). This is located in the upstream activation site, UAS2. In *S. cerevisiae*, the CCAAT-binding factor has been shown to be a heteromeric transcription factor composed of the heme-dependent regulatory proteins, HAP2, HAP3, HAP4 and HAP5. The subunits HAP2/3/4 were identified by mutations which abolished the transcriptional activation of the *CYC1* gene from the UAS2 element, and thus suppressed growth on a non-fermentable carbon source (Olesen *et al.*, 1987; Forsburg and Guarente, 1989b). HAP5 was recently identified by using the two-hybrid screening system (McNabb *et al.*, 1995). Two other undefined proteins were also obtained from the hybrid screen and may be required for the complete CCAAT-binding factor, or may encode ancillary factors for CCAAT-binding transcriptional activation (McNabb *et al.*, 1995).

None of the subunits of the yeast CCAAT-binding factor contain any homology to other known classes of DNA-binding motifs. Therefore, the precise mechanism for subunit association is unknown and will most likely prove to be novel. HAP4 has been shown to encode the principle transcriptional activation domain (Forsburg and Guarente, 1989b; Olesen and Guarente, 1990). It contains a highly acidic domain responsible for transcriptional activation which can be functionally exchanged with the GAL4 acidic activation domain. Furthermore HAP4, unlike HAP2 and HAP3, can independently activate transcription at high levels through a lexA operator-driven promoter. The

HAP2/3/5 subunits have been shown through mobility shift experiments to each be essential for DNA binding activity *in vitro*, thus representing the first heterotrimeric transcription factor (McNabb *et al.*, 1995).

Mutational analysis of the HAP2 protein has identified an essential 60 amino acid core region required for all functions of the HAP2 subunit (Olesen and Guarente, 1990). This core region contains an 18 amino acid subunit association domain (SAD) and a DNA binding domain of 21 amino acids (Olesen and Guarente, 1990; Xing et al., 1993). Amino acid residues critical for HAP2 subunit association have been identified (Xing et al., 1994), and structural studies of this region indicate that the SAD forms an alphahelical structure which interacts with another HAP subunit (Xing et al., 1994). HAP2 has been shown to bind to the HAP3 subunit (Olesen and Guarente, 1990), and most likely also binds to the HAP5 protein, which was isolated through protein interactions with the HAP2 protein (McNabb et al., 1995). Through these studies it was suggested that HAP2 and HAP3 proteins interact through a weak interaction involving their respective DNA binding domains, and through a strong interaction between the SADs. The HAP5 subunit was suggested to be central to the complex, and acts by bringing HAP2 and HAP3 together through the binding of their SADs. This action would allow the weak DNA binding domains of HAP2 and HAP3 to come together and interact with the CCAAT box for cytochrome c activation.

The S. cerevisiae HAP2, HAP3 and HAP5 proteins appear to be highly conserved through yeast to humans. Functional mammalian equivalents (Chodosh et al., 1988) and homologs of HAP2 and HAP3 have been isolated from Schizosaccharomyces pombe (Olesen et al., 1991; Xing et al., 1993), Kluyveromyces lactis (NGuyen et al., 1995; Mulder et al., 1994), mouse (Hooft van Huijsduijnen, 1990), and rat (Maity et al., 1990). A homolog of HAP2 has been isolated from humans (Becker et al., 1991), and a HAP3 homolog (hapC) and a HAP5 homolog (hapE) have been isolated from A. nidulans (Papagiannopoulos et al., 1996a; Papagiannopoulos et al., 1996b). Studies with the rat CCAAT-binding complex (CBF) suggest that the CBF-C component might represent the mammalian HAP5 homolog (Maity et al., 1992).

A number of yeast genes required for respiration (for example *HEM1*, *COX4*, and *CYT1* genes) are activated by the yeast CCAAT-binding factor. Therefore, the HAP2/3/5 heterotrimer has been suggested to be a global yeast gene regulator (Zitomer and Lowry, 1992; Chodosh *et al.*, 1988). In addition, since only homologs of HAP2/3/5 have been obtained, it was suggested that this heterotrimer corresponds to the general transcription factor found in other species.

#### 1.3.3 Other factors affecting CYC7 expression

Minor factors affecting transcription of the yeast CYC7 gene but not the CYC1 gene are heat shock, stationary-growth phase and cAMP levels (Pillar & Bradshaw, 1991). During stationary-growth phase, transcription of CYC7 is strongly induced (40-70 fold), whereas transcription of CYC1 is greatly decreased. Heat shock increases the level of CYC7 transcription by 4-6 fold aerobically and 30 fold anaerobically. Low cAMP levels (which have also been shown to induce several heat shock proteins) also induce the CYC7 gene. Therefore the response of the CYC7 gene to these environmental factors may possibly reflect a unique role for this gene.

The heme-dependent regulatory protein HAP1 which activates transcription of CYC7 via the UAS is unlikely to be involved in both heat shock and stationary induction of the CYC7 gene. HAP1 only activates CYC7 in aerobic conditions and therefore cannot be involved in heat shock induction which occurs both aerobically and anaerobically. Stationary growth phase is probably not heme mediated, because CYC1 which is activated by HAP1 is repressed during these conditions.

# 1.4 REGULATION OF THE KLUYVEROMYCES LACTIS CYTOCHROME C GENE (KICYCI)

The aerobic respiratory yeast *K. lactis*, has one cytochrome *c* gene, *KlCYC1*, which produces two transcripts of different sizes (Freire-Picos *et al.*, 1995). This is in contrast to the fermentative yeast *S. cerevisiae* which produces a single transcript for each of *CYC1* and *CYC7*. Studies with this aerobic yeast may help to provide further understanding into the regulation of respiration in aerobic organisms, such as *A. nidulans*.

Consensus sequences for the recognition of the *S. cerevisiae* regulatory HAP2/3/4/5 complex and for the HAP1 transcription factor have been found in the upstream region of the *KlCYC1* gene (Freire-Picos *et al.*, 1993).

The *KlCYC1* gene was found to be transcriptionally activated by oxygen through the control of heme in the same way as found for the *S. cerevisiae CYC1* gene (Freire-Picos *et al.*, 1995). In addition, heterologous expression of the *KlCYC1* gene in a *hap1* mutant strain of *S. cerevisiae* decreased the level of cytochrome *c* expression in the mutant strain compared to wild type (Freire-Picos *et al.*, 1995). These results support the hypothesis that *KlCYC1* expression is regulated by the presence of oxygen through a factor similar to HAP1.

Functional homologs of the *S. cerevisiae HAP2* and *HAP3* genes have been isolated and cloned from *K. lactis* (NGuyen *et al.*, 1995; Mulder *et al.*, 1994). However, the significance of putative HAP2/3 regulatory sites remains unclear due to apparently conflicting results from two groups.

One group showed from the heterologous expression of *KlCYC1* in a *S. cerevisiae hap2* mutant, that the *HAP2* gene was essential for the expression of the *K. lactis* cytochrome *c* gene as it is for that of *S. cerevisiae* (Freire-Picos *et al.*, 1995). In addition the *K. lactis* strain Y1140 was found to be subject to glucose repression (Freire-Picos *et al.*, 1995), thus supporting the function of the HAP2/3/4/5 complex in *KlCYC1* expression.

Another group's results showed that similarly to *S. cerevisiae*, expression of the *KICYCI* gene is significantly induced on a non-fermentable carbon source (Mulder *et al.*, 1995). However, in contrast to *S. cerevisiae*, they found that the *KICYCI* gene was not glucose repressed, because expression levels of *KICYCI* were not affected with the addition of glucose to a non-fermentable carbon source. However, glucose repression has been indicated to be a strain-dependent feature in *K. lactis*. The *K. lactis* nuclear genes *KIQCR7* and *KIQCR8* (encoding subunits VII and VIII respectively of the mitochondrial bc<sub>1</sub>--complex) were also not repressed by glucose (Mulder *et al.*, 1995). These results are in agreement with observations made in 1966 by De Deken, who observed that various *K. lactis* respiratory enzymes were not affected by glucose repression, but contrasts to the findings above by Freire-Picos *et al.* (1995). However, both groups employed different strains of *K. lactis*, which may explain the different results obtained.

Further work by Mulder et al. (1994) and NGuyen et al. (1995), indicated that the K. lactis HAP2 and HAP3 genes were not essential for expression of the mitochondrial respiratory system. Both KIHAP2 and KIHAP3 genes were inactivated in K. lactis, but in contrast to the situation in S. cerevisiae, these hap null mutants did not alter the ability of K. lactis to respire and grow on non-fermentable carbon sources. Interestingly, the KIHAP2 gene was able to be correctly expressed when placed in S. cerevisiae and to complement a HAP2 deficiency (NGuyen et al., 1995). Similar results were obtained for the cytochrome c gene from the starch fermenting yeast Schwanniomyces occidentalis (CYCI<sub>o</sub>), which was not repressed by glucose in native S. occidentalis cells, but was glucose repressed when placed in S. cerevisiae strains (Amegadzie et al., 1990). Both these results tend to suggest that the HAP2/3/4/5 complex has no role in glucose repression in either K. lactis, or S. occidentalis cytochrome c genes, but will regulate these genes in response to carbon source when placed in a S. cerevisiae background.

In summary, the *KICYC1* gene appears to be transcriptionally activated in response to oxygen through the binding of a HAP1-like transcription factor to its upstream regulatory region. However, the role of the HAP2/3/4/5 complex in the regulation of the *KICYC1* is more unclear. The disruptions of the *KIHAP2* and *KIHAP3* genes performed in *K. lactis* by Mulder *et al.* (1994) and NGuyen *et al.* (1995), are more meaningful than the *KIHAP2* disruption performed in *S. cerevisae* by Freire-Picos *et al.* (1995) because the gene disruptions were carried out under the true environmental influences of the organism. Therefore these results suggest that either the HAP2/3/4/5 complex has a different function in *K. lactis*, or perhaps that additional controls or factors (specific to respiratory or fermentative yeasts) may be required to adjust the general function of the HAP2/3/4/5 complex.

# 1.5 REGULATION OF THE ASPERGILLUS NIDULANS CYTOCHROME C GENE (cycA)

Like most lower eukaroytes, A. nidulans contains only one copy of the cytochrome c gene (cycA) (Raitt et al., 1994). The predicted nucleotide sequence has approximately 72% identity to the nucleotide sequence of the S. cerevisiae CYC1 gene (Raitt et al, 1994). Like the S. cerevisiae cytochrome c genes, expression of the cycA gene is transcriptionally regulated by oxygen availability, and is induced by heat shock conditions (Raitt, 1992). However, in contrast to S. cerevisiae, cycA expression is not affected by glucose repression. The cycA gene may be transcriptionally activated by oxygen in a similar manner to that established for the cytochrome c genes of S. cerevisiae (see Section 1.3), but further experimental evidence is required to determine this.

## 1.4.1 Regulation of the cycA gene in response to Environmental Parameters

Transcription of the *cycA* gene is activated by oxygen with a 10 fold increase observed during the transition from anaerobic to aerobic conditions (Raitt *et al.*, 1994). Similar expression levels of the *S. cerevisiae CYC7* gene occur in aerobic conditions (Prezant *et al.*, 1987).

CycA transcript levels remain constant when A. nidulans is grown in the presence of fermentable (glucose) or non-fermentable (galactose) carbon sources (Raitt et al., 1994). Therefore the cycA gene is not glucose repressed as observed for the S. cerevisiae cytochrome c genes. However, it may be that glucose repression in A. nidulans is a strain-dependent feature, as found for K. lactis. However, the lack of glucose repression is not surprising since A. nidulans is an obligate aerobe, where reduced activity of the

respiratory chain may compromise the organism's ability to generate ATP, and to support cell growth.

Under heat shock conditions, a 3-4 fold induction of the *cycA* transcript occurs (Raitt *et al.*, 1994). A similar response is observed for the *S. cerevisiae CYC7* gene (Pillar and Bradshaw, 1991).

#### 1.4.2 Analysis of the 5' Region of the cycA gene

#### 1.4.2.1 Putative HAP1 Binding Site

A putative HAP1 binding site was located 382-400 nucleotides upstream of the published translation initiation site, and was found to contain the greatest homology with the CYC7 HAP1 binding site (Raitt, 1992). The sequences responsible for HAP1 mediated oxygen regulation of the yeast CYC1 and CYC7 genes, are shown aligned to the cycA putative HAP1 binding sequence (Figure 1.2; Raitt, 1992). Ten out of eleven nucleotides which contact HAP1 in the CYC7 promoter are also conserved in the cycA promoter region. Due to these sequence similarities, it was proposed that a HAP1-type gene product may be involved in the oxygen induced transcriptional activation of the A. nidulans cycA gene.

#### 1.4.2.2 Putative Upstream Opening Reading Frame (uORF)

At 120 nucleotides before the translational start point, an upstream open reading frame was found which could encode a 10 amino acid polypeptide including five consecutive serines (Raitt et al., 1994). This uORF may be involved in a translational regulatory mechanism, similar to that observed in some genes of *S. cerevisiae* such as *GCN4*. The *GCN4* gene is a transcriptional activator of many genes involved in the biosynthesis of ten amino acids. It contains four ORFs in its mRNA leader sequence which are involved in translational control. In response to amino acid availability, these uORFs can control ribosome flow so initiation occurs at the correct codon (Abastado et al., 1990).

#### 1.5 AIMS AND OBJECTIVES

The aim of this study was to investigate the role of promoter sequences, in particular, the putative HAP1 binding site, in the regulation of the A. nidulans cytochrome c gene.

In vitro methods for promoter analysis include gel mobility shift experiments and DNaseI footprinting. However, promoter-reporter vectors present a powerful in vivo method to

# Figure 1.2 Comparison of HAP1 binding sites in S. cerevisiae and A. nidulans

Nucleotides shown in blue are directly involved in the binding of the HAP1 protein. Red nucleotides represent those conserved between the *CYC7* and the *cycA* promoters. The numbering of the nucleotide is relative to the ATG initiation codon.

		-3	20 I	-330 I	-340 I	)
S. Cerevisiae	5 '	GTCAT	CGTCCGTA	AACCCCG	GCCAA	3
CYCI	3 '	CAGTA	GCAGGCAT	TTGGGGC	CGGTT	5
		-250 I	-24 I	0	-230 I	
S. Cerevisiae	5 '	AGCTA	ATAGCGAT	AATAGCG	AGGGC	3
CYC7	3 ′	TCGAT	TATCGCTA	TTATCGC	TCCCG	5
		-400 T	-390 T	-38	3 O	
A. niclulans	5 '	GGCTA	AGGGCGGT	GACAGCT.	ACTCT	3
eveA	3 '	CCGAT	TCCCGCCA	CTGTC GA	TGAGA	5

investigate the functional role of promoter elements involved in gene regulation. This type of analysis, unlike the *in vitro* methods mentioned above, allows gene expression to be investigated under the true environmental influences of the organism. I therefore chose to use this method to study the putative HAP1 motif in the *cycA* gene.

For the filamentous fungus A. nidulans, integration vectors have been constructed containing the lacZ reporter gene of Escherichia coli (van Gorcom et al, 1986) to allow investigation of promoter sequences involved in gene regulation. Transcriptional or translational gene fusions can be obtained, and quantitative analysis of expression signals is attainable, since these vectors preferentially integrate at a unique place in the genome (the argB locus), in one copy (van Gorcom et al, 1986). Streatfield et al (1991), have successfully used this type of lacZ gene fusion vector to carry out a functional analysis of the expression of the 3`-phosphoglycerate kinase (pgk) gene in A. nidulans.

In this project my initial aim was to prepare vectors which contained the *Escherichia coli lacZ* gene translationally fused to either the entire *cycA* promoter region, or the entire promoter region with deletion of the putative HAP1 binding site. Following transformation of these constructs into *A. nidulans*, and confirmation by Southern blotting that they had integrated into the *argB* locus, the expression of the *lacZ* reporter would be assessed. This assessment would involve β-galactosidase assays and Northern blots to measure transcription and translation of the *lacZ* reporter under different environmental conditions. This study should allow us to determine the significance of the putative HAP1 binding site for the regulation of the *A. nidulans* cytochrome *c* gene. In addition I hoped to confirm that observed differences in mRNA levels under low and high oxygen levels are due to transcriptional regulation and not simply due to differences in mRNA half-life. Genuine transcriptional regulation should be reflected in the promoter reporter fusions whilst differences in mRNA half-life would not. Information regarding possible translational regulation by the uORF should also be elucidated from this study.

# CHAPTER 2 MATERIALS AND METHODS

## 2.1 FUNGAL AND BACTERIAL STRAINS, LAMBDA CLONES AND PLASMIDS

These are listed in Table 2.1.

#### 2.2 MEDIA

All media was prepared with Milli-Q water and sterilised by autoclaving at 121°C for 15 minutes. Both solid and liquid media was cooled to approximately 50°C before the addition of media supplements.

- **2.2.1** Luria Broth (LB) Medium (g/L): tryptone, 10.0; yeast extract, 5.0; NaCl, 5.0; 1% D-glucose. The pH was adjusted to 7.0 before autoclaving. For solid media, agar was added to 1.5%. When required, ampicillin, isopropylthio- $\beta$ -D-galactoside (IPTG) and 5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside in dimethylformamide (X-gal) were added at a final concentration of 100  $\mu$ g/ml, 30  $\mu$ g/ml and 60  $\mu$ g/ml respectively.
- **2.2.3 MYG Media** (g/L): malt extract, 5.0; yeast extract, 2.5; 2% glucose; CuSO<sub>4.5</sub>H<sub>2</sub>O, trace; agar, 20.0.
- 2.2.4 SOC Medium: 2% (w:v) Tryptone, 0.5% (w:v) Yeast Extract, 10 mM NaCl, 10 mM MgCl<sub>2</sub>, 10 mM MgSO<sub>4</sub>, 20 mM Glucose.
- **2.2.5** NZCYM (g/L): NZ amine (type-A hydrolysate of casein), 10.0; casamino acids, 1.0; NaCl, 5.0; yeast extract, 5.0; MgSO4.7H<sub>2</sub>O, 2.0. The pH was adjusted to pH 7.5 before autoclaving.
- **2.2.6 TB Top Agarose** (g/L): tryptone, 10.0; NaCl, 5.0; agarose, 8.0. After autoclaving, the media was cooled to 45-50°C and supplemented with 1 M MgSO4.7H<sub>2</sub>O.

#### 2.3 COMMON BUFFERS AND SOLUTIONS

2.3.1 1 x TAE Buffer: 40 mM Tris-acetate, 11.4 ml/L glacial acetic acid, and 1 mM EDTA; pH 8.5.

Table 2.1 Strains,  $\lambda$  clones and plasmids

Strain, Plasmid or I Clone	Relevant Characteristics	Source or Reference
Fungal Cultures		
A.nidulans		
A18	(R153) wA3, pyroA4	Waldron and Roberts (1973)
Bacterial Cultures		
E. coli		
XL-1	supE44 hsdR17 recA1 endA1 gyrA96 thi-1relA1 lac <sup>-</sup> F <sup>-</sup> [pro AB <sup>+</sup> lac <sup>IQ</sup> lacZDM15 Tn10(tet <sup>I</sup> )]	Bullock et al. (1987)
DH1	F' supE44 recA1 endA1 gyrA96 thi1	Hanahan (1983)
LE392	relA1 hsdR171 <sup>-</sup> supE44 supF58 hsdR514 galK2 galT22 metB1 trpR55 lacY1	Borck et al. (1976)
<u>Plasmids</u>		
R40	0.9 kb <i>Hin</i> dIII- <i>Xho</i> I fragment of <i>A. nidulans cycA</i> gene contained in pICI9R	Raitt (1992)
R41	2.3 kb PstI-HindIII fragment of A. nidulans cycA gene contained in pBS	Raitt (1992)
R109	(pAN923-42B) lacZ Ap <sup>r</sup> t <sub>trp</sub> C	Van Gorcom et al. (1986)
R116	(pAN5-d1) gpdA lacZ Ap <sup>r t</sup> trpC argB-BglII	Punt et al. (1990)
R117	(pAN923-42B $_{Bg\Pi I}$ ) lacZ Ap <sup>r t</sup> trpC argB- $Bg\Pi I$	Punt et al. (1990)
R120	0.68 kb BamHI fragment of A. nidulans cycA gene contained in R117 (wrong orientation)	this study
R121	0.68 kb BamHI fragment of A. nidulans cycA gene contained in pUC18	this study
R122	0.68 kb BamHI fragment of A. nidulans cycA gene contained in R117 (correct orientation)	this study
R127	2.1 kb <i>Eco</i> R1 fragment of <i>A. nidulans</i> cycA gene contained in pUC18	this study
pUC18	Ap <sup>r</sup> lacZ'	Yanish-Perron et al. (1995)
$\lambda$ clones		
λGEM-11	The arms are derived from EMBL3A and the stuffer fragment is derived from λ2001	Frischauf <i>et al.</i> (1983); Karn (1984)
λLM5	λGEM-11 clone containing cycA from A18	this study
λLM9	λGEM-11 clone containing cycA from	this study
λLM19	A18 λGEM-11 clone containing cycA from A18	this study

- 2.3.2 1 x TBE Buffer: 89mM Tris HCl, 89 mM boric acid, 2.5 mM Na<sub>2</sub>EDTA.
- 2.3.3 TE Buffer (10/1): 10 mM Tris-HCl, and 1 mM Na<sub>2</sub>EDTA.
- 2.3.4 TE Buffer (10/0.1): 10 mM Tris-HCl and 0.1 mM Na<sub>2</sub>EDTA.
- 2.3.5 10 x Sequencing TBE Buffer: (g/l): Tris, 162; Na<sub>2</sub>EDTA, 9.5; and boric acid, 27.5.
- **2.3.6** TES (10/1/100): 10 mM Tris-HCl (pH 8.0), 1 mM Na<sub>2</sub>EDTA (pH 8.0), and 100 mM NaCl.
- **2.3.7 STET Buffer:** 8% sucrose, 5% Triton X-100, 50 mM Na<sub>2</sub>EDTA (pH 8.0), and 50 mM Tris-HCl (pH 8.0).
- **2.3.8 SM Buffer:** (g/L): NaCl, 5.8; MgSO47H<sub>2</sub>O, 2.0; 1M Tris-HCl (pH 7.5), 50 ml; and 2% gelatin, 5 ml.
- **2.3.9 Hybridisation Solution:** (per litre): 150 ml of 20 x SSC (Section 2.3.12), 20 ml of 50 x Denhardt's Solution (Section 2.3.10), 30 ml of 10% SDS, 5 ml of salmon sperm DNA (at 50 μg/ml).
- 2.3.10 50 x Denhardt's Solution: (g/L): ficoll, 10.0; polyvinylpyrolidone, 10.0; bovine serum albumin, 10.0. Filter sterilised.
- 2.3.11 Tris-Equilibrated Phenol was prepared by melting solid phenol at 50°C followed by the addition of 0.1 M Tris-HCl (pH 8.0) and thorough mixing. After 15-30 minutes, the buffer was decanted and the procedure repeated twice. Hydroxyquinoline was added to a final concentration of 0.1% (w/v) and the equilibrated phenol stored under 100 mM Tris-HCl (pH 8.0) at 4°C.
- 2.3.12 20 x SSC: 3 M NaCl and 0.2 M tri-sodium citrate.
- **2.3.13 10 x Gel Loading Buffer:** 2 M Urea, 50% glycerol, 50 mM Tris acetate, 0.4% Bromophenol Blue, 0.4% Xylene cyanol.
- **2.3.14** Standard DIG Buffer: 5 x SSC, 1.0% (w/v) blocking reagent stock solution (Section 2.3.18), 0.1% N-lauroylsarcosine, and 0.02% SDS.

- 2.3.15 DIG Buffer 1: 100 mM maleic acid, and 150 mM NaCl: pH 7.5.
- **2.3.16 DIG Buffer 2:** 1% blocking reagent stock soluton (Section 2.3.18) diluted in Buffer 1 (Section 2.3.15).
- 2.3.17 DIG Buffer 3: 100 mM Tris-HCl (pH 9.5), and 100 mM NaCl.
- **2.3.18** Blocking Reagent stock solution: 10% (w/v) blocking reagent (Boehringer Mannheim DIG kit) diluted in DIG buffer 1 (Section 2.3.15).
- **2.3.19** Alkaline Lysis Solution I: 50 mM glucose, 25 mM Tris HCl pH 8.0, and 10 mM EDTA: pH 8.0.
- 2.3.20 Alkaline Lysis Solution II: 0.2 M NaOH, and 1% SDS.
- **2.3.21** Alkaline Lysis Solution III: 60ml 5 M potassium acetate, 11.5ml glacial acetic acid, and 28.5ml H<sub>2</sub>O.

#### 2.4 GROWTH AND MAINTENANCE OF CULTURES

Bacterial cultures were grown at 37°C on LB agar plates, supplemented with antibiotic when appropriate, and sealed with parafilm for storage at 4°C. Fungal cultures were grown on MYG plates, either at 37°C overnight or at 30°C for 2 days, and sealed with parafilm for storage at 4°C.

#### 2.5 DNA PREPARATIONS

#### 2.5.1 Rapid Boiling Plasmid Preparation

This method for preparing plasmid DNA was performed as described by Holmes and Quigley, (1981).

#### 2.5.2 Small Scale Alkaline Lysis Plasmid Preparation

The pAN923-42B vector (Table 2.1) was prepared by the small scale alkaline lysis method followed by phenol-chloroform extraction and ethanol precipitation (Section 2.5.7).

This method was based on that of Sambrook *et al.*, (1989). 3 ml of LB was inoculated with a single bacterial colony and incubated overnight at 37°C with vigorous shaking. 1.5 ml of the culture was pelleted by centrifugation at 12000 rpm (in a microcentrifuge) for 30 seconds. After removing the supernatant the pellet was resuspended in 100 µl of solution I (Section 2.3.19). 200 µl of freshly prepared solution II (Section 2.3.20) was added and mixed by inverting the tube rapidly five times and placing on ice. 150 µl of solution III (Section 2.3.21) was added, followed by gentle vortexing, and incubated on ice for 5 minutes. After centrifugation at 12000 rpm (in a microcentrifuge) for 5 minutes, the supernatant was transferred to a fresh tube. An equal volume of phenol:chloroform was added, the tube vortexed and centrifuged for 2 minutes. The aqueous layer was transferred to a fresh tube, and the DNA was precipitated with 2 volumes of 100% ethanol. The DNA was pelleted by centrifugation for 5 minutes at 12,000 rpm (in a microcentrifuge) and then dried under vacuum before resuspension in TE buffer (10/0.1) or H<sub>2</sub>O.

#### 2.5.3 Large Scale Alkaline Lysis Plasmid Preparation

The modified pAN923-42B $_{BgIII}$  vector (Table 2.1) was prepared by the large scale alkaline lysis method and purified by using a CsCl gradient (Section 2.5.4).

This method was based on that of Ish-Horowicz and Burke (1981). 250 ml of LB was inoculated with a 1/100 dilution from a 3 ml overnight culture of E. coli, and incubated overnight at 37°C on a shaking platform (at 250 rpm). Cells were harvested by centrifugation at 8000 rpm (centrifugation was carried out in a GSA rotor throughout) for 10 minutes, then washed in 100 ml of TE (10/0.1) by resuspension and pelleted by centrifugation at 8000 rpm for 10 minutes. The cells were resuspended in 30 ml of solution I (Section 2.3.19), 3 ml of lysozyme (at 50 mg/ml) was added and the tube incubated at room temperature for 10 minutes. 60 ml of solution II (Section 2.3.20) was added, mixed by inversion and the tube placed on ice for 10 minutes. 45 ml of solution III (Section 2.3.21) was added, mixed by inversion followed by incubation on ice for 10 minutes. After centrifugation at 8000 rpm for 10 minutes, the supernatant was transferred to a fresh tube and the DNA precipitated by the addition of 0.6 volumes of cold isopropanol followed by incubation at room temperature for 20 minutes. DNA was pelleted by centrifugation at 12000 rpm for 10 minutes, washed with 75 ml of 70% ethanol, and air dried before resuspension in 3.5 ml of TE buffer (10/0.1) with heating for 10 minutes at 65°C.

## 2.5.4 Purification of Plasmid DNA by Cesium Chloride-Ethidium Bromide Gradient.

CsCl was added to the plasmid DNA solution in a ratio of 1.05 g/ml, and ethidium bromide (at 10 mg/ml) was added in a ratio of 75 µl/ml. The mixture was vortexed and put at 4°C overnight. After this time, the solution was centrifuged at 12000 rpm in a SS34 rotor for 10 minutes to remove precipitated material, and the refractive index adjusted to between n=1.3860-1.3920 by the addition of either TE buffer or CsCl. Plasmid DNA was purified by centrifugation at 55000 rpm in a Sorvall combi TV865 rotor for 5 hours. The plasmid band was removed with an 18 gauge hypodermic needle and syringe under long wavelength UV light. Ethidium bromide was removed from the DNA sample by repeated extraction with equal volumes of SSC saturated isopropanol (prepared by stirring equal volumes of 20 x SSC and isopropanol for several hours). The DNA was dialysed at 4°C against TES (10/1/100) (Section 2.3.6) with 4 changes of buffer and stored at -20°C. DNA purity and concentration were calculated by spectroscopy as in Section 2.6.4.2.

#### 2.5.5 Preparation of DNA from Fungal Cultures

This method for isolating DNA from fungi was based on that of Raeder and Broda (1985). Mycelium was grown on MYG plates overlaid with sterile cellophane disks. 50 μl of spore suspension (Section 2.11.1) was spread on to the cellophane disks and after incubation for 20 hours at 37°C mycelium was scraped off and freeze dried. Approximately 400 mg (dry weight) of mycelium was ground to a fine powder with liquid nitrogen, 4 ml of extraction buffer (22 mM Tris pH 8.5, 250 mM NaCl, 25 mM EDTA, 0.5% SDS) was added and the mixture vortexed. At this stage 2.8 ml of phenol (equilibrated with extraction buffer) and 1.2 ml of chloroform was added, followed by centrifugation at 12000 rpm, (centrifugations were performed in a GSA rotor at 4°C through out) for 40 minutes. Supernatant was transferred to a fresh tube and the sample was re-extracted with phenol and chloroform as before, followed by centrifugation at 12000 rpm for 15 minutes. To remove RNA, 100µg/ml of RNase was added to the supernatant and incubated at 37°C for 30 minutes. An equal volume of phenol:chloroform was added and the tube centrifuged for 15 minutes at 12000 rpm. Chloroform was added to the aqueous phase and centrifuged for 20 minutes at 12000 rpm. DNA was precipitated with 0.54 volumes of isopropanol and centrifuged for 5 minutes at 12000 rpm. The DNA pellet was washed with 70% ethanol, dried and resuspended in TE buffer (10/0.1).

#### 2.5.6 Preparation of Lambda Phage DNA by Liquid Lysate

λ phage DNA was prepared using a modification of the liquid lysate method from Current Protocols in Molecular Biology (1994). 3 ml of LB supplemented with 0.2% maltose and 10 mM MgSO4.7H2O, was inoculated with a single bacterial colony of LE 392 and incubated overnight at 37°C. 100 µl of overnight bacterial culture was combined with 100 μl of titred diluted phage to give 10<sup>6</sup> or 10<sup>7</sup> phage/100 μl. The phage mixture was transferred to 50 ml of NZCYM and shaken at 300 rpm at 37°C until lysis occurred, indicated by the culture clearing. 200 µl of chloroform was added upon clearing to lyse any remaining cells and the culture shaken at 37°C for a few minutes. The culture was then immediately harvested by centrifugation at 10000 rpm (in a GSA rotor) for 10 minutes. The lysate was transferred into a fresh tube and stored at 4°C. RNase and DNase was added to the liquid lysate to give a final concentration of 10 µg/ml and incubated at 37°C for 30 minutes. Phage were precipitated by the addition of NaCl (to 0.5 M), and PEG 6000 (to 10% (w/v). Following incubation on ice for 1 hour, the phage were pelleted at 5000 rpm (in a GSA rotor) for 10 minutes at 4°C, and the supernatant discarded. The pellet was resuspended in 1 ml of SM buffer (Section 2 3.8) and the solution transferred to an eppendorf tube. Remaining bacterial debris was pelleted by centrifugation for 10 minutes at 12000 rpm, the supernatant transferred to a fresh tube and proteinase K added to a final concentration of 0.1 mg/ml. After incubation for 30 minutes at 37°C, the phage suspension was extracted with phenol:chloroform, vortexed for 20 minutes on an eppendorf platform, and centrifuged for 5 minutes at 10000 rpm (in a microcentrifuge). Phenol:chloroform extractions were repeated, then an equal volume of chloroform was added, vortexed for 5 minutes, and centrifuged for 5 minutes as above. The upper aqueous phase was taken and DNA precipitated by the addition of 2 volumes of ethanol and 0.3 M NH4Ac. DNA was pelleted by centrifugation for 10 minutes, washed with 70% ethanol, vacuum dried and resuspended in TE buffer (10/0.1).

## 2.5.7 Purification of DNA by Phenol-Chloroform Extraction and Ethanol Precipitation

To the DNA sample an equal volume of Tris-equilibrated phenol:chloroform (1:1) was added, vortexed, and centrifuged at 13000 rpm for 5 minutes. The aqueous phase was transferred to a fresh tube, and an equal volume of chloroform added. The mixture was vortexed and then centrifuged for 5 minutes and the aqueous phase transferred to a fresh tube. To ethanol precipitate the DNA, 1/10 volume of 0.3M sodium acetate plus 3 volumes of 95% ethanol was added to the aqueous phase and mixed. The DNA was pelleted by centrifugation for 15 minutes. The ethanol was poured off, and the pellet

washed with 70% ethanol by centrifugation for 3 minutes, dried under vacuum for 5 minutes and resuspended in TE buffer (10/0.1) or H<sub>2</sub>O.

#### 2.6 STANDARD PROCEDURES

#### 2.6.1 Restriction Enzyme Digests

DNA digests were carried out with the appropriate restriction enzyme buffer at 37°C, for 1.5-3 hours, in a total volume at least 10 times the combined volume of DNA and enzyme. When possible, commercial buffers specifically matching the restriction enzyme were used. RNA was removed by adding RNase to a final concentration of 0.5  $\mu$ g/ $\mu$ l and incubating for a further 10 minutes at 37°C. Digests were checked for completion by electrophoresis on a 1-2.5% agarose mini-gel.

#### 2.6.2 Agarose-gel Electrophoresis of DNA

DNA fragments were size fractionated by electrophoresis through a gel ranging from 0.8-2.5% agarose (depending on the expected fragment sizes), in TAE or TBE buffer at 80-100 V. Large preparative gels were usually electrophoresised overnight at 25-30 V. Agarose gels were stained in ethidium bromide, 10 minutes for mini-gels or 30 minutes for preparative gels, and then destained in Milli-Q water, observed under short wave UV light and photographed with polaroid 667 film.

#### 2.6.3 Determination of Molecular Weights

DNA fragments were sized by running the DNA sample alongside any of the following standards:

- (i) HindIII/EcoRI double digest of lambda DNA
- (ii) 1 kb DNA ladder (BRL)
- (iii) HinfI digest of pBR322

Relative mobilities of DNA fragments were measured, and the molecular weights determined using the GEL 1.01 software program (1993 Jean-Michel Lacroix).

#### 2.6.4 Determination of DNA Concentration

#### 2.6.4.1 Quantification of DNA Concentration by Electrophoresis

A series of Lambda DNA or pBR322 DNA concentration standards were run on a 1.0-2.5% agarose minigel alongside the DNA sample. DNA sample concentration was estimated by comparing the intensity of fluorescence to that of the DNA concentration standards.

#### 2.6.4.2 Spectrophotometric Determination of DNA Concentration and Purity

This method was used to the estimate the DNA concentration of DNA solutions of high concentration and purity. The DNA sample was appropriately diluted and the absorbance measured at both 260 nm and 280 nm. The DNA concentration was calculated on the assumption that an absorbance of 1.0 at 260 nm is equivalent to 50  $\mu$ g/ml. Purity was indicated by the 260 nm/280 nm ratio, where pure DNA has an OD260/OD280 ratio of 1.8-2.0 (Current Protocols in Molecular Biology, 1989; Glasel, 1995).

### 2.7 AMPLIFICATION OF DNA BY THE POLYMERASE CHAIN REACTION (PCR)

PCR reactions were set up on ice using a cocktail which contained all common reagents used for n+1 PCR reactions. Uncommon reagents were aliquoted out separately. The PCR reactions were carried out in either 25 μl or 50 μl volumes in 0.2 ml strip tubes (Biotek). The final concentrations of each component in 1 reaction were, 1 x Tfl buffer (Boehringer Mannheim), 1.5 mM MgCl<sub>2</sub>, 1.25 mM dNTPs, 1 unit/25μl of Tfl polymerase (Boehringer Mannheim), 2 μM forward primer, 2 μM reverse primer, 0.1 ng to 1 ng of template DNA, and MilliQ water up to the total reaction volume. A negative control containing water instead of DNA template was included in each PCR run. After mixing, the PCR reaction tubes were placed in a thermal cycler (Corbett FTS-960) preheated to 95°C.

For PCR amplification of the A. nidulans cycA promoter region from cloned genomic DNA (Raitt et al. (1994), published version), the following PCR cycle conditions were used:

Initial Step	95°C	2.5 minutes
Denaturation	95°C	1 minute
Annealing	60°C	1 minute x 30
Elongation	72°C	2 minutes
Elongation	72°C	5 minutes
Cool	4°C	1 minute

For repeated PCR amplification of the 596bp and 298bp A. nidulans cDNA products, the following optimised PCR cycle conditions were used:

Initial Step	95°C	2.5 minutes
Denaturation	95°C	30 seconds
Annealing	65°C	30 seconds x 25
Elongation	72°C	1 minute
Elongation	72°C	5 minutes
Cool	4°C	1 minute

On completion of PCR, the reactions were checked on a 2.5% agarose minigel. Primers used in the PCR reactions are listed in Table 2.2.

#### 2.8 DNA CLONING PROCEDURES

Fragments to be cloned were gel purified (Section 2.8.1), ligated into the appropriate CAP-treated vector (Sections 2.8.2 and 2.8.3), and transformed into a suitable *E. coli* host strain (Section 2.8.4, Table 2.1). Transformants were screened for recombinants by one of the following methods. When blue/white selection of recombinants was available (eg. when using pUC18 vector), the preferred method involved restriction digest analysis (Section 2.6.1) of plasmid DNA obtained by the rapid boiling method (Section 2.5.1). When there was no blue/white selection and the colony numbers were large, colonies were screened either by colony hybridisation (Section 2.9) or PCR (Section 2.7) using primers designed to specifically detect the presence of insert (refer to Table 2.2).

Table 2.2 PCR and Sequencing Primers

Primer	Size	Tm°C*	Sequence (5' to 3')	Source
RB1	24 mer	72	gTggATCCTACCTCTA TTggTCA	Saunders (1993)
RB2	24 mer	74	gTggATCCTCgAAgAgC TTAgCAC	Saunders (1993)
RB3	26 mer	76	gTgTCgACTCTCCTggT AAgTAgTTT	Saunders (1993)
RB4	25 mer	72	gTgTCgACAAgggTTATg TAAAgAg	Saunders (1993)
LG1	24 mer	72	gTAgATCTCgTCTAATC gCgAACg	this study
LG2	25 mer	74	CCAATCAgAggTAggAT GATTCAgC	this study
LG3	24 mer	76	gACgCTggTTACTTggAg CTCTgC	this study
LG4	24 mer	70	AgTgCAgAgCTCCAAgT AAccAgC	this study
LM5	23 mer	76	CgAggTgTTCCCTCAgC ACgAgC	this study
LM6	25 mer	76	CAggAATAggATTgCTC gTgCTgAg	this study
LM7	25 mer	72	ggTAgTTAACgACTTTA CCATTgCg	this study
CYC8	23 mer	66	gTggTgAAgCTAAATgT TgCAAg	this study
pUC/M13 Forward	22 mer	70	gCCAgggTTTTCCCAgT CACgA	Perkin Elmer
pUC/M13 Reverse	24 mer	70	gAgCggATAACAATTTC ACACAgg	Perkin Elmer

<sup>\*</sup> Calculated as  $Tm^{\circ}C=2(A+T)+4(G+C)$  from Itakura et al., (1984).

#### 2.8.1 Gel Purification of DNA

DNA fragments to be purified from either a PCR reaction or restriction enzyme digest, were run on a 1% sea-plaque low melting point agarose gel in TAE buffer. After staining in ethidium bromide, the DNA fragments were visualised under long wave UV light, and the appropriate band cut out with a sterile scalpel blade. The DNA was extracted from the agarose using the following commercial kits according to the manufacturers instructions:

- (i) Gibco BRL Glassmax DNA Isolation Spin Cartridge System (Life Technologies Ltd).
- (ii) Bio 101 Geneclean Kit (Bio 101).
- (iii) Wizard PCR Preps DNA Purification System for Rapid Purification of DNA Fragments (Dade Diagnostics Ltd).

#### 2.8.2 CAP Treatment of DNA

Vector DNA (5-10  $\mu$ g) was digested to completion with the desired restriction enzyme, and the enzyme in-activated by heat (if applicable) or by phenol and chloroform extraction (Section 2.5.8). 1 unit of calf alkaline phosphatase (CAP) was added to the DNA and incubated for 30 minutes at 37°C. This was repeated once, then proteinase K and SDS were added to a final concentration of 50  $\mu$ g/ml and 1%, respectively followed by incubation for 1 hour at 37°C. The DNA was subsequently purified as described in Section 2.5.7.

#### 2.8.3 Ligation Reactions

Ligation reactions usually contained 1-3 times the molar ratio of insert:vector, (using a minimum of 20 ng of vector), 2.0 μl of 10 x ligation buffer, 1.0 μl of undiluted T4-DNA ligase, and Milli-Q water to a total volume of 10 μl. Ligation was carried out at room temperature for at least 1 hour and then overnight at 4°C, (or ligated at room temperature only). For ligations that did not work under these conditions, reactions were incubated at 14°C for 2 hours and than at 4°C overnight. Ligations were checked by taking a 1 μl aliquot before the addition of ligase, and 1 μl after the ligation incubation period. Both samples were run on an agarose minigel and compared. The presence of higher molecular weight bands and the disappearance of insert DNA was indicative of successful ligation. Transformation of the ligation mixture was then carried out as described in Section 2.8.4.

#### 2.8.4 Transformation of Escherichia coli

Preparation of electro-competent *E. coli* cells (Section 2.8.4.1) and electroporation of *E. coli* (Section 2.8.4.2) was based on the method by Dower *et al.* (1988). Electroporation was the preferred method for transformation of *E. coli*, as the procedure could be performed both rapidly and efficiently. However, the CaCl<sub>2</sub> transformation method (Sections 2.8.4.3 and 2.8.4.2) was advantageous to use when difficulties were experienced with arcing (due to the DNA sample having a high ionic strength) during electroporation. Dialysis could be used to overcome this problem, but it was found to be an unreliable method. In addition, large volumes of ligation reactions could be easily transformed by the CaCl<sub>2</sub> method to avoid ethanol precipitation of dilute DNA samples (particularly small PCR fragments, ranging from 190-680 bp), and possible complete loss of DNA.

#### 2.8.4.1 Preparation of Electroporation Competent E. coli Cells

One litre of LB broth was inoculated with 10 ml of an overnight culture of the desired E. coli strain (XL-1 or DH1), and grown at 37°C with vigorous shaking to mid-log phase (OD600 0.5-1.0, about 3 hours). The cells were chilled on ice for 20 minutes then harvested by centrifugation for 10 minutes at 5000 rpm in a GSA rotor. The cells were washed sequentially (by resuspension, centrifugation at 5000 rpm to pellet the cells, then draining the supernatant) in 1.0 litre of ice cold sterile water, 500 ml of water, 20 ml of ice cold 10% glycerol, then finally resuspended in 4 ml of 10% glycerol. The cells were stored at -70°C in 200  $\mu$ l aliquots.

#### 2.8.4.2 <u>Electroporation of E. coli</u>

2-5  $\mu$ l of DNA (ligation mixture or controls) was added to 50-100  $\mu$ l of electro-competent *E. coli* cells in an ice cold eppendorf and gently mixed. The DNA/cell mixture was immediately transferred to an ice cold 0.2 cm cuvette, tapped to the bottom and electroporated in a Biorad gene pulser set at 25  $\mu$ F, 2.5 kV and 200  $\Omega$ . A time constant of 4-5 msec was indicative of successful electroporation and cells were immediately resuspended in 1 ml of SOC medium (Section 2.2.4) and incubated for 1 hour at 37°C. A positive (circular plasmid DNA) and negative (water only) control were always included. Cells were spread on LB agar plates supplemented with ampicillin (100  $\mu$ g/ml) and incubated overnight at 37°C.

#### 2.8.4.3 <u>Preparation of CaCl2-Competent Cells</u>

*E. coli* cells were made competent by CaCl<sub>2</sub> treatment as described by Cohen *et al.* (1972). LB (25 ml) was inoculated 1:50 with an overnight culture of *E. coli* and grown at 37°C on a shaking platform at 225 rpm to mid log phase (A550=0.5, usually 2-4 hours). The cells were then pelleted by centrifugation (in a SS34 rotor) for 10 minutes at 5000 rpm at 4°C, resuspended in 10 ml of 60mM CaCl<sub>2</sub>, and incubated on ice for 20 minutes, pelleted again and resuspended in 250 μl of 60 mM CaCl<sub>2</sub>.

#### 2.8.4.4 CaCl<sub>2</sub> Transformation of *E. coli*

DNA (diluted with water to a volume of 50  $\mu$ l) was added to 250  $\mu$ l of CaCl<sub>2</sub> competent *E. coli* cells and incubated on ice for 30 minutes. The transformation mix was heat shocked for 2.5 minutes at 42°C, then 900  $\mu$ l of LB broth (prewarmed to 37°C) was added. The cells were shaken at 250 rpm for 1 hour at 37°C, then spread on LB plates supplemented with ampicilin (100  $\mu$ g/ml) and incubated overnight at 37°C.

### 2.9 COLONY HYBRIDISATION USING THE DIG DETECTION SYSTEM

This method was based on the Boehringer manual: The DIG System Users Guide for Filter Hybridisation (1994).

#### 2.9.1 Colony Lifts

Colonies to be screened were picked and patched onto duplicate nylon filters (Biotechnology Systems, 82 mm disks), placed in plates containing selective LB media, and incubated at 37°C for approximately 5 hours (or until the colonies were 1-2 mm in diameter). Each filter was marked asymmetrically with a needle to record its orientation on the plate. Filters were lifted off, and placed colony side up on 3 layers of blotting paper, soaked in 1.5 M NaCl, 0.5 M NaOH for 2 minutes, then transferred to blotting paper soaked in 1.5 M NaCl, 0.5 M Tris-HCl (pH 7.2), 1 mM EDTA for 5 minutes, and finally transferred to 2 x SSC for 2 minutes. Filters were air dried on blotting paper, then the DNA was fixed by UV-crosslinking for 2 minutes (DNA side up) on a transilluminator. Filters were washed in 3 x SSC to remove bacterial debris, dried and then stored at room temperature until required.

#### 2.9.2 Colony Hybridisation

Filters were placed in a small plastic container and prehybridised in 25 ml of standard DIG buffer (Section 2.3.14) for 2 hours. After prehybridisation, buffer was discarded, and 7 ml of fresh standard DIG buffer was added. Denatured DIG-labelled probe (Section 2.9.3) was added to a final concentration of 5-25 ng/ml and hybridisation in a rotating oven at 65°C overnight was carried out. Filters were washed with 2 x wash solution (2 x SSC containing 0.1% SDS) at room temperature (2 x 5 minutes), then washed with 0.1 x wash solution (0.1 x SSC containing 0.1% SDS) at 68°C (2 x 15 minutes). After washing, the DIG detection procedure (Section 2.9.4) was performed to develop the hybridisation signal.

#### 2.9.3 Preparation of DIG-Labelled Probe

DNA was labelled with Digoxigenin-11-dUTP (DIG) using the random primer method (Boehringer manual: The DIG Systems Users Guide for Filter Hybridisation, 1994). Approximately 10 ng of DNA was denaturated for 5 minutes in a boiling water bath, and immediately placed on ice to cool. The following reagents were added to the DNA template: 4 µl of high prime reaction mix (Boehringer, vial 1), 4 µl of DIG DNA labelling mix (containing 1 mM dATP, dCTP, dGTP, 0.65 mM dTTP, and 0.35 mM DIG-dUTP), and Milli-Q water to a total volume of 20 µl. The reaction was incubated at 37°C for 20 hours and stopped by the addition of 2 µl of 0.2 M EDTA. The labelled DNA was than precipitated by adding 0.1 volumes of 4 M LiCl, 3 volumes of 100% ethanol and incubating at -20°C overnight. The DNA was pelleted by centrifugation at 13000 g for 15 minutes, washed with 70% ethanol and then dried under vacuum. The probe was resuspended in 50 µl of TE buffer (10/1) and stored at -20°C.

To estimate the concentration of the DIG-labelled probe and to check the efficiency of the labelling reaction, 10-fold serial dilutions (in TE buffer) of labelled control DNA (Boehringer) and DIG-labelled probe were performed. 1 µl of each of these dilutions were spotted on to a piece of membrane (Hybond-N, Amersham) marked so each dilution could be identified. The DNA was fixed onto the membrane by UV crosslinking for 1 minute. After the detection procedure was carried out (Section 2.9.4), the concentration of the labelled probe was estimated by comparing the spot intensities of the control DNA (of known concentration) with the DIG-labelled probe DNA dilutions.

#### 2.9.4 Detection of Hybridised DIG-labelled Probe

The filters that had been hybridised to the DIG-labelled probe (Sections 2.9.2 and 2.9.3) were washed in DIG buffer 1 (Section 2.3.15 ) for 5 minutes, than in DIG buffer 2 (Section 2.3.16) for 30 minutes at room temperature with gentle shaking. After this time buffer 2 was discarded and replaced with the anti-DIG-alkaline phosphatase conjugate (antibody, Boehringer) diluted 10<sup>-4</sup> in DIG buffer 2. After incubation for 30 minutes at room temperature the antibody solution was poured off, and the filter placed in a fresh, clean container where it was washed at room temperature with DIG buffer 1 for 2 x 15 minutes. The filter was incubated for 2 minutes in DIG buffer 3 (Section 2.3.17) at room temperature then placed on a sheet of acetate. 100 µl drops of CSPD (chemiluminescent alkaline substrate) diluted 10<sup>-2</sup> in DIG buffer 3, (using a total volume of 0.5 ml/100 cm<sup>2</sup> of filter ) was spotted onto the filter. Another sheet of acetate was lowered onto the filter and the CSPD was spread evenly over the filter by gently wiping the surface of the acetate. The CSPD was left in contact with the filter for 5 minutes at room temperature and then incubated with the edges of the acetate sheets sealed for 15 minutes at 37°C. The acetate sealed filter was placed in a X-ray cassette with intensifying screens, either Fuji Medical or Kodak Scientific Imaging X-ray film, and exposed at room temperature for 20-30 minutes. After exposure the film was developed using standard procedures.

#### 2.10 DNA SEQUENCING

Initial DNA sequencing work was performed using the Sequenase Version 2.0 Kit from USB. The majority of sequencing used the Amplicycle Kit from Perkin Elmer. Both methods were based on the Sanger method of chain termination (Sanger *et al.*, 1977). The Amplicycle system also utilises the Polymerase Chain Reaction (PCR), enabling small amounts of template DNA to be labelled quickly and efficiently, hence this kit was ideal for directly sequencing small quantities of PCR products.

#### 2.10.1 Sequenase Version 2.0 Sequencing Protocol

To denature the template, 5-8 μg of template DNA was incubated with, 0.2 M NaOH and 0.2 mM EDTA for 30 minutes at 37°C. 0.1 volumes of 3M sodium acetate was added to neutralise the DNA and mixed well. DNA was precipitated by adding 2-4 volumes of 95% ethanol and was pelleted by centrifugation for 10 minutes. The DNA pellet was washed in 2 volumes of 70% ethanol, dried under vacuum and resuspended in 7 μl of water, 2 μl of Sequenase reaction buffer and 1 μl of the appropriate sequencing primer (0.5-1 pmol/μl). Annealing of the primer to the template DNA, was accomplished by warming the reaction to 65°C for 2 minutes, and slowly cooling to 30°C (over 30

minutes), then placing on ice. During the annealing reaction, 2.5  $\mu$ l of each of the G,A,T,C termination mixes was aliquoted out per sequencing reaction to a labelled microtitre tray (Nunc). Upon completion of the annealing reaction, 1.0  $\mu$ l of 0.1 M DTT, 2.0  $\mu$ l of dGTP labelling mix (diluted 5x in Milli-Q water), 0.5  $\mu$ l of [a-35S]dATP, and 2.0  $\mu$ l of Sequenase (diluted 8x in enzyme dilution buffer) were added to the annealed template and mixed. These labelling reactions were incubated for 2-5 minutes at room temperature. After incubation, 3.5  $\mu$ l of the labelling reaction was added to each of the four termination mixes in the microtitre tray. A further incubation at 37°C for 5 minutes was carried out and stopped by the addition of 4  $\mu$ l of stop solution. Sequencing reactions could then be stored at -20°C.

#### 2.10.2 Amplicycle Sequencing Kit Protocol

For each DNA sequencing sample, 2 µl of each of the G,A,T,C termination mixes were aliquoted into four labelled 0.2 ml strip tubes (for use with the Corbett FTS-960 thermal cycler). A cocktail was prepared for each DNA sequencing template as follows: 1.0 µl of 20 µM primer, 1.0 µl of [a-33P]dATP or dCTP, 4.0 µl of 10x cycling mix, template DNA (10 fmol of PCR product, or approximately 1-2 µg of Rapid Boil plasmid DNA [Section 2.6.1]), and Milli-Q water to 30 ml. 6 ml of this cocktail was then dispensed to each of the four termination tubes previously prepared. The tubes were then placed in a Corbett FTS-960 thermal cycler preheated to 95°C. The following cycling conditions were used:

Initial Step	95°C	1 minute	
Denaturation	95°C	30 seconds	
Annealing	65°C	30 seconds x 2	25
Elongation	72°C	1 minute	
Elongation	72°C	1 minute	
Cool	4°C		

Within 45 minutes of PCR completion, 4  $\mu$ l of stop solution was added to the tubes, and the reactions stored for up to 1 week at -20°C.

#### 2.11 RNA PROCEDURES

#### 2.11.1 Precautions Taken for Working With RNA

Glassware was acid washed in a chromic acid bath and baked for at least 2 hours at 180°C. Previously unopened bags of disposable tips and eppendorfs were autoclaved in

RNA treated glassware. Fresh disposable gloves were always worn when handling both RNA equipment and during RNA experiments. RNA solutions were only prepared from unopened stocks of reagents. A set of gilson pipettes dedicated to RNA work were always used.

#### 2.11.2 Agarose-gel Electrophoresis of RNA

Electrophoresis equipment dedicated to RNA work was used to avoid contamination with RNases. The gel box, gel combs and glass plates were presoaked overnight in 1% SDS and wiped clean with ethanol. A surface tension 1% agarose TAE gel was poured onto the glass plate. Approximately 7  $\mu$ g of RNA (prepared as in Section 2.13.2) was loaded in 1x RNA loading dye. The gel was electrophoresised at 80 V at room temperature until the dye had migrated 3/4 of the gel length. The gel was stained in ethidium bromide for 10 minutes and the RNA observed under UV light and photographed.

#### 2.11.3 Determination of Fungal RNA Concentration and Purity

The RNA sample was appropriately diluted and the absorbance measured by a Shimazdu spectrophotometer at a wavelength of 260 nm. The RNA concentration was calculated on the assumption that an absorbance of 1.0 at 260 nm is equivalent to 40  $\mu$ g/ml. RNA purity is indicated by the 260 nm/280 nm ratio, where pure RNA has an OD260/OD280 ratio of 2.0 (Current Protocols in Molecular Biology, 1989; Glasel, 1995).

#### 2.12 RNA PREPARATION

#### 2.12.1 Aspergillus nidulans Spore Suspension Preparation

MYG (Section 2.2.3) plates were inoculated with *A. nidulans* spores and incubated at 37°C for 2-3 days or until the culture was sporulating well. 5 ml of Tween 80/Saline solution (5 ml of tween 80, 1.6 g NaCl, Milli-Q water to 200 ml), was pipetted onto the surface of the MYG plate, and rubbed over the surface with a glass spreader to release spores. This spore suspension was pipeted onto another *A. nidulans* culture and the procedure repeated. The final spore suspension was passed through sterile cheese cloth to remove debris and pelleted by centrifugation at 3 K (Heraeus Sepatech Megafuge 1.0) for 5 minutes. The spore pellet was then resuspended in 2 ml of Saline (0.8% NaCl) and aliquoted into eppendorfs. The spore concentration was estimated by counting, using a haemocytometer.

#### 2.12.2 Isolation of RNA from Aspergillus nidulans

Total RNA was prepared for RT-PCR (Section 2.13) from strain A18 (Table 2.1) based on the method of Shirzadegan et al. (1991). Due to degradation of RNA and time constraints, total RNA (A18) was obtained from laboratory stocks prepared by R. Bradshaw. Approximately 400 mg of dry weight mycelium was ground to a fine powder with liquid nitrogen in an ice cold mortar and pestle. The powder was added to 4 ml of 80°C phenol extraction buffer (containing 0.1 M Tris HCl pH 8.0, 0.01 M EDTA, 1% SDS, in an equal volume of Tris-equilibrated phenol), immediately mixed by vortexing and incubated for 5 minutes at 80°C. An equal volume of chloroform was added, mixed by vortexing and centrifuged at 11 K for 15 min at 4°C. The top aqueous phase was then extracted with phenol/chloroform and centrifuged at 11 K for 15 minutes (centrifugations were performed in a SS34 rotor throughout). The aqueous phase was extracted with an equal volume of chloroform and centrifuged again as above. To precipitate the RNA, 1/2 volume of 8 M LiCl was added to the aqueous phase, and left at -20°C overnight. The RNA was pelleted at 12 K for 30 min at 4°C and then washed with 1 volume of 2 M LiCl, 0.05 M EDTA and centrifuged at 12 K for 30 min. Finally, the pellet was washed in 3 M NaOAc (pH 5.5) and resuspended in 200 µl of TE (10/0.1).

#### 2.13 RT-PCR ANALYSIS OF RNA

#### 2.13.1 Reverse Transcriptase Synthesis of cDNA

Before the RNA was transcribed into cDNA, it was DNaseI treated to remove any traces of DNA which may be preferentially amplified during the PCR step. The method for DNaseI treatment was based on that of Collett (1995). To 10 μg of RNA (Section 2.12.2), 1.0 μl of 100mM dithiothreitol (DTT) in 200 μl of DNaseI buffer (100 mM sodium acetate, 5 mM magnesium sulphate, pH 5), 50 U of RNase inhibitor, and 20 U of DNaseI (free of RNase) were added and incubated for 30 minutes at 37°C. Another 20 U of DNaseI was added and incubated for 30 minutes at 37°C. The RNA was then Phenol/chloroform extracted and ethanol precipitated (as in Section 2.5.7, except 3 M NH4Ac was used instead of 3 M NaOAc). RNA was resuspended in 20 μl of TE (10/0.1).

The DNaseI treated RNA and untreated RNA was reverse transcribed into cDNA using the method based on that by Kawasaki (1990). 1 µg of RNA was denaturated at 90°C for 5 minutes, then immediately put on ice for 5 minutes before it was added to the PCR cocktail. The cocktail contained the following: 2 µl of 10 x PCR buffer (500 mM KCl,

200 mM Tris-HCl [pH 8.4 at 23°C], 25 mM MgCl<sub>2</sub> and 1 mg/ml bovine serum albumin), 1 mM of each dNTP, 100 pmol of random primers (BRL), 40 units of RNase inhibitor, 200 units of Superscript reverse transcriptase (BRL) and Milli-Q water to 20 μl. Upon addition of denatured RNA to the cocktail, the reaction was mixed and incubated for 10 minutes at room temperature, at 42°C for 60 minutes, and then denatured at 95°C for 5 minutes. The reaction was then cooled on ice for 5 minutes and stored at -20°C.

#### 2.13.2 Amplification of cDNA by PCR

10  $\mu$ l of the cDNA reaction (prepared in Section 2.13.1) was added to the following: 4  $\mu$ l of 10 x PCR buffer, 2  $\mu$ M 'upstream' primer, 2  $\mu$ M 'downstream' primer, 2.5 units of Taq DNA Polymerase, and Milli-Q water to 50  $\mu$ l. The cDNA reaction tubes were then placed in a thermal cycler preheated to 95°C. The following cycle conditions were used to amplify the cDNA:

Initial Ste	95°C	2 minutes	
Denaturat	on 95°C	30 seconds	$\neg$
Annealing	60°C	45 seconds	x35
Elongation	72°C	60 seconds	/ <del></del>
Elongation	72°C	7 minutes	
Store	4°C		

#### 2.14 SOUTHERN BLOTTING AND HYBRIDISATION

#### 2.14.1 Southern Blotting

This method was based on that of Southern (1975). DNA to be blotted and hybridised was electrophoresed through a 1% agarose gel, stained and photographed as described in Section 2.6.2. The wells were cut off the gel and the gel dimensions measured. The gel was gently agitated in a tray containing depurination solution (250 mM HCl) for 15 minutes, then gently shaken in denaturating solution (500 mM NaOH, 500 mM NaCl) for 30 minutes and finally gently shaken in neutralisation solution (500 mM Tris (pH 7.4), 2 M NaCl) for 30 minutes. The gel was washed in 2 x SSC for 2 minutes.

While the gel was being treated, the blotting apparatus was assembled. Two sheets of Whatman 3 MM chromatography paper soaked in 20 x SSC (Section 2.3.12), were placed over a plastic trough with large wells at each end, so that the ends of the paper

dipped into these wells. The wells were then filled with 20 x SSC. The blotting trough was overlaid with gladwrap, and a grid was marked on the gladwrap of dimensions 2 mm less than the gel size. The grid was then cut out and the treated gel was placed, with well side down over the grid, so that the edges of the gel are on top of the gladwrap. A piece of nylon membrane (Hybond-N, Amersham; or Sartolan Blotting membranes) was cut out 2 mm less than the gel size, soaked in 2 x SSC, and placed over the gel. Any bubbles between the gel and filter were carefully removed. Two pieces of Whatman 3 MM were cut to 2 mm less than the gel, soaked in 2 x SSC and placed on top of the nylon filter. Another 2 sheets of dry Whatman 3MM of the same dimensions as above were placed on top of the wet 3 MM sheets. A pile of paper towels (5-8 cm deep) were then placed on top of the dry 3 MM sheets, followed by a 250 ml glass bottle filled with liquid as a weight. After blotting overnight, the blotting apparatus was disassembled and the nylon filter washed in 2 x SSC for 5 minutes. The filter was left to air dry between blotting paper and then baked under vacuum at 80°C for 2 hours.

#### 2.14.2 Preparation of [a-32P]dCTP-Labelled DNA Probe

DNA was labelled using the Ready-To-Go DNA Labelling Kit (Pharmacia). 25-50 ng of DNA (in a volume up to 45  $\mu$ l), was denatured in a boiling water bath for 3 minutes, then immediately put on ice for 2 minutes and centrifuged briefly. The DNA solution was then added to the tube of reagent mix and mixed by pippeting the solution up and down. 3-5  $\mu$ l of [a-<sup>32</sup>P]dCTP (3000 Ci/mmol) and water to a final volume of 50  $\mu$ l was added. The reaction was incubated for 30 minutes at 37°C.

Unincorporated nucleotides were removed from the labelled probe by using the ProbeQuant G-50 Micro Columns (Pharmacia Biotech) as described by the manufacturers.

#### 2.14.3 Southern Blot Hybridisation

The Southern blot (Section 2.14.1) to be probed was prehybridised in a sealed glass hybridisation tube for a minimum of 2 hours at 65°C in hybridisation solution (Section 2.3.9). After prehybridisation, the hybridisation solution was discarded, and replaced with 7 ml of fresh hybridisation solution. The [a-32P]dCTP labelled probe (Section 2.14.2) was denatured by boiling for 10 minutes immediately before it was added to the hybridisation tube. Hybridisation was carried out in a rotating oven at 65°C for a minimum of 6 hours, but usually overnight.

After hybridisation, the hybridisation fluid was discarded and about 100 ml of wash solution (3 x SSC, 0.2% SDS) was added. The filter was washed 3 times for 20 minutes at 65°C, and then wrapped (still wet) in gladwrap. The filter was placed in a X-ray cassette with intensifying screens against either Fuji Medical or Kodak Scientific Imaging X-ray film, and exposed for an appropriate period of time at -70°C. After exposure the film was developed (using standard procedures) in the dark room by placing in developing solution for 3 minutes, then in fixing solution for 3 minutes.

#### 2.14.4 Stripping Probes from Southern Blots

Filters to be stripped were gently shaken in a container containing a solution of boiling 0.1% SDS. The solution was allowed to cool to room temperature, the procedure repeated and the filter autoradiographed (Section 2.14.3), to check that the hybridised probe had been removed. If probe was still present, this process was repeated.

#### 2.15 LIBRARY SCREENING BY PLAQUE HYBRIDISATION

This method was based on that of Sambrook *et al.* (1989). Three rounds of library screening were performed to select the final positive plaques. Each round of screening involved plating the recombinant phage (Section 2.15.1), taking plaque lifts (Section 2.15.2), hybridisation of the filters to labelled probe (Section 2.15.3), and the selection of positive plaques (Section 2.15.3).  $\lambda$  phage DNA was then prepared from the positive plaques as detailed in Section 2.5.6.

#### 2.15.1 Plating Phage $\lambda$

The phage population to be screened was diluted in SM buffer (Section 2.3.8) at 3 x 10<sup>4</sup> pfu/ml for the primary screen, or 3x 10<sup>2</sup> to 3x 10<sup>3</sup> pfu/ml for the secondary and tertiary screens. 100 µl of diluted phage was then added to 100 µl of an overnight bacterial culture of LE 392 grown in LB (Section 2.2.1) supplemented with 0.2% maltose and 10 mM MgSO4.7H2O. The phage/bacteria mix was incubated for 30 minutes at 37°C, added to 3 ml of TB top agarose (Section 2.2.6) at 45-50°C, then poured onto fresh LB agar plates. The top agarose was allowed to set, then the plates were incubated at 37°C until small plaques formed (about 7 hours). Plates were stored at 4°C for a minimum of 1 hour or overnight before plaque lifts were taken.

#### 2.15.2 Plaque Lifts

Nylon filters (Biotechnology Systems, 82 mm disks) were marked asymmetrically with a needle for orientation purposes. Duplicate lifts were taken from each plate for both the primary and secondary rounds of library screening. The filter was placed on the LE 392 phage lawn and left for 1 minute for the first lift and 2 minutes for the duplicate lift, once the filter was completely wet. The filters were then placed DNA side up on 3 layers of filter paper soaked in 500 mM NaOH, 500 mM NaCl for 2 minutes, then 500 mM Tris-HCl (pH 7.4), 2 M NaCl for 5 minutes, and finally in 2x SSC for 1-2 minutes. The filters were air dried and the DNA fixed onto the filters by microwaving on high for 1 minute.

#### 2.15.3 Hybridisation of λ Phage DNA to [a-32P]dCTP-Labelled Probe

Library filters were hybridised to [a-32P]dCTP-labelled DNA (Section 2.14.2) and autoradiographed as described in section 2.14.3, except plastic pots (10 cm diameter) were used (10 filters/pot), and the hybridisation and washing steps were performed in a shaking water bath.

Positive plaques were identified by a hybridisation signal on the autoradiogram at identical sites on duplicate filters. The corresponding plaques were located on the plate by alignment of the orientation markings. A plug of agar containing the positive plaques was removed by using a 1 ml Gilson pipette tip from which the end 5-10 mm had been cut off. The plugs were then put into an eppendorf containing 1 ml of SM buffer and 1 drop of chloroform and the phage left to elute into the buffer at 4°C overnight.

# CHAPTER 3 CONSTRUCTION OF PROMOTER-REPORTER VECTORS AND DISCOVERY OF A SEQUENCING ERROR

#### 3.1 INTRODUCTION

To analyse the promoter region of the Aspergillus nidulans cytochrome c (cycA) gene, in particular the role of the putative HAP1 binding site (Raitt, 1992), a promoter reporter vector was constructed containing the cycA promoter fused to the lacZ gene of Escherichia coli. The construction of a second vector which was to contain the cycA promoter with the putative HAP1 site deleted ( $\Delta$  HAP1) fused to the lacZ gene, was also attempted.

#### 3.2 CLONING STRATEGY

#### 3.2.1 The lacZ Reporter Vector

The 11.5 kb reporter vector used in this study, pAN923-42B<sub>Bg/II</sub> (R117, Table 2.1), constructed by Van Gorcom *et al.* (1986), and modified to create a mutant *argB* gene by Punt *et al.* (1990), (Figure 3.1), allows in-phase translational fusion to the *E. coli lacZ* gene. pAN923 contains a unique *Bam*HI site in one of three possible translational reading frames (41B, 42B, 43B) in front of the *lacZ* gene. By insertion of the *cycA* 5' fragments into the *Bam*HI site at 42B, a translational fusion was obtained with the 0.68 kb fragment containing 578 bp of promoter and upstream region, along with 54 bp of coding sequence (hereafter called the 'promoter' region).


The reporter vector which contains a mutant argB gene, can be transformed into an A.  $nidulans\ argB2$  mutant strain and integrated at the argB gene locus. Complementation of the argB mutations to yield an  $argB^+$  phenotype only occurs if homologous integration occurs, thus allowing selection for this event. After confirmation by Southern blotting of single copy integration,  $\beta$ -galactosidase assays and Northern blots can be performed to determine  $in\ vivo$  promoter activity, and to measure transcriptional and translational lacZ reporter expression.

#### 3.2.2 PCR Amplification of the cycA Promoter Fragments

PCR (Section 2.7) was used to amplify the *cycA* promoter fragments using plasmid R41 DNA (Table 2.1) which contains a 2.3 kb *cycA* genomic fragment (Raitt *et al.*, 1994). Primers for PCR were designed to incorporate a restriction enzyme site at each

#### Figure 3.1 The lacZ reporter vector pAN923

The cycA fragments containing the promoter and 5' coding region were inserted into the BamHI (B) site of pAN923-42B. Thick lines represent A. nidulans DNA; thin lines represent E. coli DNA. B, BamHI; Bg, BglII; E, EcoRI; H, HindIII; N, NruI; P, PstI; S, SalI; Sp, SphI; Ss, SstI; X, XhoI; Xb, XbaI.



end of the PCR product (Figure 3.2). Primers RB1 and RB2 (Table 2.2) include BamHI recognition sequences at the 5' end and were used to amplify the entire cycA promoter region (published by Raitt et al. (1994)), along with the 5' coding region to give a total of 0.68 kb. Both the vector and the 0.68 kb fragment were then digested with BamHI, followed by ligation to allow the cycA fragment to be fused in frame at the BamHI site of the lacZ gene. To obtain a translational fusion, the promoter fragment must be orientated so that the protein-coding region of the cycA gene is fused next to the protein-coding region of the lacZ gene. Diagnostic restriction digests were necessary to distinguish the orientation of the ligated promoter fragments (see Table 3.2, Figure 3.3).

To amplify the  $\Delta$  HAP1 *cycA* promoter fragment for the second construct, primers RB1 and RB4 were used to amplify the region upstream of the  $\Delta$  HAP1 site, producing a 190 bp fragment. Primers RB3 and RB2 were used to amplify the region downstream of the  $\Delta$  HAP1 site, producing a 400 bp fragment. The internal primers, RB3 and RB4 both include a *Sal* I restriction site. After *Sal* I digestion both the 400 bp and 190 bp fragments can be ligated together to form the 0.59 kb  $\Delta$  HAP1 promoter fragment which will contain a 54 bp deletion of the putative HAP1 binding region. The 0.59 kb fragment contains 536 bp of promoter region, along with 54 bp of coding region. Digestion of the  $\Delta$  HAP1 promoter with *Bam*HI, allows this promoter to be cloned, as with the 0.68 kb promoter, into pAN923-42B<sub>BgIII</sub>.


PCR conditions for the amplification of the *cycA* promoter fragments were established by Saunders, 1993. However, R41 plasmid (Table 2.1) DNA concentration was optimised to produce a high yield and purity of PCR product; a concentration of 0.1 ng/μl was found to be optimal as a template for amplification of the promoter fragments.

### 3.2.3 General Preparation of pAN923-42B Vector and 0.68 kb cycA Promoter

Plasmid pAN923-42B (R109, Van Gorcom et al., 1986) was initially used as the cloning vector, before the modified pAN923-42B $_{BgIII}$  (R117) vector (Punt et al. 1990) was available. After plasmid preparation, pAN923 vectors were digested to completion with BamHI and CAP treated as described in section 2.8.2. This was necessary because pAN923 provides no effective selection system for recombinants in E. coli and CAP treatment prevents self-ligation of the vector, thus abolishing high vector background on selective LB ampicillin plates.

Figure 3.2 PCR primer binding sites within the cycA promoter fragments

The relative positions of the primer sites used for amplification of the *cycA* promoter fragments are shown. B and S represent *BamHI* and *SalI* restriction enzyme sites respectively. The blue coloured box identifies the putative HAP1 binding region.





The 0.68 kb and Δ HAP1 promoter fragments obtained by PCR (Section 2.7), were gel purified (Section 2.8.1) followed by phenol/chloroform and ethanol precipitation purification (Section 2.5.7) to remove any residual Taq polymerase terminal transferase activity which can add adenosine bases to the digested 3' ends of PCR products (Clark, 1988; Bennet and Molenaar, 1994). After purification, the promoter fragments were digested with *Bam*HI and/or *Sal*I as appropriate (Section 2.6.1), resulting in the removal of a small number of base pairs from each end. Since the change of length would be too slight to detect on an agarose gel, a control digest (to check the enzymes) was performed on plasmid DNA to give known fragment sizes. Following restriction enzyme digestion (Section 2.6.1) the promoter fragments were purified (Section 2.5.7) to remove the enzyme activity.

### 3.3 LIGATIONS AND TRANSFORMATIONS OF THE 0.68 kb cycA PROMOTER

#### 3.3.1 Ligation of 0.68 kb promoter directly into pAN923-42B

The efficiencies of vector digestion, CAP-treatment and ligation were checked by comparing yields of transformants from electroporation of the following samples; self-ligated *Bam*HI-digested and CAP treated pAN923 vector (Cut, Cap, Lig), self-ligated *Bam*HI digested pAN923 vector (Cut, Lig), *Bam*HI-digested and CAP treated pAN923 vector (Cut, Cap) and uncut pAN923 vector. The results of these controls are included in Tables 3.1, 3.3 and 3.6.

The controls for transformation experiments A, B, and C, involved in the ligation of the 0.68 kb promoter into pAN923-42B (Table 3.1), suggested that the vector was poorly CAP treated but efficiently *Bam*HI digested (shown by the high numbers of transformants per µg of Cut, Cap, Lig vector DNA and low numbers for Cut, Cap vector respectively). However, the comparison of control samples to determine capping efficiency was inaccurate due to the low numbers of transformants produced per plate (2-10 colonies). The transformation of control samples with a higher DNA concentration would have been more accurate. Electroporation and ligation efficiencies were satisfactory (shown by the numbers of transformants for uncut vector and Cut, Lig vector respectively).

The first attempt at ligating the 0.68 kb promoter fragment into pAN923-42B and transforming into *E. coli* XL-1 cells was unsuccessful (Transformation A, Table 3.1). After transformation of the ligation reactions into *E. coli* XL-1 cells, 7 transformants were obtained (Table 3.1, (A)) and checked by *Bam*HI digestion (Section 2.6.1) of plasmid DNA prepared by the rapid boiling method (Section 2.5.1). If ligation was

Table 3.1 Transformations A, B & C (with pAN923-42B vector)

DNA transformed into E. coli XL-1 cells <sup>a</sup>	DNA/plate (ng)	Number of colonies per Plate	Number of transformants per μg DNA (mean/experiment)
A 0.68 kb insert and pAN923 vector ligation (1:1)	1.2 1.5	0	0
0.68 kb insert and pAN923 vector ligation (2:1)	1.5 1.5 3.0 3.0	2 1 4 0	1.1x10 <sup>3</sup>
B 0.68 kb insert and pAN923 vector ligation (1:1) <sup>b</sup>	1.6 1.6	2 2	1.3x10 <sup>3</sup>
0.68 kb insert and pAN923 vector ligation (2:1)	1.9 1.9 1.6 1.6	3 2 1 0	$1.0 \times 10^3$
0.68 kb insert and pAN923 vector ligation (2:1) <sup>c</sup>	1.5 1.5	3 3	2.0x10 <sup>3</sup>
0.68 kb insert and pAN923 vector ligation (3:1)	2.0 2.0 1.7 1.7	0 1 3 1	8.0x10 <sup>2</sup>
C 0.68 kb insert and pAN923 vector ligation (3:1) <sup>d</sup>	6.2	53	8.5x10 <sup>3</sup>
Controls Cut, Cap pAN923 vector	1.5 8.5	0 0	0
Cut, Cap, Lig pAN923 vector	1.4	2	$1.4 \times 10^3$
Cut, Lig pAN923 vector	2.5	10	$4.0x10^3$
Uncut pAN923 vector	0.6	235	4.0x10 <sup>5</sup>

a numbers in brackets indicate insert:vector ratio used

b indicates re-electroporation of (1:1) ligation from section A

c indicates re-electroporation of (2:1) ligation from section A

d indicates re-electroporation of (3:1) ligation from section B

successful, the 11.5 kb vector band, and the 0.68 kb promoter band should both be present (Table 3.2). For all 7 transformants, only the 11.5 kb vector band was present, indicating that ligation of promoter into pAN923 had not occurred. The ligation reactions were repeated and both old and new ligations were electroporated into E. coli XL-1 cells (Transformation B, Table 3.1). In total, 21 transformants were obtained, and PCR (Section 2.7) was used to screen the transformants by using primer pair RB1 and RB2 (Table 2.2) to detect the 0.68 kb promoter. Two transformants contained a PCR product of approximately 0.68 kb and were further checked by BamHI digestion of rapid boil plasmid DNA (Section 2.6.1) to confirm the presence of ligated promoter. Promoter orientation and copy number were determined by SphI digestion. Restriction digest patterns showed that one transformant (R120, Table 2.1) contained the 0.68 kb promoter ligated into pAN923 in the wrong orientation (Table 3.2, Figure 3.3). The other transformant produced a 7.2 kb fragment with BamHI digestion and 6.1 kb plus 1.3 kb fragments with SphI digestion. This restriction profile was not expected for either vector only or vector with insert, and must therefore indicate either a contaminant or a deletion/re-arrangement event.

The remaining ligation mixture which produced the R120 plasmid from transformation B (Table 3.1), was electroporated into *E. coli* XL-1 cells (Section 2.8.4.2), resulting in 53 transformants (Transformation C, Table 3.1). These transformants were checked by PCR (Section 2.7) as described above, and all were shown to contain the promoter fragment. *Bam*HI digestions of 25 transformants showed that 23 contained insert, 1 contained vector only, and 1 contained no plasmid DNA. The 23 insert-containing transformants were checked by *Sph*I digestion to check promoter orientation. All transformants contained insert ligated into the vector in the wrong orientation which was surprising, since 50% of inserts would be expected to be ligated into the vector in the other orientation. Perhaps there was some unknown preference for the insert ligating in the wrong orientation. Another possibility was that all transformants had arisen from one transformants may have arisen from contamination by plasmid R120, but this is unlikely since transformation control experiments were negative. In summary, time and resources seemed wasted by further checking of the remaining transformants.

Further pAN923/0.68 kb promoter ligations were performed with freshly prepared vector (Table 3.3). However, the vector was not cut to completion by *BamHI*, (as shown by the number of transformants with cut, capped, unligated vector) and the cut ends were poorly capped (as shown by the increased number of transformants with capped and ligated vector). This resulted in many transformants to screen for the presence of insert (Transformation D, Table 3.3). Despite the large number it was thought worthwhile to


Table 3.2 Expected Restriction Digestion Fragment Profiles for Ligation of 0.68 kb Promoter into pAN923-42B

estriction Digest	Fragment Sizes (kb)
BamHI	11.5, 0.68
SphI	7.16, 4.15, 0.87
BamHI	11.5, 0.68
SphI	7.68, 4.15, 0.37
9	SphI

Figure 3.3 Restriction enzyme analysis of plasmids pAN923, R122 and R120

Restriction digests of plasmids pAN923, R120 (from transformation B, Table 3.1), and R122 (from transformation G, Table 3.6) were carried out individually with *Bam*HI and *Sph*I. Lane 2: pAN923, *Bam*HI. Lane 3: pAN923, *Sph*I. Lane 4: R122, *Bam*HI. Lane 5: R122, *Sph*I. Lane 6: R120, *Bam*HI. Lane 7: R120, *Sph*I. Lane 1 contains a 1 kb ladder (Gibco BRL, Life Technologies). The size of the bands indicated are in kb.

### 1 2 3 4 5 6 7



12,216\_\_ 10,180\_\_ 8,144\_\_ 6,108\_\_ 5,090 4,072

3,054

2,036

1,636

1,018

506,517 396 344

Table 3.3 Transformation D (with pAN923-42B vector)

DNA transformed into <i>E. coli</i> XL-1 cells	DNA/plate (ng)	Number of colonies per Plate	Number of transformants per µg DNA (mean/experiment)
D			
0.68 kb insert and	5.5	9	$1.2x10^4$
pAN923 vector	6.9	113	
ligation	9.6	179	
Controls			
Cut, Cap pAN923 vector	50	91	1.8x10 <sup>3</sup>
Cut, Cap, Lig pAN923 vector	5.0	800	1.6x10 <sup>5</sup>
Uncut pAN923 vector	0.5	360	7.2x10 <sup>5</sup>

screen for recombinants by colony hybridisation (Section 2.9.2). Unfortunately, there was a lot of background present on the filters affecting the detection of positive colonies. However ten positive colonies were selected for further investigation, but upon restriction enzyme analysis of rapid boil plasmid DNA (Section 2.5.1) were found to contain no insert.

#### 3.3.2 Ligation of 0.68 kb Promoter into T-Tailed pUC18 Vector

Difficulties experienced with cloning the 0.68 kb promoter directly into pAN923 vector (Section 3.3.1) were due to a number of factors. One major problem was probably inefficient cutting of *Bam*HI sites on a large proportion of the PCR product ends, thus decreasing the ligation efficiency. The T-tailed vector cloning approach (Marchuk *et al.*, 1991) avoids this problem by making use of the adenosine base overhang generally added to PCR products by Taq polymerase, allowing ligation to a 3' T-Tail attached to the pUC18 vector (kindly provided by R. Johnson). Once the promoter is cloned into the T-Tailed vector, it can be digested using flanking sites in the polylinker and gel purified (Section 2.8.1) before ligation into pAN923.

Transformants obtained from the T-tail ligations produced unexpected restriction digestion profiles. Eighteen white transformants were obtained from three separate ligation experiments and were checked by PCR (Section 2.7) using primers RB1 and RB2 (Table 2.2) to detect insert. Eight transformants contained a 0.68 kb PCR product and were further checked by digestion individually with BamHI and SphI. All eight transformants produced a 5.4 kb fragment with BamHI and a 4.1 kb and 1.3 kb fragment with SphI (see Table 3.4 for expected restriction enzyme profiles). After considering all possible combinations of insert and vector, no possibilities matched the data observed for these transformants. The total construct size (5.4 kb) suggested that the transformants contained pUC18 (2.7 kb) ligated to four 0.68 kb promoters, or two ligated pUC18 plasmids, but the SphI digestion results did not support this hypothesis. The alternative explanation was that DNA contamination had occurred.

Lack of success with the T-tailing cloning approach may be due to a number of reasons. The ligation efficiency may not have been optimal, causing a decrease in the number of white transformants obtained. Also, the terminal transferase activity of Taq polymerase is only weak (Parry, 1993) and thus may add low numbers of adenosine bases to the PCR fragments.

Table 3.4 Expected Restriction Enzyme Profiles for Ligation of 0.68 kb Promoter into Plasmid pUC18

Restriction Digest	Expected Fragment Sizes (kb)
BamHI	2.7, 0.68
SphI	2.8, 0.60
	OR*
	3.3, 0.13

<sup>\*</sup> depending on orientation

## 3.3.3 Ligation of 0.68 kb Promoter into Commercially CAP-treated pUC18

Another approach to facilitate the cloning of the PCR-amplified 0.68 kb promoter into pAN923, was to digest with *Bam*HI and subclone it into commercially CAP treated pUC18 plasmid (BRL/Life Technologies). Successful ligation of insert in either orientation should show that both *Bam*HI ends of the promoter were intact and functional. Cloning into pUC18 was expected to be easier than cloning directly into pAN923 since it has the advantage of blue/white selection. Another advantage of ligating the promoter into pUC18, a multiple copy number plasmid, was that large quantities of digested promoter could be obtained, gel purified (Section 2.8.1), then ligated into pAN923 to increase the chances of success.

The pUC18 ligations with the PCR derived 0.68 kb promoter, produced either no transformants or they were not stable transformants (Transformation F, Table 3.5). However, ligations with pUC18 and the 0.68 kb promoter derived from plasmid R120 (in which the promoter had previously been cloned in reverse orientation, Table 2.1) produced 12 transformants (Transformation E, Table 3.5). Restriction enzyme analysis of the 12 transformants from transformation E showed that four contained an insert of the expected size for the promoter, two of which were ligated in each of the different orientations. One representative of the pUC18 clones was stored in glycerol (R121, Table 2.1) and used for all further sources of 0.68 kb promoter. Since the only difference between transformants E and F was the source of the 0.68 kb promoter, it was suggested that the problem with obtaining recombinants with transformation F (using the PCR amplified promoter), was due to inefficient digestion of the promoter ends. This may have been because the enzyme recognition site (incorporated by the PCR primers) was too close to the ends of the PCR product (2 bp) for efficient cutting.

#### 3.3.4 Ligation of 0.68 kb promoter into pAN923-42B

A BamHI digestion of plasmid R121, followed by gel purification, was carried out to obtain the 0.68 kb promoter fragment for ligation into pAN923-42B $_{BgIII}$ . Now that large quantities of insert with BamHI ends were easily obtainable for ligation, larger scale ligation reactions comprising 50 ng of insert in a 3:1 or 1:1 insert to vector ratio were set up to increase ligation frequency. Due to large volume sizes (30-40  $\mu$ l), these samples were transformed by the calcium chloride method (Section 2.8.4.3). Table 3.6 displays the transformation results.

Table 3.5 Transformations E & F (with CAP-treated pUC18 vector)

DNA transformed into E. coli XL-1 cells <sup>a</sup>	DNA/plate (ng)	Number of colonies per Plate	Number of transformants per µg DNA
E			
0.68 kb insert (from R120) and pUC18 vector	5.3	12	$2.3 \times 10^3$
ligation (1:1)			
F			
0.68 kb insert (PCR) and	7.5	0	0
pUC18 vector ligation (2:1)			
0.68 kb insert (PCR) and	8.8	0	0
pUC18 vector ligation			
(3:1)			

a numbers in brackets indicate insert:vector ratio used

Table 3.6 Transformation G (with pAN923-42B vector)

DNA transformed into E. coli XL-1 cells <sup>a</sup>	DNA/plate (ng)	Number of colonies per Plate	Number of transformants per µg DNA
G 0.68 kb insert and pAN923 vector ligation (3:1)	242	1280	5.3x10 <sup>3</sup>
0.68 kb insert and pAN923 vector ligation (1:1)	655	704	1.1x10 <sup>3</sup>
Controls Cut, Cap pAN923 Vector	20	48	2.4x10 <sup>3</sup>
Cut, Cap, Lig pAN923 vector	10	37	3.7x10 <sup>3</sup>
Uncut pAN923 Vector	0.5	360	7.2x10 <sup>5</sup>

a numbers in brackets indicate insert:vector ratio used

The CAP treatment controls for transformation G (Table 3.6) indicate that the vector has not been cut or CAP treated to completion. In addition, the number of transformants from the promoter ligation experiments are approximately equivalent to the number of transformants present in the cut, cap, lig control. Therefore the majority of transformants were expected to be circular pAN923 vector. However, two transformants from each of the pAN923 vector:insert ligations were checked to confirm the nature of the transformants. One of these transformants contained, in addition to the vector band, the 0.68 kb promoter when digested with *BamHI*. An *SphI* digest of this transformant demonstrated that the 0.68 kb promoter was ligated into pAN923 in the correct orientation with the following band sizes: 7.1 kb, 4.15 kb, 0.80 kb (Table 3.2, Figure 3.3). This construct was named R122 (Table 2.1).

### 3.4 LIGATIONS AND TRANSFORMATIONS OF THE cycA $\triangle$ HAP1 PROMOTER

All ligations and transformations of the  $\Delta$  HAP1 promoter were unsuccessful (results not shown). As with the 0.68 kb promoter, ligations of the  $\Delta$  HAP1 promoter were attempted with both pAN923 and commercially CAP treated pUC18 plasmids. The lack of success with the  $\Delta$  HAP1 promoter ligations was probably due to the increasingly more difficult nature of three way ligations and hence a lower ligation frequency. Similar problems with the enzyme sites at the ends of the PCR products, (as discussed with the 0.68 kb promoter) were also expected to be the cause of unsuccessful ligation with the  $\Delta$  HAP1 promoter fragments. Due to unexpected results arising in the sequence of the whole promoter (Section 3.5), cloning of the cycA  $\Delta$  HAP1 was not pursued further.

#### 3.5 SEQUENCING OF THE 0.68 kb PROMOTER IN PLASMID R122

Sequencing across the junction in the new R122 vector construct, containing the 0.68 kb promoter in the pAN923 reporter vector, was essential to determine the exact position of the *cycA* gene ATG start codon relative to the *Bam*HI insertion site in the MCS. This information was required to confirm that a functional (in-frame) translational fusion was obtained. In addition, the promoter was amplified by PCR using Taq polymerase, which has a mis-incorporation error rate of 8.9x10<sup>-5</sup> errors/bp (Cariello *et al.*, 1991). Therefore, sequencing of the promoter was required to check there were no mutations present in the promoter sequence which could affect promoter expression studies, or in the coding sequence to put the *cycA* coding region out of frame.

pUC/M13 universal forward and RB1 primers (Table 2.2) were used to obtain double stranded sequence using the Sequenase Version 2.0 (Section 2.10.1) and Amplicycle

(Section 2.10.2) sequencing protocols. The preferred method of sequencing was the Amplicycle method, since it consistently produced high quality sequence using a quick and simple protocol. Plasmid DNA was prepared by small scale alkaline lysis preparation (Section 2.5.2) and purified by Wizard PCR preps DNA purification system (Section 2.8.1) for use with the Sequenase kit. Rapid boil DNA (Section 2.5.1) was used for sequencing with the Amplicycle kit.

Upon comparing the sequence obtained from plasmid R122 with the published sequence (Raitt *et al.* 1994), a discrepancy was noticed, with six thymidine bases obtained rather than the expected four (Figure 3.4). Unfortunately, the region of this discrepancy occured within the published coding region of the *cycA* gene, thus disrupting the reading frame. Either Taq polymerase had slipped during the amplification of the promoter, incorporating two extra thymidine bases, or the published sequence was incorrect. To determine which of these alternatives was correct, sequence was obtained from plasmid R41, which contains a sub-clone from a genomic library (Table 2.1), using primer RB2 (Table 2.2). Sequence from R41 over the region with the discrepancy clearly confirmed that six thymidine bases were present (Figure 3.5). Therefore the published sequence of the *cycA* gene by Raitt *et al.* (1994) was incorrect over this region. Figure 3.6 shows the published nucleotide sequence of the *cycA* gene as published by Raitt *et al.* (1994).

### Figure 3.4 Comparison of the cycA nucleotide sequence from plasmid R122 with the published cycA sequence (Raitt, 1992)


The region spanning the sequencing discrepancy has been presented. The top strand represents sequence obtained from plasmid R122 (see Figure 4.5), whereas the bottom strand represents the published nucleotide sequence of the *cycA* gene (Figure 3.6, Raitt *et al.*, 1994). The boxed area in red highlights the region in which the sequencing error occured, and the published ATG start codon is shown in green.

# Figure 3.5 Autoradiograph of sequence obtained from the cycA gene cloned into plasmid R41

Nucleotide sequence from the non-coding strand of plasmid R41 is presented. Six adenosine bases were observed in the A lane (lane 2) instead of the expected four.

R122 403 ATGCTAATCAATTGTTTTTCCAGGCGACT 432

cycA +1 ATGCTAATCAATTGTTTT CCAGGCGACT 28



GATC1

<sup>1</sup> The sequence presented in this autoradiograph was obtained from the non-coding strand of plasmid R41 (Table 2.1), therefore this sequence represents the reverse and complement of the coding strand from plasmid R122 (Table 2.1) in Figure 3.4.

Figure 3.6 The published nucleotide sequence of the cycA gene (Raitt et al., 1994)

The nucleotide sequence of the *cycA* gene as published by Raitt *et al.* (1994) is displayed. Nucleotides are numbered from the A of the published methionine translational start residue (+1). The alternative translational start site proposed in this study is indicated by the pink ATG. The major transcriptional start site (published by Raitt *et al.*, 1994) for the 2 intron model is indicated below the red asterisk. The two intervening regions (Introns I, II) are displayed in blue with the corresponding splice consensus sequences underlined. Consensus splice sequences for the proposed additional intron are underlined and coloured blue. The predicted amino acid sequence is shown below the coding strand using the standard one-letter code. The conserved CKQCH domain and the Gly 8 residue are indicated in green in the amino acid sequence. The position of the observed sequencing error is indicated in red. The putative HAP1 site proposed by Raitt (1992) is marked by green highlighting.

						•							•						,
-649	TACTO	TGA	GCAC	AGT	GCI	'GAA	TCA	TCC	TAC	CTC	TGA	TTG	GTC	AAT	TCC	CAG	ATC	ACG	GGI
-589	GTCGI	CGG	GGGC	GCG	ACC	AAG	AAA	CCA	GCT	CTA	CAA	ATT	ccc	TCC	AAG	· TTT	TTI	TCT	TCC
-529	CTTTG	GCC	AGTO	CGC	TTG	ACT	TGA	ATT	CTG	TCT	TTC	ATC	TTC	TCT	GCT	ACA	TAC	AAC	TCI
-469	TTGTA	CTA	TACC	ACT	TAC	CTC	TTT.	ACA!	FAA	ccc	TTT	CTC	TCT	ACC	TCT	TTA	TTI	TTT	ATC
-409	ACTCA	CAA	· TGGC	TAA	GGG	ICGG	TGA	CAG	· CTA	CTC	TCC	TGG	TAA	GTA	GTT	TGA	ITA	CAT	· CTC
-349	TTCGT	TTC	GGCI	AGG	CGC	TCT	CTG	CTG	GGG	GTG.	ATA	TTC.	ACT	CCA	CTG	CGT	GCA	TTG	TCG
-289	AAGTG	AAC	GATA	TGT	GAG	ACA	CAG	GCT	GTG.	TAA	GAT	GTG	· TTG	GGT	CTG	GTG	AGA	ACA	TTG
-229	TCCGA	TCC	AACA	CAA	GCI	CAA	AGT	TGC	CCA	CTT	CTG	GAT	cgc	CAT	TTG	ATC	GCC	AGC	ACA
-169	ATACA	ATT	TCTA	CTT	CTA	TCG	CGC	GCT	GGC.	AAT	CCT	CAC	· TTT	GCA	GCG	GAT	GCI	TTA	TTC
-109	TTCAT	CGT	CGTC	GCC	ACA	ATG	ACG.	AGC!	rTC	GAC	GCT	T'AC'	TGC	TTC	ACG	ACC	ACI	CTC	AGC
-49	ATGCG	CGA	AGCC	GAA	CTG	GAA	GAG	CTG	GAG	AAA	GAG.	AAA	GCG	GAC	CAG	AAT M	GCI L	T'AA'I I	CAA N
11	TTGTT C F	TTC		CGA D	SECOND ST	TAC		GGG'			-		CGA E	GAC T	CCG R	TTG	CAA	GCA	GTG
71	CCACA		TCGA E	GAA N	.CGG	CGG G	CGG	CCA(	CAA	GGT V	CGG	CCC P	CAA	CCT L	CCA H		TCI L	CTT F	CGG G
	((()) 25		•					115014			510	(E)	•	1020			1000	1752	
131	CCGTA R K	111444	G G		.GGC	TGG G	AGG G	Y Y	2GC	Y Y	T	D D	TGC A		K	GCA Q	GGC A	CGA D	CGT V
191	CACCI T W			GAA N		TCT L	GGT	ACA	ATC	CCA	TGA	CAG	стс	TAA	CAG	CTC	TGG	GCC	GAT
251	TGCT		•																0.000
							F	K	Y .	L	E	N	. P	K	K	. Y	1	F	
311	GTACC T		ATGG M A																1000
371	ATGTA	ACG	CTGC	TTA	CCA	CGG	ATA	GGG	CAC	ATA	GGC	TAA	<u>C</u> AG	GAT	GCA	CAG	CTA Y		CAA K
431	GGAGA E S					CGT	TCG	CGA'	TTA	GAC	GAG	ATA	AAC	CGC	ccc	CCT			
491	CGAGO	GCC'	TCTO	GCI	AGG	TGA	CAG	GCG	GGT	ACT	GTA	ACA	· TTA	CAC	CTA	GAC	CTG	GTI	TTC
551	AAGGT	CGT	CGGG	SACA	TGG	GAGG	ATA	TTA	rag.	ATC	TTG	TTT	CCT	TCG	CCA	TCC	TTO	TCT	ATA
611	TCTTA	TTC	TTTC	CTI	TAC	CTI	GAC	GAG	TGT	TTT	TTC	AGC	· TTT	GTG	GTA	cc			

# CHAPTER 3 DISCUSSION

### 3.6 PROBLEMS WITH CLONING INTO pAN923-42B VECTOR

The difficulties experienced with cloning the cycA promoter fragments into pAN923-42B were numerous and are summarised as follows. Firstly, the cloning strategy required ligation to occur through BamHI sticky ends. Since the BamHI recognition site was incorporated into the PCR product (positioned 2 nucleotides from the terminus) by the PCR primers, inefficient digestion of the promoter fragments near the ends of the PCR molecules was probably the cause of reduced ligation efficiency. For example, subcloning of the 0.68 kb promoter into pUC18 (Section 3.3.3) was successful with promoter derived from the BamHI digestion of plasmid R120 (Transformation E, Table 3.5), but unsuccessful (under identical conditions) when using PCR-amplified and BamHI digested 0.68 kb promoter (Transformation F, Table 3.5). Thus, the enzyme was probably unable to efficiently digest the PCR product. Secondly, the pAN923 vector contains no effective selection system for recombinants in E. coli. The vector therefore required CAP treatment to abolish high vector background, which was usually inefficient. Attempts were made to increase the capping efficiency by using a fresh batch of CAP enzyme, altering CAP enzyme concentration, incubation times, and concentrations of proteinase K. However, no overall improvement was observed from these modifications. Thirdly, the requirement for a translational fusion to investigate a potential translational control mechanism required clones to be screened for insert orientation, thus in retrospect a directional cloning approach would have been desirable.

### 3.7 IMPLICATIONS FROM THE DISCOVERY OF THE SEQUENCING ERROR IN THE PUBLISHED cycA GENE BY RAITT et al (1994)

Discovery of the presence of six thymidine bases in the *cycA* nucleotide sequence rather than the expected four as published by Raitt *et al* (1994) had many implications. The addition of two extra thymidine bases in the coding region of the *cycA* gene disrupts the published translational reading frame. This means that the long sought after translational fusion of the *cycA* promoter with the *lacZ* gene of pAN923-42B (R122, Table 2.1) was not obtained. Moreover, the discovery of this sequencing error brought into question the validity of the open reading frames of the cytochrome *c* gene as published by Raitt *et al* (1994). The published amino acid sequence downstream from the sequencing error would now code for different amino acids since the reading frame has altered. However, the published amino acid composition of the *cycA* gene shows extensive homology to both the yeast *CYCI* gene and the *Neurospora crassa* cytochrome *c* gene and contains all

the functional features associated with the cytochrome c protein. These features include a conserved heme binding domain CKQCH (position 21 to 25 in the amino acid sequence, Figure 3.6), and conserved glycine residues required for winding of the polypeptide around the heme molecule (Raitt, 1992). One of these invariant glycine residues, Gly 8 (Figure 3.6), falls immediately after the sequencing error, and is therefore no longer in frame with the published ATG codon with the sequencing correction (Figure 3.7). Since all the above features are found and conserved in the cycA protein as published by Raitt et al. (1994), it would appear that the published reading frame is correct. However, the published ATG start codon is no longer in the correct reading frame to produce the conserved cytochrome c protein, and therefore cannot be the translational start site.

Closer scrutiny of the published *cycA* nucleotide sequence revealed that both the six thymidine bases and the published ATG codon may fall within an intron region not previously detected. Filamentous fungal consensus splice sites for an additional intron were noticed at the 5' site (5'-GTANGT-3'), a possible branch point (5'-PyGCTAAC-3') and the 3' site (5'-CAG-3') at nucleotide positions -371, -77, and +19, respectively as seen in Figure 3.6 (Gurr *et al.* 1987). Evidence of this intron would further establish that the published translational start point is not correct.

Further examination of the cycA nucleotide sequence revealed a new potential translational start point, upstream of the putative intron, at position -402 nt in Figure 3.6 (highlighted in pink). This ATG codon would give the correct reading frame to produce the conserved cycA protein after the splicing of the putative additional intron. Moreover, re-calulation of the transcription start point (by moving the transcription site point in the 5' direction to take account of the additional intron) indicates the mRNA transcript commences at -230 nt. The suggested ATG codon would also become the first AUG in the mRNA, and translation of fungal genes usually occurs from the first AUG (Kozak, 1991; Gurr et al 1987). The sequence context of the AUG triplet is also important as it affects the fidelity and efficiency of translation initiation. For example, the scanning 40 S ribosome may occasionally initiate at the second AUG codon if the context of the first AUG is weak (Kozak, 1991). The Kozak sequence or consensus sequence for initiation of translation for filamentous fungi is CNNNCA(A/C)NATGGC (Jon et al. 1993). The surrounding nucleotides of the proposed new ATG codon in this study form a strong Kozak sequence of CACTCACAATGGC, whereas the context of the published ATG codon is weak with the Kozak sequence of GGACCAGAATGCT.

If the proposed additional third intron exists, the putative HAP1 binding site would be situated after the putative ATG start codon in the coding region of the gene. Consequently, this HAP1 site cannot be an upstream regulatory element for analysis! In

Figure 3.7 How the sequencing error affects the cycA aminoterminal sequence

The twelve amino acids from the amino-terminal of the predicted translated sequence is shown using the standard one-letter code. The top line represents the amino acid sequence produced from the published *cycA* gene (Raitt *et al.*, 1994). The bottom line shows how the correction for the two extra Thymidine bases affect the predicted amino acid sequence if the translational start site proposed by Raitt *et al.* (1994) was maintained. Homologous amino acid residues are indicated by the colour red. The invariant amino acid residue Gly 8 conserved in all cytochrome c proteins is shown in blue.

 $\frac{\text{Predicted}}{\text{amino-terminal sequence:}} + 1 \text{ M L I N C F F Q A T L P R V L } + 12$  after sequencing correction (6T)

addition, the 0.68 kb *cycA* promoter cloned into the R122 plasmid contains only a small portion of promoter sequence, and therefore this construct would be of no further use. Thus for the remainder of my project I endeavoured to confirm my suspicions about the presence of a third intron and thus an altered translation initiation site, and to re-examine the 'new' *cycA* promoter region.

# CHAPTER 4 RT-PCR ANALYSIS OF cycA RNA

#### 4.1 INTRODUCTION

RT-PCR analysis was performed to determine if the *cycA* gene contains two or three introns and therefore the ATG translational start codon proposed from this study (see Section 3.7), located upstream of the putative intron (position -402nt, Figure 3.6). The precise location of the introns was confirmed by sequencing the *A. nidulans* cDNA products over the intron/exon boundaries.

The transcriptional start site for two introns was deduced from primer extension analysis by Raitt *et al.* (1994). If there are three introns in the *cycA* gene as proposed in this study, a revised transcriptional start site can be calculated. The transcription start site proposed by Raitt *et al.* (1994), was deduced by primer extension analysis and does not account for the length of the third putative intron, which has been spliced from the RNA. Therefore the position of the revised transcriptional start site would be moved upstream by the length of the additional intron (395 bp) from the transcription start site calculated for two introns (Figure 4.1). Confirmation of the transcriptional start site is presently being determined in this laboratory.


#### 4.2 DEVELOPMENT OF A RT-PCR STRATEGY

mRNA is reverse transcribed into single stranded cDNA by the reverse transcriptase enzyme. Since introns are spliced from the mRNA, the cDNA strand synthesised will contain only exon information and 5' and 3' non-translated regions. To determine whether the *cycA* gene contains two or three introns, PCR primers were used which would produce a PCR product from the cDNA strand if there were three introns as in the theory outlined above, but no PCR product if there were two introns as shown in the published *cycA* sequence (Table 4.1, Figure 4.2). If three introns were present in the *cycA* gene, primer pair RB1/LG1 would amplify the entire cDNA of 596 bp and primer pair RB1/RB2 would produce a 298 bp product (Table 4.1). However, if two introns were present, no PCR product would be expected with either of these primer combinations because the RB1 primer site is located upstream of the major transcriptional start site (deduced from primer extension analysis (Raitt *et al.* (1994)). Therefore, the RB1 primer site would not be included in the mRNA.

A third primer pair, RB3/LG1 was used as a negative control for the amplification of products from the cDNA. No RT-PCR product should be amplified from the cDNA for

### Figure 4.1 Re-estimation of the cycA transcription start point

How the transcriptional start site for a *cycA* gene with three introns was estimated is shown. Asterisks with 3 and 2 represent the major transcriptional start point for three or two introns respectively. The proposed additional intron (Intron I), is shown by a large box. The published two introns (Introns II and III) are represented by blue blocks. Thin black lines represent exons and 5' and 3' untranslated regions. The approximate position of the primer originally used for primer extension analysis (Raitt, 1992) is indicated and the length of the mRNA product produced from this primer is shown.



Approximate Scale
0 500 1000 bp

### Figure 4.2 RT-PCR outline showing the relative positions of primers

The position of the primers (RB1, RB3, RB2, and LG1) used for RT-PCR are shown with respect to Introns I, II and III. The proposed third intron (Intron I) is shown by a red block, whereas the two published introns (Introns II and III) are indicated by blue blocks. Thin black lines represent exons and 5' and 3' untranslated regions. Asterisks represents the major transcriptional start point for 3 or 2 introns respectively.




Table 4.1 PCR Products Expected From RT-PCR Analysis

	RT-PCR Products Expected From:											
Primer Pairs	Splicing of 3 introns (bp)	Splicing of 2 introns (bp)	DNA template (bp)									
RB1-LG1	596	0	1100									
RB1-RB2	298	0	691									
RB3-LG1	0	0	860									

either two or three introns, since the RB3 primer site would fall within the proposed intron region for the three intron model, or would lie upstream of the transcriptional start point for the two intron model, and therefore would not be transcribed into cDNA.

#### 4.3 RT-PCR RESULTS


RT-PCR products are shown in the gel photograph in Figure 4.3. The 596 bp and the 298 bp RT-PCR products expected from the splicing of three introns were successfully amplified with the RB1/LG1 primer pair (Lane 2) and RB1/RB2 primer pair (Lane 4) respectively, indicating that the number of introns present in the cycA gene is three, rather than the published two (Raitt et al., 1994). The 596 bp product contains the full length cDNA sequence of the published cycA gene, whereas the 298 bp product contains the upstream cDNA region of the cycA gene, including the region from which the third intron has been spliced. A non-specific product of approximately 240 bp was also amplified consistently with the RB1/LG1 primer pair (Lane 2). This amplification was probably as a result of the primers binding non-specifically to the cDNA, because the product was too small to represent amplification from either DNA or partially spliced cDNA. Although Raitt et al. (1994) found that A. nidulans contains only one cytochrome c gene, the 240 bp product may have been amplified from an undiscovered second A. nidulans cytochrome c gene. However, this was unlikely due to the result of recent work in this laboratory (Gardiner, unpublished) which confimed that A. nidulans contains only one cytochrome c gene.

The three sets of primer pairs (RB1/LG1, RB1/RB2, RB3/LG1) were also used with a DNA template as a positive control (from total RNA samples that were not DNaseI treated prior to RT-PCR). These samples contained enough residual DNA for PCR. Both the 596 bp and the 298 bp RT-PCR products were distinguishable from the DNA template amplified products of 1100 bp (Lane 3) and 691 bp (Lane 5) respectively. A product of 691 bp was also amplified from the DNaseI treated RT-PCR sample with primers RB1/RB2 (Lane 4), probably as a result of DNA amplification due to incomplete DNaseI treatment. In addition, with these primers a faint non-specific product of approximately 610 bp was also observed for both DNaseI and non-DNaseI treated templates (Lanes 4 and 5), and probably represents the primers binding non-specifically to the DNA. For primer pair RB3/LG1 (Lanes 6 and 7), no cDNA product was amplified. This was expected as the RB3 primer site is not transcribed into cDNA for PCR amplification. The expected DNA template derived product of 860 bp was, however, produced from PCR amplification with primer pair RB3/LG1 (Lane 6 and 7). A faint band of approximately 750 bp was also amplified with these primers for both the DNaseI and non-DNaseI treated templates (Lanes 6 and 7). This product may be produced from the non-specific

Figure 4.3 RT-PCR analysis of total RNA from A. nidulans strain A18

All reactions were amplified by PCR from reverse transcribed total RNA from A18. The templates for lanes 2, 4 and 6 were DNaseI treated before RT-PCR. The following PCR primer pairs were used. Lanes 2 and 3: RB1 and LG1. Lanes 4 and 5: RB1 and RB2. Lanes 6 and 7: RB3 and LG1. Lanes 1 and 8 contain the 1 kb ladder (Gibco BRL, Life Technologies), the size of the bands indicated are in bp.

#### 



1,636

1,018

344 298

201

amplification of DNA, or a product formed from the non-splicing of Intron I (to allow binding of the RB3 primer) with either the splicing of both Intron II and III (to give a 746 bp product) or the splicing of just Intron II (799 bp) or Intron III (807 bp). Another faint band of approximately 630 bp was also observed for the non-DNaseI treated sample with the RB3/LG1 primers, and was probably the result of non-specific amplification from the DNA template.

#### 4.3.1 Sequencing of RT-PCR Products

The reverse transcribed and PCR amplified products of 596 bp and 298 bp were sequenced directly using the Perkin-Elmer Amplicycle Kit (Section 2.10.2) with primers RB1, RB2, and LG1 (Table 2.2). The PCR products were gel-purified (Section 2.8.1) before sequencing to remove additional PCR products present from the PCR reaction. To obtain sufficient quantities of DNA for sequencing, the cDNA fragments were further amplified by PCR using the PCR cycle conditions for the amplification of the cycA promoter region (Section 2.7). Initial sequence obtained from the cDNA fragments was generally of poor quality and often contained termination bands across all four lanes. Optimisation of the PCR reaction was performed to increase the purity of the PCR generated cDNA fragments for sequencing. Optimisation was performed by increasing the primer annealing temperature, decreasing the number of PCR cycles from 30 to 25, decreasing DNA concentration and decreasing the denaturation, annealing and elongation times (see Section 2.7 for optimised PCR cycle conditions). After PCR optimisation, increased quantities of DNA were obtained, and reduced DNA smearing (from the amplification of non-specific products) during agarose electrophoresis, indicated better quality DNA. Single stranded sequence of the 596 bp cDNA fragment was subsequently obtained successfully (Figure 4.4). Sequence from over the intron/exon boundaries (Figure 4.4) demonstated that three introns were spliced from the cycA gene, confirming that the cycA gene contains a total of three introns. The corrected nucleotide sequence for the cycA gene, including introns, is presented in Figure 4.5.

### Figure 4.4 cDNA sequence of the 596bp RT-PCR product.

Nucleotides are numbered from the A of the proposed methionine translational start codon (+1). The predicted amino acid sequence is shown below the coding strand in blue using the standard one-letter code. The pink asterisks above the nucleotide sequence indicates where the introns (Intron I, II and III) were spliced.

			•									*							
AC:	CAC	TAA	GGC	TAA	GGG	CGG	TGA	CAG	CTA	CTC	TCC	TGG	CGA	CTC	TAC	CAA	GGG	TGC	TA
		M	A	K	G	G	D	S	Y	S	P	G	D	S	T	K	G	A	F
•								direction.									10/4/20		1272
GC.	CTI	'CGA	GAC	CCG	TTG	CAA	GCA	GTG	ALC: NO.	CAC	TGI	CGA	GAA	CGG	CGG	CGG	CCA	CAA	GG
L	E	E	T	R	C	K	Q	C	H	Т	V	E	N	G	G	G	H	K	V
CGG	GCCC	CAA	CCT	CCA	CGG	TCT	CTT	CTG	CCA	CAC	TGT	CGA	GAA	CGG	CGG	CGG	CCA	CAA	GC
G	P	N	L	Н	G	L	F	C	H	T	$\nabla$	E	N	G	G	G	H	K	7
CG	GCCC	CAA	CCT	CCA	ccc	mcm		ccc	ccc	ת מידי	CAC	TCC	TCA	GGC	TGG	AGG	CTA	ccc	CI
000	D	L'AL		H		1	E	COG	D	TAA	m	100	Ch	7	00	700	U	7	
-0	P	18	D	п	9	L	E	131	17	1	1	6	W	£1	G	G	1	A	
																*			
CAC	CCGA	TGC	CAA	CAA	GCA	GGC	CGA	CGT	CAC	CTG	GGA	CGA	GAA	CTC	TCT	GTT	CAA	GTA	CC
T	D	A	N	K	Q	A	D	V	T	W	D	E	N	S	L	F	$Y_{i}$	Y	1
TC	GAAA	ccc	CAA	GAA	GTA	רביד	ccc	TGG	TAC	CAA	САТ	GGC	dididididi	ccc	TGG	тСп -	CDD	CAA	CI
T C	N	D	K	K	V	T	D	6	THE	K	M	A	T I	C	0	1	T	U	GI
J.u.	51.90	.4.	23	-23	(34)	7	1	Q	1	54X	100	678.5	:4:	342	0	-144	13.	100	
								*		•									
CA	AGGA	GAG	GAA	CGA	TCT	CAT	CAC	CTA	CCT	CAA	GGA	GAG	CAC	TGC	TTA	AAT	CGT	T	
CH																			

#### Figure 4.5 The corrected nucleotide sequence of the cycA gene.

The nucleotide sequence of the cycA gene published by Raitt et al. (1994), has been presented here in its corrected form. Nucleotides are numbered from the A of the proposed methionine translational start residue (+1). The published translational start site (Raitt et al., 1994) is indicated by the colour pink. The major transcriptional start site is indicated by a red asterisk. Three intervening regions (Introns I, II and III) are displayed in blue with the corresponding splice consensus sequences underlined. The predicted amino acid sequence is shown below the coding strand using the standard one-letter code. The observed sequencing error is indicated in red. The putative HAP1 site proposed by Raitt (1992) is marked in green.

-247	TACTCTGAGCACAGTGCTGATCATCCTACCTCTGATTGGTCAATTCCCAGATCACGGG
-187	GTCGTCGGGGGCGCGACCAAGAAACCAGCTCTACAAATTCCCTCCAAGTTTTTTTCTTC
-127	CTTTGGCCAGTCCGCTTGACTTGAATTCTGTCTTTCATCTTCTCTGCTACATACA
-67	TTGTACTATACCACTTACCTCTTTACATAACCCTTTCTCTCTCTACCTCTTTATTTTTT
-7	ACTCACAATGGCTAAGGGCGGTGACAGCTACTCTCCTGGTAAGTAGTTTGAATTCATCT
	MAKGGDSYSPG
54	TTCGTTTCGGCTAGGCGCTCTCTGCTGGGGGTGATATTCACTCCACTGCGTGCATTGTC
114	AAGTGAACGATATGTGAGACACAGGCTGTGAATGATGTGTTGGGTCTGGTGAGAACATT
174	TCCGATCCAACACAAGCTCAAAGTTGCCCACTTCTGGATCGCCATTTGATCGCCAGCAC
234	ATACAATTTCTACTTCTATCGCGCGCTGGCAATCCTCACTTTGCAGCGGATGCTTTATTC
294	TTCATCGTCGCCACAATGACGAGCTTCGACGCTTACTGCTTCACGACCACTCTCAG
354	ATGCGCGAAGCCGAACTGGAAGAGCTGGAGAAAGAGAAAGCGGACCAGAATGCTAATCA
414	
	D S T K G A K L F E T R C K Q
474	TGCCACACTGTCGAGAACGGCGGCGGCCACAAGGTCGGCCCAACCTCCACGGTCTCTTCCC H T V E N G G G H K V G P N L H G L F
534	. GGCCGTAAGACTGGTCAGGCTGGAGGCTACGCCTACACCGATGCCAACAAGCAGGCCGA
BATAIT	G R K T G Q A G G Y A Y T D A N K Q A D
594	GTCACCTGGGACGAGAACTCTCTGGTACAATCCCATGACAGCTCTAACAGCTCTGGGCC
	V T W D E N S L
654	ATTGCTAACTTTCTTTCACCAACAGTTCAAGTACCTCGAGAACCCCAAGAAGTACATCCC F K Y L E N P K K Y I P
714	TGGTACCAAGATGGCTTTCGGTGGTCTCAAGAAGACCAAGGAGGAGGAACGATCTCATCACGGTGTACCAAGAAGACCAAGGAGAAGACGATCTCATCACGGTTACACGATCTCATCACGATACAAGAAGACCAAGGAGAAGAACGATCTCATCACACAAGAAGAACGATCTCATCACAAGAAGAACGATCTCATCACAAGAAGAACGATCTCATCAACAAGAAGAACGAAC
774	GTATGTAACGCTGCTTACCACGGATAGGGCACATAGGCTAACAGGATGCACAGCTACCTC
834	AAGGAGAGCACTGCTTAAATCGTTCGCGATTAGACGAGATAAACCGCCCCCTGGGATTA
894	GACGAGGCCTCTGGCTAGGTGACAGGCGGGTACTGTAACATTACACCTAGACCTGGTT
954	TGAAGGTCGTCGGGACATGGAGGATATTATAGATCTTGTTTCCTTCGCCATCCTTGTCT
1014	TATCTTATTCTTTCCTTTACCTTGACGAGTGTTTTTTCAGCTTTGTGGTACC

# CHAPTER 4 DISCUSSION

### 4.4 THE cycA GENE CONTAINS A THIRD INTRON AND HAS A CORRECTED N-TERMINAL REGION

RT-PCR was performed to determine if the *cycA* gene contained two or three introns. Amplified products of 596 bp and 298 bp, expected from the splicing of three introns were produced using two different pairs of primers. Subsequent sequencing of both these products confirmed that the *cycA* gene contained three introns instead of two (Raitt *et al.*, 1994). The new intervening sequence (Intron I, 395 nt) is positioned from +32 nt to +426 nt in Figure 4.5. A consequence of the detection of Intron I was that the N-terminal region of the published ORF was affected. The published ATG start codon was positioned within the new intron region and thus could not be the correct translational start point. An ATG start codon was proposed and is located upstream of Intron I at nt position +1, Figure 4.5.

The new ORF (Figure 4.5) encodes a polypeptide of 112 amino acid residues (instead of 109 as published by Raitt et al., 1994), and contains a new N-terminal region, with the first 6 amino acid residues of the published cycA protein replaced by a new set of 9 amino acid residues (Figure 4.6). The N-terminal region of the cycA protein (as proposed in this study from the new ATG) was aligned with those from other fungi (including the predicted cycA sequence published by Raitt et al. (1994)), the plant Arabidopsis thaliana, and animals (Figure 4.6). The cytochrome c N-terminal region appears to be highly conserved within closely related groups. For example, within the yeast group, cytochrome c from the three yeasts Candida albicans, Schwanniomyces occidentalis and Kluyveromyces lactis, have identical amino-terminal regions. Within the animal kingdom, the human, rat, chicken and mouse cytochrome c genes encode only one amino acid variation within this region. However, the cytochrome c genes of the yeast S. cerevisiae encode slightly different amino acids at the N-terminal when compared with the other yeasts. Although the new A. nidulans cycA N-terminal portion (this study) is not completely homologous with the filamentous fungus Neurospora crassa, it provides a better match than the published cycA protein (with only 4 from the 9 new amino acids not identical) to any other cytochrome c genes sequenced so far. It also has the greatest amino terminal conservation with N. crassa, and with the CYC7 gene of S. cerevisiae. Interestingly, the new cycA ORF contains the same number of amino acid residues as the CYC7 gene. Once the cytochrome c genes from other filamentous fungi are sequenced, a closer match may be obtained within this group.

#### Figure 4.6 Alignment of cytochrome c N-terminal protein sequences

The N-terminal cytochrome c protein sequences of several different organisms are aligned with respect to A. nidulans (as proposed in this study), using the standard one-letter code. Alignment at conserved glycine residues shows different lengths of N-terminal sequence. Amino acid residues identical to A. nidulans (this study) are indicated by green. Dashes represent gaps introduced into the sequence to allow for optimal alignment. The highly-conserved glycine residue is indicated in red.

### Filamentous Fungi

A. nidulans (this study) A. nidulans (Raitt et al. 1994) N. crassa			K - -	Μ	L	I	N	C	F		G	D	S	T	K	G	A . A .		•
Yeast																			
S. cerevisiae CYC7 S. cerevisiae CYC1 C. albicans S. occidentalis K. lactis S. pombe	M - - -	- -		- М М	M P P	T A A	EPPP	F Y Y	K E K	A K K	GGGG	S S S S	A E E E	K K K	K K K	G D G G	A. A. A. A.		•
Plant A. thaliana Animal	-	М	Q	V	A	D	Ι	S	L	Q	G	D	Α	K	K	G	A.	•	
Human/Mouse/Rat/Chicken (Somatic)	-	-	-	-	-	-	-	_	_	М	G	D	VI	Ε	K	G	K	• •	٠

### 4.5 ALL THREE INTRONS OF THE cycA GENE ARE LOCATED AT CONSERVED POSITIONS

In the cytochrome c genes of the higher eukaryotes, mouse (Limbach and Wu, 1985), chicken (Limbach and Wu, 1983), and rat (Scarpulla and Wu, 1981), the position of intervening sequences appear to be conserved, with all of the coding sequences interupted after Lysine 55. The corresponding position in the cycA gene is at Alanine residue 64, nine amino acids upstream from the A. nidulans Intron II region (Figure 4.5 and Figure 4.7). None of the yeast cytochrome c genes sequenced so far contain intervening sequences, although introns are present in other yeast mitochondrial respiratory-chain proteins, for example the cytochrome b protein of S. cerevisiae (Lazowska et al., 1988). Figure 4.7 shows the cycA protein sequence aligned with the cytochrome c proteins of N. crassa and A. thaliana. The third intron shown to be present in the cycA gene by RT-PCR and subsequent sequencing (termed Intron I) is located in the coding sequence after the first nucleotide in the triplet coding for the conserved glycine 10 residue (Figure 4.7). This intron is also found in an identical position in the N. crassa cytochrome c gene (Bottorff et al., 1994). Intron III from A. nidulans is also conserved, being at an identical position in the N. crassa cytochrome c gene (at Thymidine 105 in the cycA protein, Figure 4.7). The plant cytochrome c gene from Arabidopsis thaliana (Kemmerer et al., 1991a) was found to cluster with N. crassa and other yeasts rather than with other higher plants. In accordance with this observation by Kemmerer et al. (1991a) Intron II, from A. thaliana, interrupts the coding sequence after Lysine 73, at the same position as for Intron II from A. nidulans (Figure 4.7). However, a recent paper by Hudson (1995) suggests that the A. thaliana cytochrome c sequence may originate from contaminating fungal material present in plant tissue, and thus is really a fungal cytochrome c sequence. The only other cloned and sequenced plant cytochrome c gene is from Oryza sativa (Rice) (Kemmerer et al., 1991b), which contains two introns located in unique positions after Glycine 38 and Lysine 82 in the cycA protein sequence.

In summary, all three intron positions from the *cycA* gene are conserved, and are found at identical positions with either *N. crassa* or *A. thaliana* cytochrome *c* genes. The length of these conserved introns at identical positions was also found to be similar but not exact (as shown in Figure 4.7).

Figure 4.7 Alignment of three cytochrome c protein sequences showing intron positions

The alignment of cytochrome c protein sequences from Aspergillus nidulans, Neurospora crassa, and Arabidopsis thaliana are shown using the standard one-letter code for amino acid residues. Dashes represent gaps introduced into the sequence for optimal alignment. The glycine residue found to be conserved in all cytochrome c genes sequenced is indicated by red.

```
A. nidulans
            Met A K G G D S Y S P G (Intron I 395 bp) D S
                                                   12
N. crassa
            Met ----- G F S A G (Intron I 547 bp) D S
                                                   7
A. thaliana
             Met Q V A D I S L Q G ----- D A
                                                   11
13
    TKGAKLFE ----- TRCKQCHTVENG
                                                   32
    KKGANLFK-------TRCAQCHTLEEG
8
                                                   28
    KKGANLFK (Intron I 78 bp) TRCAQCHTLKAG
12
                                                   32
    G G H K V G P N L H G L F G R K T G Q A G G Y A Y T D A N
                                                   61
33
    G G N K I G P A L H G L F G R K T G S V D G Y A Y T D A N
29
                                                   57
    EGNKIGPELHGLFGRKTGSVAGYSYTDAN
33
                                                   61
62
    KQADVTWDENSL (Intron II 61 bp) FKYLENPK
                                                   81
    KQKGITWDENTL ----- FEYLENPK
58
                                                   77
62
    KQKGIEWKDDTL (Intron II 59 bp) FEYLENPK
                                                   81
    KYIPGTKMAFGGLKKTKERNDLIT (Intron III 105
82
    KYIPGTKMAFGGLKKDKDRNDIIT (Intron II 101
78
82
    K Y I P G T K M A F G G L K K P K D R N D L I T ------ 105
    53 bp) Y L K E S T A
106
                     112
102
    74 bp) F M K E A T A
                      108
     ---- F L E E E T K
106
                      112
```

# CHAPTER 5 CLONING AND ANALYSIS OF THE 5' REGION OF THE cycA GENE

#### 5.1 INTRODUCTION

The *cycA* gene as published by Raitt *et al.* (1994), was found from this study to have the major translational ATG start codon positioned incorrectly (see Chapters 3 and 4). Consequently, due to the new location of the ATG codon, and the detection of one extra intron (Intron I), the clone of the *cycA* gene in plasmid R41 (Table 2.1, Figure 4.5), only contains 247 bp of cloned and sequenced promoter region (instead of 649 bp as published). It was therefore of interest to obtain additional promoter sequence in which to look for interesting regulatory regions, such as the yeast HAP1 motif involved in oxygen regulation. Therefore, library screening was performed to enable the cloning and characterisation of the 5' region of the *cycA* gene.

#### 5.2 SCREENING OF AN A. NIDULANS LIBRARY

To isolate the 5' region of the *cycA* gene, 10 filters containing approximately 30,000 pfu from a λGEM-11 *A. nidulans* genomic amplified library (kindly donated by Michael Hynes, University of Melbourne), prepared from *E. coli* MH2088, were screened by plaque hybridisation (Section 2.15). A 298 bp *A. nidulans* cDNA *Bam*H1 fragment (Section 2.15.3) which contains 224 bp of the 5' region of the *cycA* gene was used as a probe. Hybridisation and washing conditions are as in Section 2.15.3.

Thirty-five positive plaques were detected from the primary screen, and five plaques were selected for second round screening. During second round screening four out of five plaques remained positive, and after third round screening three plaques remained positive. These were designated  $\lambda$ LM5,  $\lambda$ LM9 and  $\lambda$ LM19 (Table 2.1). DNA was prepared from these clones as described in Section 2.5.6.

### 5.3 SOUTHERN BLOTTING AND RESTRICTION MAPPING OF POSITIVE $\lambda$ CLONES

The three positive  $\lambda$  clones were digested (Section 2.6.1) with double and single combinations of EcoRI, HindIII, and Xho I. These enzymes were chosen as they cut once or twice within the published region of the cycA gene (Raitt et al., 1994). In addition, EcoRI does not cut within the  $\lambda$ GEM-11 arms, XhoI cuts cloned inserts out of the  $\lambda$ GEM-11 arms, and HindIII cuts within the right arm. Digested  $\lambda$  DNA was

separated on a 1% agarose gel (Section 2.6.2), and the sizes of fragments obtained are summarised in Table 5.1. Clones λLM5 and λLM9 produced identical restriction enzyme patterns when digested with single and double combinations of EcoRI. HindIII. and XhoI, suggesting that  $\lambda$ LM5 and  $\lambda$ LM9 were identically isolated  $\lambda$  clones. However, λLM19 contained only a few identical restriction fragments with λLM5 and λLM9 (see Table 5.1) using the same combinations of enzymes, and thus appears to be an overlapping clone. A Southern blot of the two different clones, \(\lambda LM9\) and \(\lambda LM19\), was carried out to determine which restriction fragment contained the 5' region of the cycA gene. This was performed by hybridising a blot of the gel containing λLM9 and λLM19 clones digested with single and double combinations of EcoRI, HindIII, and Xho I (Figure 5.1a), to the 298 bp cDNA probe containing 224 bp of 5' cycA region (Figure 5.1b). Results from the Southern blot enabled a restriction map of both  $\lambda$  clones to be obtained (Figure 5.2), and confirmed that  $\lambda$ LM9 and  $\lambda$ LM19 are overlapping clones at the 5' region, with \(\lambda LM19\) containing less of the 3' cycA region. These results indicated that the 2.1 kb EcoRI fragment present in λLM9 and λLM19 contained the 5' region of the cycA gene.

# 5.4 SEQUENCE ANALYSIS OF CLONED $\lambda$ DNA SHOWN TO HYBRIDISE TO THE cycA GENE

### 5.4.1 Subcloning and Sequencing of the 2.1 kb Fragment

To obtain additional promoter sequence of the *cycA* gene, the 2.1 kb *Eco*RI fragment that hybridised to the 298 bp cDNA probe in both  $\lambda$ LM9 and  $\lambda$ LM19 (Table 5.1, Figures 5.1a and 5.1b), was subcloned from  $\lambda$ LM9 into pUC18 to give plasmid R133 (Table 2.1).

DNA from plasmid R133 was prepared by the rapid boiling method (Section 2.5), and sequenced by the Amplicycle sequencing system (Section 2.10.2). pUC/M13 universal forward (Perkin Elmer), pUC/M13 universal reverse (Perkin Elmer), and LG2 primers (Table 2.2) were used to obtain initial insert sequence. This information was used to design other primers (LG3, LG4, LM5, LM6, LM7, CYC8, Table 2.2) for directed sequencing. Figure 5.3 shows the sequencing strategy used to obtain the nucleotide sequence of the 5' region of the *cycA* gene from the cloned 2.1 kb  $EcoRI \lambda$  fragment. The nucleotide sequence is shown in Figure 5.4. Assembly of contigs and analysis of the 5' region of the *cycA* gene was performed using Sequencer Version 2.0.6 (Gene Codes Corporation).

### Figure 5.1a Restriction digestion profiles of λLM9 and λLM19

1 kb Gibco BRL (Life Technologies) ladder (lane 1). Lanes 2-7 are digests of λLM9: *Eco*RI (lane 2), *Hin*dIII (lane 3), *Xho*I (lane 4), *Eco*RI and *Hin*dIII (lane 5), *Eco*RI and *Xho*I (lane 6), *Xho*I and *Hin*dIII (lane 7). Lanes 8-13 are digests of λLM19: *Eco*RI (lane 8), *Hin*dIII (lane 9), *Xho*I (lane 10), *Eco*RI and *Hin*dIII (lane 11), *Eco*RI and *Xho*I (lane 12), *Xho*I and *Hin*dIII (lane 13).

### Figure 5.1b Southern Blot of λLM9 and λLM19

Autoradiograph of a Southern blot of the gel shown above in Figure 5.1a, hybridised to [a-32P]dCTP labelled 298 bp cDNA *Bam*HI fragment. Lanes as in Figure 5.1a.

### 1 2 3 4 5 6 7 8 9 10 11 12 13


212

5.2, 5.0 4.3 3.5

2.0/1.9

1.6 1.4

0.9 0.8



### 3 4 5 6 7 8 9 10 11 12 13

212

5.2, 5.0 4.3 3.5

2.0/1.9

1.6 1.4

0.9 0.8




Figure 5.2 Restriction maps of  $\lambda$ LM9 and  $\lambda$ LM19 isolated from an A. nidulans genomic library that hybridised to the cycA gene

Restriction maps of two *cycA* clones from *A. nidulans* strain A18. Arrow between two fragments indicates that the position of these fragments may be the reverse of that shown.

_		λLN	119						1 kb	
-						λLM9				1.5
<del></del>					•					
<i>Eco</i> RI			1.1	2.1	3	.6				
HindIII			6.1			4.5				
XhoI	6.0			4.9	1.7	<→ 1.2	3.5	<	> 1.6	


Table 5.1 Restriction Enzyme Digestions of  $\lambda$  clones

λ Clone	Restriction digest	Fragment Size (KB)
λLM9	EcoRI	22.7, 17.5, 14.9, 3.6 <sup>a</sup> , 2.1 <sup>ab</sup> ,
	Hind $III$	1.1 <sup>b</sup> 20.0, 8.1, 4.5 <sup>a</sup> , 4.5 <sup>b</sup> , 1.2, 1.0,
	XhoI	0.8, 0.7 22.7 <sup>a</sup> , 10.8, 3.5, 1.7 <sup>b</sup> , 1.6, 1.2 <sup>b</sup>
	EcoRI and HindIII	20.0, 8.1, 4.5 <sup>b</sup> , 3.6 <sup>a</sup> , 1.9 <sup>b</sup> , 1.2, 1.1 <sup>b</sup> , 1.0, 0.8, 0.7
	EcoRI and Xho I	22.7, 18.6, 10.8, 3.5, 2.1ab, 1.7b 1.6, 1.2b, 1.1b, 0.7ab
	Xho I and HindIII	20.0, 6.4, 4.5, 1.7 <sup>b</sup> , 1.4, 1.2 <sup>b</sup> , 1.0 <sup>ab</sup> , 0.7, 0.6
λLM19	<i>Eco</i> RI	22.6, 15.7, 2.1 <sup>ab</sup> , 1.1 <sup>b</sup>
/CLIVII)	HindIII	22.7 <sup>a</sup> , 8.4, 6.1, 4.5
	XhoI	24.4, 20.0, 4.9 <sup>a</sup> , 1.7 <sup>b</sup> , 1.2 <sup>b</sup>
	$Eco{ m RI}$ and $Hin{ m dIII}$	22.7, 8.4, 4.5 <sup>b</sup> , 2.6, 2.1 <sup>a</sup> , 2.0, 1.2 <sup>b</sup>
	EcoRI and XhoI	24.2, 20.0, 14.9, 2.1 <sup>c</sup> , 1.9 <sup>b</sup> , 1.7 <sup>b</sup> , 1.2 <sup>b</sup> , 1.1 <sup>b</sup> , 0.7 <sup>ab</sup> , 0.6
	XhoI and HindIII	20.0, 8.4,4.6 <sup>c</sup> , 4.5, 3.6, 2.0, 1.7 <sup>b</sup> , 1.2 <sup>b</sup> , 1.0 <sup>ab</sup>


<sup>&</sup>lt;sup>a</sup> indicates strong hybridisation signal with 298 bp cDNA probe.

b indicates identical restriction enzyme fragments present in both λLM9 and λLM19.

<sup>&</sup>lt;sup>c</sup> indicates partial restriction fragment hybridised with 298 bp cDNA probe.



The sequencing stratergy used to obtain additional promoter sequence from the 2.1 kb *Eco*RI fragment is shown, primer name is indicated above each arrow.



### Figure 5.4 Nucleotide sequence of the 5' region of the cycA gene

The nucleotide sequence of the 5' region of the A. nidulans cycA gene is presented here. Nucleotides are numbered from the A of the proposed methionine translational start residue (+1) (this study, Chapter 4). The HAP1 consensus sequences are indicated by red characters, with nucleotides matching the CGG N3 TA N CGG N3 TA consensus underlined. The putative HAP2/3/4/5 binding site is shown in blue, TATA motifs are coloured green, and CREA sequences are indicated by pink.

1247		TCACATA	AGCTCCCAACC	CAGAAAGCA	GTTTGCGGGTAA	AT
						C 7
-1207	GAGTACGCACAAA	AGCAATCCAGACA	TGAATCCACCC	BACICGICAA	MAACCGAAACAI	GA
-1147	CCGTCCCCTCGGG	CGGGAACATATTC	GGGTACTTCTT	TTTTTGCCCG	CTCCGCCTCTTC	TT
		1962				
-1087	TCTCAGAGAACTT	GGGACCGGGGTAG'	TTAACGACTTI	PACCATTGCG	TGTTGGAACGAC	GC
1007	GGCGCGCAGGATG		• • • • • • • • • • • • • • • • • • • •		• CACCCCCCCAMM	mп
-102/	GGCGCGCAGGATG	1GAGGGCGGCGA.	AAIIIAICIGO			11
-967	GGCTGTTGTCACT	TGAGGAATTTGTT	GATGCGCAGCC	GTGCGTCGG	TGGGAGCCTGCC	GA
		186	*	*	*	
-907	TGCGGGAGGACCG	GGACAGGAATAGG.	ATTGCTCGTGC	CTGAGGGAAC	ACCTCGGGGAAG	GA
-847		GGCTGATGGGGGC	• •	• • • • • • • • • • • • • • • • • • • •	TOTTCCCTCCCA	thit.
-04/	GGAAIGGGGAAI		ATTCTGTACGE		·	11
-787	TCTCTATATGCTA	GCTTCTGGTCGCC	GGCATAACATI	TTTGGTGTTG	ATATAATCATGT	GA
		· •	•	*		
-727	CTTCTGCCGCCGG	GAATAAGGCATCA	AGGCATCAAGG	GCACAAACAC	ATTTTTTCTAAT	GG
-667		AGGCCACTTCGGA	· mmaccccccc		• •	TC.
-007	COGCIAMGGCATC	AGGCCACTICGGA	11AGGGC <u>CGG</u>	JONGHOCOGG	ANAMACT COCCA	10
-607	ACTAGGCGAATGA	AAGGATGCAGATT	TGTTATTACGO	GGAGGGCTA	CTCCGGCCTCCG	TA
	*	•	:•	*		
-547	GCCCACCGTTGCC	CATTCCCCGAGAC.	AGACAGTGCA0	GAGCTCCAAG	TAACCAGCGTCT	CI
-487	ATGCGCTGGAATG	AGGTCATGCCATG	ACGGCATGCAC	EA ATCACCAA		СT
407				·	·	01
-427	ATAGTAGTTAGGC	TCTATGATAGATA	GATGTCATAGA	AGGTGCATT	GTTGCTATCTAG	AG
	32					
-367	CTGCATAACTGAG	CCCTTAGACGTAG	TATATAGGATI	PACAATAGTC	TCTAAATAAAGC	TI
-307	CATCCAGCCAGGC	GTTATTTGCAGTG	ACCGATTCCT	GACCTCAGAC	CCGGCAGCGCCC	GT
A 75.05 A 1		* .		•		
-247	TACTCTGAGCACA	GTGCTGAATCATC	CTACCTCTGAT	TGGTCAATT	CCCAGATCACGG	GT
102						00
-187	GTCGTCGGGGGCG	CGACCAAGAAACC	AGCTCTACAAA	TTCCCTCCA	AGTTTTTTTTCTT	
-127	CTTTGGCCAGTCC	GCTTGACTTGAAT	TCTGTCTTTCA	TCTTCTCTG	CTACATACAACT	CI
PSACAR-						
-67	TTGTACTATACCA	CTTACCTCTTTAC	ATAACCCTTTC	CTCTCTACCT	CTTTATTTTTTA	TC
7	· · · · · · · · · · · · · · · · · · ·	A A CCCCCCCCACACA	· COMA COMO COM	200		
-7	Managara and San	AAGGGCGGTGACA K G G D S		G		
	1.1 1.7		1 5 1	O .		

### 5.4.2 Confirmation that the 2.1 kb Fragment Corresponds to Regions of the A. nidulans A18 Genome

To verify that the 2.1 kb EcoRI fragment present in both  $\lambda LM9$  and  $\lambda LM19$ , and isolated from  $\lambda LM9$ , truly represents the 5' region of the cycA gene, Southern blots were performed against A. nidulans A18 (Table 2.1) genomic DNA with the 2.1 kb EcoRI  $\lambda$  fragment as a probe (Figures 5.5a and 5.5b). Southern blot analysis of A. nidulans A18 genomic DNA digested with EcoRI, and both EcoRI and HindIII, revealed that the  $\lambda LM9$  2.1 kb EcoRI and the 1.9 kb EcoRI-HindIII fragments (which were shown to hybridise to the 298 bp cDNA probe in Figure 5.1b) were also present in the A. nidulans A18 genome, thus confirming that the sequenced 2.1 kb EcoRI fragment represents the 5' region of the cycA gene as found in the genome. Unfortunately partial fragments were also detected by Southern blotting with EcoRI, and both EcoRI and HindIII digests, but the resulting fragment sizes were found to match the expected sizes for partial fragments using the information obtained from the restriction map of  $\lambda LM9$  (Figure 5.2).

### 5.4.3 Analysis of the cycA promoter sequence

#### 5.4.3.1 Putative HAP1 UAS

The cycA upstream sequence was searched to detect sequences which were similar to the 'optimal' HAP1 binding site CGG N3 TA N CGG N3 TA (Ha et al., 1996). It was anticipated that only imperfect versions of the 'optimal' consensus sequence would be found in the cycA gene, like all other known HAP1 binding sites (Ha et al., 1996). The potential A. nidulans HAP1 sites were aligned with other known Yeast HAP1 binding sites (Figure 5.6). From this alignment, the HAP1 binding site at -634 nt from A. nidulans appears to best match the optimal sequence, however the other HAP1 sites cannot be disregarded (see Section 5.4.6 for discussion of the A. nidulans HAP1 regulatory sequences).

### 5.4.3.2 Putative HAP2/3/4/5 (Yeast CCAAT-binding Factor) UAS

The region upstream of the *cycA* gene was also screened for the HAP2/3/4/5 complex (or Yeast CCAAT-binding factor) regulatory sequence. In the yeast *S. cerevisae*, this complex of proteins binds CCAAT sequences and activates many genes required for respiration (Zitomer and Lowry, 1992 for a review). An *A. nidulans* sequence was located at position -446 (Figure 5.4) that was 100% identical to the HAP2/3/4/5 consensus binding site TNATTGGT (or ACCAATNA).

# Figure 5.5a Restriction digestions of A. nidulans A18 genomic DNA Lanes 2 and 3 contain restriction digestions of A. nidulans A18 genomic DNA: EcoRI (lane 2), EcoRI and HindIII (lane 3). Lane 1 contains λ DNA digested with EcoRI and

HindIII.

### Figure 5.5b Southern blot of A. nidulans A18 genomic DNA

Autoradiograph of a Southern blot of the gel shown above in Figure 5.5a, hybridised to [a-32P]dCTP labelled 2.1 kb EcoRI fragment from  $\lambda LM9$ : EcoRI (lane 1), EcoRI and HindIII (lane 2).

1 2 3

212

5.2, 5.0

4.3

3.5

2.0/1.9

1.6

1.4

0.9

1 2



3.5

2.1 1.9

Figure 5.6 Comparison of known yeast and putative A. nidulans HAP1 UASs.

The known (functional) HAP1 binding sites from yeast are aligned with the putative HAP1 binding sites from the *A. nidulans cycA* gene. Characters in red indicate nucleotides which match the conserved CGG triplets or TA repeats given in the 'optimal' HAP1 sequence.

### Known Yeast UASs of HAP1:

CYC1	TGGC	CGG	GGTTTA	CGG	ACGATGA
CYC7	CCCT	CGC	TATTAT	CGC	TATTAGC
CTT1	GGAA	TGG	AGATAA	CGG	AGGTTCT
CYB2	GGCA	AGG	AGATAT	CGG	CAGGCTT
CYT1	CCGC	CGG	AAATAC	CGG	CCGCCCA
CYT1 (reverse)	CGGC	CGG	TATTTC	CGG	CGGCCAA
KlCYC1 (reverse)	ATTT	CGG	GAACAT	CGG	TCAAGAC

### A. nidulans putative HAP1 UAS:

CYCA	(-634)	CCGC	CGG	GGAGAG	CGG	GAAAAGG
CYCA	(-669)	TAAT	GGC	CGCTAA	GGC	ATCAGGC
CYCA	(-905)	GATG	CGG	GAGGAC	CGG	GACAGGA
CYCA	(+3, original)	CAAT	GGC	TAAGGG	CGG	TGACAGC

'OPTIMAL' CGG NNNTAN CGG NNNTA

#### Genes with HAP1 sites:

CYC1: Iso-1-cytochrome c from S. cerevisiae (Pfeifer et al., 1987).

CYC7: Iso-2-cytochrome c from S. cerevisiae (Prezant et al., 1987).

CTT1: Catalase T from S. cerevisiae (Wrinkler et al., 1988).

CYB2: Cytochrome b2 from S. cerevisiae (Lodi and Guiard, 1991).

CYT1: Cytochrome c1 from S. cerevisiae (Schneider and Guarente, 1991).

KICYC1: Cytochrome c from K.lactis (Freire-Picos et al., 1995).

CYCA: Cytochrome c from A. nidulans (+3, original; Raitt, 1992)

#### 5.4.3.3 Putative CREA Binding Site

CREA protein (encoded by the *creA* gene) is a negative regulator involved in glucose repression in *A. nidulans* (Cubero and Scazzocchio, 1994). CREA is a GC box-binding protein (Ronne, 1995), and binds sequences with the consensus SYGGRG (Cubero and Scazzocchio, 1994). The binding ability of CREA is also context dependent, and seems to require an AT rich 5' context *in vitro* (Cubero and Scazzocchio, 1994) Three candidate CREA sites were found in the *cycA* gene at positions -561 (reverse and complement position), -635, and -1070 (Figure 5.4). Multiple CREA regulatory elements have been found in the promoters of other *A. nidulans* genes (Mathieu and Felenbok, 1994).

#### 5.4.3.4 Other Regulatory Elements.

Three possible TATA boxes were present in the upstream region of the *cycA* gene at -293, -319, -341 nt positions (Figure 5.4). No significant pyrimidine rich elements were found, although an 18 bp pyrimidine tract extending from -124 to -139 nt (Figure 5.4), was located downstream from the proposed transcripitonal start site (Section 4.1). Therefore, either this element has no significance, or the proposed major transcriptional start site has been calculated incorrectly. Confirmation of the *cycA* transcription start site is currently under study in this laboratory.

The following *cis*-elements were not found in the *cycA* gene; a heat shock protein element, an iron responsive element (IRE), and a Yeast Ras/cAMP responsive element (PDS).

# CHAPTER 5 DISCUSSION

An A. nidulans genomic library was successfully screened to obtain the 5' region of the cycA gene. The nucleotide sequence upstream of the cycA gene revealed a number of putative regulatory elements. The relevance of some of these cis regulatory elements will be discussed below.

#### 5.4.4 TATA Box

Three possible TATA boxes were present in the 5' region of the *cycA* gene at 64, 90, and 112 nucleotides upstream of the estimated major transcriptional start site (Figure 5.4). Usually, TATA consensus sequences are found within 30-120 bp of the major transcriptional start site in fungal genes (Punt and van den Hondel, 1992). All three putative TATA boxes present in the *A. nidulans* promoter are located within this range, and therefore any one of them might represent the TATA box.

However, not all filamentous fungal promoters contain TATA boxes and, when they are present, promoters usually contain only one copy (Punt and van den Hondel, 1992; Gurr et al., 1987). In contrast, in S. cerevisiae, multiple TATA boxes are often found at various positions (Gurr et al., 1987). This is true for the S. cerevisiae CYC1 gene, which contains three TATA boxes (Smith et al., 1979). The cycA gene of A. nidulans therefore might be unusual for a filamentous fungal gene and contain three functional TATA boxes, although this seems unlikely.

In filamentous fungi, no regulatory proteins have been identified which bind to the TATA elements, thus the regulatory role of these elements in fungal genes is uncertain. However, deletion analysis of A. nidulans trpC promoter has suggested that the TATA box may be important in transcription iniation (Hamer and Timberlake, 1987). Whether one or all three of the TATA like boxes present in the A. nidulans cycA promoter are functional, will need to be determined experimentally.

### 5.4.5 The AnCF (A. nidulans CCAAT binding Factor) complex may regulate the cycA gene

A CCAAT motif was found in the upstream region of the cycA gene at position -446 nt (Figure 5.4), which shows complete homology to the binding site of the S. cerevisae HAP2/3/4/5 complex (or yeast CCAAT-binding factor). In the yeasts S. cerevisiae and

S. pombe, this heteromeric complex is involved in the activation of respiratory genes, such as those encoding cytochrome c, in response to a non-fermentable carbon source.

Research into the regulation of the A. nidulans cycA gene has revealed that its expression is not repressed by glucose (Raitt, 1992; Raitt et al., 1994). Since the HAP2/3/4/5 complex is responsible for glucose repression, the homology to this regulatory site may only be fortuitous. However, this study in A. nidulans was not performed on a number of different strains to determine if A. nidulans is like the aerobic yeast K. lactis, whose cytochrome c gene (KlCYCI) was only glucose repressed in some strains (Freire-Picos et al., 1995; Mulder et al., 1995).

The A. nidulans ands gene, which encodes an acetamidase enzyme, is controlled by both carbon and nitrogen regulation, and contains a CCAAT sequence in the 5' regulatory region which is required for setting basal levels of expression (Bonnefoy et al., 1995; van Heeswijck and Hynes, 1991). A protein complex named AnCF (A. nidulans CCAAT binding Factor) was found to bind to the CCAAT sequence (van Heeswijck and Hynes, 1991). Recent results suggest that the AnCF complex is analogous to the yeast HAP2/3/4/5 complex (Bonnefoy et al., 1995; Papagiannopoulos et al., 1996a; Papagiannopoulos et al., 1996b). Firstly, the yeast complex was found to recognise and activate expression of the amds gene through the amds CCAAT sequence (Bonnefoy et al., 1995). Secondly, homologs of the S. cerevisiae HAP3 and HAP5 components of the yeast CCAAT-binding factor have been isolated from A. niduilans (Papagiannopoulos et al., 1996a; Papagiannopoulos et al., 1996b). Deletion of the A. nidulans HAP3 homolog (hapC), resulted in decreased levels of amds expression under carbon limiting conditions, suggesting that it may encode a component of the AnCF complex. The HAP5 homolog (hapE) has yet to be disrupted. Interestingly, sequences other than the CCAAT sequence in the amds 5' regulatory region were required to achieve a response to carbon limitation, suggesting a difference in the response of the HAP2/3/4/5 complex to glucose repression in A. nidulans compared to that seen in S. cerevisiae.

Research on the *K. lactis HAP2* and *HAP3* genes has suggested that these genes, in contrast to their *S. cerevisiae* counterparts, were not essential for the expression of the *KlCYC1* gene (Mulder *et al.*, 1994; NGuyen *et al.*, 1995). Thus, the equivalent of the *S. cerevisiae* HAP2/3/4/5 complex in *K. lactis* does not seem to be involved in the response to glucose repression, (which only occurs in some strains), but will probably play a more general role in gene regulation, as found for the *A. nidulans amds* gene. In addition, the human HAP2 homolog, CP1, in combination with related factors, binds to CCAAT sequences in promoter regions of human genes to play a constitutive and cell type specific role in gene expression (Becker *et al.*, 1991).

In summary, it appears that the yeast HAP2/3/4/5 complex has been conserved from humans (Becker et al., 1991) through to filamentous fungi (Bonnefoy et al., 1995; Papagiannopoulos et al., 1996a; Papagiannopoulos et al., 1996b), due to the discovery of an analogous complex in A. nidulans (AnCF complex). The AnCF complex and the CCAAT-binding complex in K. lactis and humans seems to differ from the S. cerevisiae HAP2/3/4/5 complex by having no role in the control of glucose repression. Instead, the AnCF complex is required to set the basal level of amds transcription. Thus, the cellular role of the CCAAT-binding transcription factors seems to have diverged through evolution. Due to the modular nature of the HAP proteins, it is possible that the loss of carbon control by the CCAAT binding factors, is through the evolution of the HAP4 subunit which confers the yeasts complex's response to glucose repression.

Regardless of whether the *cycA* gene is found to be regulated by carbon source in some strains of *A. nidulans*, if the AnCF complex acts via the CCAAT sequence, it will probably affect the expression of the *cycA* gene by setting a basal level of transcription. Verification of this will have to await further experimental studies on the *cycA* gene of *A. nidulans*.

### 5.4.6 The A. nidulans cycA gene may by regulated by a HAP1-like Protein.

Three candidate HAP1 binding sites were found in the *cycA* 5' region, and are aligned with the known yeast HAP1 UASs in Figure 5.6. Due to the diversity of natural HAP1 binding sites (Ha *et al.*, 1996), it is difficult to determine which one, if any, are the most likely HAP1 binding site(s) before performing further experiments. The original HAP1 site chosen at the start of this study (on the basis of available results by Pfiefer *et al.*, 1987), and later determined to be located in the coding region of the gene (Section 3.7), matches least well with the 'optimal' HAP1 consensus sequence. In the light of recent results by Ha *et al.* (1996), this putative HAP1 site would not have been chosen as a candidate sequence. The study by Ha *et al.* (1996) showed that HAP1 will only bind to direct CGG repeats with a 6 bp spacer. If the CGG repeats are not conserved (ie. are degenerate forms), the TA repeats positioned asymmetrically in the spacer region are then essential for HAP1 binding. Thus the original HAP1 site at position +3 (Figure 5.6) would not have been chosen primarily because it has an <u>inverted</u> repeat of the GC-rich triplets.

The A. nidulans candidate HAP1 site at -669 forms a strong HAP1 binding site, it has a direct repeat of GGC, and has the TA sequence to stabilise protein binding. The other putative HAP1 site at -634 forms a possibly stronger HAP1 site, it has a direct repeat of

the optimal CGG triplet, but does not have the TA sequence, (which was determined by Ha et al. (1996) not to be necessary when the CGG triplet repeats were maintained). The sequence at -905 is virtually identical to the -634 site, but is missing one preferred A nucleotide when compared to the -634 site. However, the -905 site is situated 900 bp upstream from the ATG codon, whereas most regulatory binding sites are usually found closer to the ATG. Thus the -634 HAP1 site is proposed to be the most likely binding site for HAP1, but the possibility exists that both the -634 and -669 sites, only 35 bp apart, may both be HAP1 sites. Other regulatory proteins have been found to have two closely located sites which are both required, for example the A. nidulans CREA GC-rich binding protein (Cubero and Scazzocchio, 1994). In addition, only sites which have a 6 bp spacer between the CGG triplet repeats have been selected as putative HAP1 sites, but it is interesting to note that there were sequences present in the cycA promoter which had CGG repeats with a 5 bp spacer.

Since there are two HAP1 sites in the *cycA* gene which closely fit the HAP1 'optimal' consensus sequence, I speculate that a HAP1-like protein homologous to those which regulate oxygen transcriptional activation of many yeast respiratory genes, will probably also regulate the *A. nidulans cycA* gene. The functional significance of these sites in the regulation of the *cycA* gene is currently being investigated in this laboratory.

# CHAPTER 6 SUMMARY AND CONCLUSIONS

A promoter-reporter vector containing the A. nidulans cycA promoter fused to the lacZ gene of Escherichia coli was constructed to allow investigation of the role of upstream regulatory elements, primarly the putative HAP1 binding site, in the regulation of the A. nidulans cycA gene. However, sequencing of the cycA promoter in the new R122 reporter vector revealed a sequencing discrepancy when compared to the published cycA sequence (Raitt et al., 1994). Confirmation that the published sequence was incorrect was obtained by sequencing the R41 plasmid, which contains a sub-clone of the cycA gene from an A. nidulans genomic library. These results clearly confimed that six thymidine bases, rather than the published four, were present in the coding region of the cycA gene. Discovery of a sequencing error in the coding region of the cycA gene brought into question the validity of the open reading frames of the published cycA gene by Raitt et al. (1994). However, by comparison to other cytochrome c sequences it was considered that the deduced published amino acid sequence downstream from the sequencing error was correct. This implies that the published translation start codon was not correct, because it was no longer in the correct reading frame to produce the highly conserved cytochrome c protein. Further examination of the cycA gene showed that both the sequencing error and the published translation start codon may fall within a previously undetected intron region. To determine if an additional intron was present in the cycA gene, and thus provide further confirmation that the published translational start point was incorrect, RT-PCR analysis was performed on cycA RNA.

Two amplified products of 596 bp and 298 bp, indicative for the splicing of three introns, were produced by RT-PCR using two different pairs of primers. Subsequent sequencing of both these products proved that the *cycA* gene contained three introns instead of the two as published by Raitt *et al.* (1994).

Consequently, the published ATG initiation codon falls within the region of the previously undetected intron (Intron I). Therefore, a new translational start site was proposed, which contained a strong Kozak sequence for initiation of translation, and was the first AUG in the mRNA. Thus, the *cycA* gene has a corrected N-terminal region which was found to be very similar with the N-terminal region of the *S. cerevisae CYC7* gene. In addition, the position of the additional intron (Intron I) was conserved, being found at an identical position to that of an intron in the *Neurospora crassa* cytochrome *c* gene.

Due to the re-location of the translational ATG codon, the *cycA* promoter cloned into the reporter vector would now contain only a small portion of promoter sequence, and therefore this construct would be of no further use. In addition, the putative HAP1 binding site (Raitt, 1992) was now known to be situated in the coding region of the gene, and hence is not an upstream regulatory element for analysis.


To obtain the entire cycA promoter region for further analysis, an A. nidulans genomic library was screened with the 298 bp A. nidulans cDNA BamH1 fragment, which contains 224 bp of the 5' region of the cycA gene. Three cycA clones were isolated, and from restiction enzyme analysis of these three clones, the λLM9 clone was found to be identical to the λLM5 clone (since they contained the same restriction fragments), and the λLM9 and λLM19 clones were thought to be overlapping. Restriction enzyme and Southern blotting analysis of the two overlapping cycA clones, showed that a 2.1 kb EcoRI lambda fragment contained the 5' region of the cycA gene. This fragment was then subcloned from λLM9 into pUC18 to give plasmid R133. Sequencing of plasmid R133 (Table 2.1) enabled 1247 bp of cycA promoter sequence to be obtained. Analysis of this region revealed possible consensus sequences for the HAP1, AnCF complex, and the CREA transcription factors. In vitro and in vivo analysis of the cycA promoter is currently being performed in this labarotory to determine the significance of these sites. In vitro analysis will include gel mobility shift assays, where fragments of the cycA promoter will be tested with nuclear extracts from both A. nidulans and S. cerevisae. In vivo analysis in A. nidulans will be performed using reporter vector constructs containing either the entire cycA promoter, or the cycA promoter with appropriate deletions.

### **APPENDICES**

### Appendix 1.0 Vector Maps


Restriction map of pUC18 showing all restriction enzyme sites.

Plasmid R121 was created by insertion of the 0.68 kb *Bam*HI fragment of *A. nidulans* into the *Bam*HI site of pUC18 (Section 3.3.3). Plasmid R127 was constructed by insertion of the 2.1 kb *Eco*RI fragment of *A. nidulans* into the *Eco*RI site of pUC18 (Section 5.4.1).



Restriction map of pAN923-41B, 42B (R109) and 43B, showing sites for B, BamHI; Bg, BglII; E, EcoRI; H, HindIII; N, NruI; P, PstI; S, SalI; Sp, SphI; Ss, SstI; X, XhoI; Xb, XbaI.

Plasmids R120 (Section 3.3.1) and R122 (Section 3.3.4) were created by the insertion of the 0.68 kb *Bam*HI fragment of *A. nidulans* into the *Bam*HI site of pAN923-42B in the wrong and correct orientation respectively.




### Appendix 2.0 Sequence Data

### Summary of sequence obtained from the 2.1 kb EcoRI cycA fragment

<b>√</b> CYC8		TCACATATTA	GTCACATAGC	TCCCAACCCA	GAAAGCAGTT
CYC8				CCA	GAAAGCAGTT
LM5		TCACATATTA	GTCACATAGC	TCCCAACCCA	GAAAGCAGTT
1 LM5					GAAAGCAGTT
16.	1	114			
		TCACATATTA	GTCACATAGC	TCCCAACCCA	GAAAGCAGTT
1 CAC8		TGCGGGTAAA	TGAGTACGCA	CAAAAGCAAT	CCAGACATGA
CAC8		TGCGGGTAAA	TGAGTACGCA	CAAAAGCAAT	CCAGACATGA
1 LM5		TGCGGGTAAA	TGAGTACGCA	CAAAAGCAAT	CCAGACATGA
1 LM5		TGCGGGTAAA	TGAGTACGCA	CAAAAGCAAT	CCAGACATGA
	41				-
		TGCGGGTAAA	TGAGTACGCA	CAAAAGCAAT	CCAGACATGA
1 CAC8		ATCCACCGAC	TCGTCAAAAA	CCGAAACATG	AC
1 € CAC8		ATCCACCGAC	TCGTCAAAAA	CCGAAACATG	ACCGTCCCCT
1 LM5		ATCCACCGAC	TCGTCAAAAA	CCGAAACATG	AC
1 LM5		ATCCACCGAC	TCGTCAAAAA	CCGAAACATG	ACCGTCCCCT
(150 <del></del>	81				
		ATCCACCGAC	TCGTCAAAAA	CCGAAACATG	ACCGTCCCCT
NA-11-1-11-11-11-11-11-11-11-11-11-11-11-					
1 CAC8		CGGGCGGAA	CATATTCGGG	TACTTCTTTT	TTGCCCGCTC
1 LM5		CGGGCGGAA	CATATTCGGG	TACTTCTTT	TTGCCCGCTC
1 LM5		CGGGCGGGAA	CATATTCGGG	TACTTCTTTT	TTGCCCGCTC
	121				
		CGGGCGGAA	CATATTCGGG	TACTTCTTTT	TTGCCCGCTC
-					
<b>1</b> ☐ CYC8			TTCTCAGAGA		
1 LM5			TTCTCAGAGA		
1 LM5		CGCCTCTTCT	TTCTCAGAGA	ACTTGGGACC	GGGGTAGTTA
	161	CCCCMMMM	mmomonono	10mm000100	000000000000000000000000000000000000000
		CGCCTCTTCT	TTCTCAGAGA	ACTTGGGACC	GGGGTAGTTA
E cyce		ACGACTTTAC	CATTGCGTGT	TGGAACGACC	CGGCGCGCAG
□ CYC8			CATTGCGTGT		
1 LM5			CATTGCGTGT		
16	201				
		ACGACTTTAC	CATTGCGTGT	TGGAACGACG	CGGCGCGCAG

CAC	8	GATGTGAGGG	CGGGCGAAA		
LM7			CGAAAT	TTATCTGGTT	GCGCGAGGAC
1 LM5		GATGTGAGGG	CGGG		
1 LM5	0.44	GATGTGAGGG	CGGGCGAAAT	TTATCTGGTT	GCGCGAGGAC
	241	GATGTGAGGG	CGGGCGAAAT	TTATCTGGTT	GCGCGAGGAC
					occonomic
LM7		GCGCGGATTT	TGGCTGTTGT	CACTTGAGGA	ATTTGTTGAT
1 LMS		GCGCGGATTT	TGGCTGTTGT	CACTTGAGGA	ATTTGTTGAT
1G3		GCGGATTT	TGGCTGTTGT	CACTTGAGGA	ATTTGTTGAT
	281	Company (1964) (Company)			
		GCGCGGATTT	TGGCTGTTGT	CACTTGAGGA	ATTTGTTGAT
					<del>(1)</del>
1 LM7		GCGCAGCGGT	GCGTCGGTGG	GAGCCTGCCG	ATGCGGGAGG
1 LM5		GCGCAGCGGT	GCGTCGGTGG		
1 LG3		GCGCAGCGGT	GCGTCGGTGG	GAGCCTGCCG	ATGCGGGAGG
	321	** **********************			
		GCGCAGCGGT	GCGTCGGTGG	GAGCCTGCCG	ATGCGGGAGG
DIM		ACCCCCACAC	C	GCTCGTGCTG	200022000
18 TW		ACCOGGACAG	GARIAGGAII	GC1CG1GC1G	CACC
18 LG3		ACCGGGACAG	GAATAGGATT	GCTCGTGCTG	
1	361				
		ACCGGGACAG	GAATAGGATT	GCTCGTGCTG	AGGGAACACC
_					
1 IM		TCGGGGAAGG			
1 LM7		TCGGGGAAGG	AGGAATGGGG	GAATGGCTGA	TGGGGGCATT
1 □ LM6		mccccc a a cc	ACCA A MCCCCC	CAAMCCCMCA	GGGGGCATT
<b>4</b> □ LG3		TCGGGGAAGG		GAATGGCTGA GAATGGCTGA	
1 LG3	401		10000	GARIGGEIGR	IGGGGGCAII
	10,0	TCGGGGAAGG	AGGAATGGGG	GAATGGCTGA	TGGGGGCATT
200					
1 IM			TGCTGGTTCT		Education Laboratoria
1 IM				TGGCTGCGAT	
1 □ rc3				TGGCTGCGAT	
1 163				TGGCTGCGAT	
<b>1</b> □ LG3	441	CIGIACGAAI	IGCIGGIICI	TGGCTGCGAT	TICICIATAT
		CTGTACGAAT	TGCTGGTTCT	TGGCTGCGAT	TTCTCTATAT
0		0000000000	0.000.000.00		0.0000000000
1 LM7		GCTAGCTTCT	GGTCGCCGGC	ATAACATTT	
1 LM7		GCTAGCTTCT	GGTCGCCCCC	ATAACATTTT	GGTGTTGATA
TES TWO			GGTCGCCGGC		GGIGIIGAIA
4 EN LG3				ATAACATTTT	GGTGTTGATA
16 rc3				ATAACATTT	
16	481				



		4			
1 LIM6		TGCCCATTCC	CCGAGACAGA	CAGTGCAGAG	CTCCAAGTAA
1 LG2		TGCCCATTCC	CCGAGACAGA	CAGTGCAGAG	CTCCAAGTAA
<b>1</b> LG2			CGAGACAGA	CAGTGCAGAG	CTCCAAGTAA
LG2					AA
•••	721	4 ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) (		*****	_
		TGCCCATTCC	CCGAGACAGA	CAGTGCAGAG	CTCCAAGTAA
1 LM6		CCAGCGTCTC	TATGCGCTGG	AATGAGGTCA	TGCCATGACG
LG4					CCATGACG
1 LG2		CCAGCGTCTC	TATGCGCTGG		
1 LG2		CCAGCGTCTC	TATGCGCTGG	AATGAGGTCA	TGCCATGACG
1 LG2		CCAGCGTCTC	TATGCGCTGG	AATGAGGTCA	TGCCATGACG
16	761				
		CCAGCGTCTC	TATGCGCTGG	AATGAGGTCA	TGCCATGACG
		00000.0.0			
-FOI TME		GCATGCAGAA	TCACCAATC		
1 □ LM6			TCACCAATCA	TCTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT	m x m x c m x c m m
IG4			TCACCAATCA		TATAGTAGTT
<b>4</b> □ LG2					
1 LG2	001	GCATGCAGAA	TCACCAATCA	TCTTTTACAG	TATAGTAGTT
	801	001000000		mammma.a.a	
		GCATGCAGAA	TCACCAATCA	TCTTTTACAG	TATAGTAGTT
_					
IG4		AGGCTCTATG	ATAGATAGAT	GTCATAGAAG	GTGCATTGTT
<b>4</b> ☐ LG2		AGGCTCTATG	ATAGATAGAT	GTCATAGAAG	GTGCATTGTT
1 LG2		AGGCTC			
	841				
		AGGCTCTATG	ATAGATAGAT	GTCATAGAAG	GTGCATTGTT
JE LG4		GCTATCTAGA	GCTGCATAAC	TGAGCCCTTA	GACGTAGTAT
4€ LG2		GCTATCTAGA	GCTGCATAAC	TGAGCCCTTA	GACGTAGTAT
1 puc	Reverse	TAGA	GCTGCATAAC	TGAGCCCTTA	GACGTAGTAT
- ( <del></del> )	881				
		GCTATCTAGA	GCTGCATAAC	TGAGCCCTTA	GACGTAGTAT
LG4		ATAGGATTAC	AATAGTCTCT	AAATAAAGCT	TCATCCAGCC
1G2		ATAGGATTAC	AATAGTCTCT	AAATAAAGCT	TCATCCAGCC
	Reverse	ATAGGATTAC	AATAGTCTCT	AAATAAAGCT	TCATCCAGCC
.0	921				
		ATAGGATTAC	AATAGTCTCT	AAATAAAGCT	TCATCCAGCC
LG4		AGGCGTTATT	TGCAGTGACC	GATTCCTGAC	CTCAGACCCG
LG2		AGGCGTTATT	TGCAGTGACC	GATTCCTGAC	CTCAGA
	Reverse	AGGCGTTATT	TGCAGTGACC	GATTCCTGAC	CTCAGACCCG
16.	961				
		AGGCGTTATT	TGCAGTGACC	GATTCCTGAC	CTCAGACCCG

LG4		GCAGCGCCCG	TTACTCTGAG	CACAGTGCTG	AATCATCCTA .
IG4					TCATCCTA
1 puc	Reverse	GCAGCGCCCG	TTACTCTGAG	CACAGTGCTG	AATCATCCTA
	1001				
		GCAGCGCCCG	TTACTCTGAG	CACAGTGCTG	AATCATCCTA
LG4		cc			
LG4		CCTCTGATTG	GTCAATTCCC	AGATCACGGG	TGTCGTCGGG
1 puc	Reverse	CCTCTGATTG	GTCAATTCCC	AGATCACGGG	TGTCGTCGGG
	1041				*****
		CCTCTGATTG	GTCAATTCCC	AGATCACGGG	TGTCGTCGGG
J∰ LG4		GGCGCGACCA	AGAAACCAGC	TCTACAAATT	CCCTCCAAGT
	Reverse	GGCGCGACCA	AGAAACCAGC	TCTACAAATT	CCCTCCAAGT
		GGCGCGACCA	AGAAACCAGC	TCTACAAATT	CCCTCCAAGT
LG4		TTTTTTCTTC	CCTTTGGCCA	GTCCGCTTGA	CTTGAATTC
1 puc	Reverse 1121	TTTTTTCTTC	CCTTTGGCCA	GTCCGCTTGA	CTTGAATTC
		TTTTTTCTTC	CCTTTGGCCA	GTCCGCTTGA	CTTGAATTC

### REFERENCES

Abastado, J., Miller, P.F., Jackson, B.M. and Hinnebusch, A.G. (1990). Suppression of ribosomal reinitiation at upstream open reading frames in amino acid-starved cells forms the basis for GCN4 translational control. *Mol Cell Biol* 11: 486-496.

Amegadzie, B.Y., Zitomer, R.S., Hollenberg, C.P. (1990). Characterization of the cytochrome c gene from the starch-fermenting yeast Schwanniomyces occidentalis and its expression in Baker's yeast. Yeast 6:429-40.

Arst, H.N. and Scazzocchio, C. (1985). Formal genetics and molecular biology of the control of gene expression in *Aspergillus nidulans*. In: Bennet J.W., Lasure L.L. (eds). Gene Manipulation in fungi. Academic Press, New York, pp 306-343.

Becker, D.M., Fikes, J.D. and Guarente, L. (1991). A cDNA encoding a human CCAAT-binding protein cloned by functional complementation in yeast. *Proc Natl Acad Sci* 88: 1968-1972.

Bennet, B.L. and Molenaar, A.J. (1994). Cloning of PCR products can be inhibited by Taq Polymerase carryover. *Biotechniques* **16:** 32-36.

Bonnefoy, N., Copsey, J., Hynes, M.J., Davis, M.A. (1995). Yeast proteins can activate expression through regulatory sequences of the *amdS* gene of *Aspergillus nidulans*. *Mol Gen Genet* **246**: 223-227.

Borck, K., Beggs, J.D., Brammar, W.J., Hopkins, A.S., and Murray, N.E. (1976). The construction *in vitro* of transducing derivatives of phage lambda. *Mol Gen Genet* **146**: 199-207.

Bottorff, D. A., Parmaksizoglu, S., Lemire, E. G., Coffin, J. W., Bertrand. H., Nargang, F. E. (1994). Mutations in the structural gene for cytochrome c result in deficiency of both cytochromes aa3 and c in Neurospora crassa. Curr Genet 26: 329-335.

Bullock, W.O., Fernandez, J.M., and Short, J.M. (1987). XL-1-Blue: a high efficiency plasmid transforming *recA Escherichia coli* strain with beta-galactosidase selection. *Biotechniques* **5:** 376-378.

Cariello, N.F., Swenberg, J.A., and Skopek, T.R. (1991). Fidelity of *Thermococcus Litoralis* DNA polymerase (Vent) in PCR determined by denaturing gradient gel electrophoresis. *Nucleic Acids Res* **19:** 4193-4198.

Chodosh, L.A., Olesen, J., Baldwin, A. S., Guarente, L. and Sharp, P.A. (1988). A yeast and a human CCAAT-binding protein have homologous subunits that are functionally interchangeable. *Cell* 53: 25-35.

Clark, J. M. (1988). Novel non-templated nucleotide addition reactions catalyzed by procaryotic and eucaryotic DNA polymerases. *Nucleic Acids Res* **16:** 9677-9686.

Cohen, S.N., Chang, A.C.Y., Hsu, L. (1972). Non-chromosomal antibiotic resistance in bacteria: Genetic transformation of *E. coli* by R-factor DNA. *Proc Natl Acad Sci* 69: 2110.

Collett, M.A. (1995). Cloning, characterisation and evolutionary relationships of two *pyr4* genes from an *Acremonium* endophyte of perennial ryegrass. Ph.D. Thesis, Massey University, Palmerston North, New Zealand.

Creusot, F., Verdiere, J., Gaisne, M. and Slonimski, P.P. (1988). Regulator of oxygen-dependent gene expression in yeast. *J Mol Biol* **204**: 263-276.

Cubero, B. and Scazzocchio, C. (1994). Two different, adjacent and divergent zinc finger binding sites are necessary for CREA-mediated carbon catabolite repression in the proline gene cluster of Aspergillus nidulans. EMBO J 13: 407-415.

Davis, M.A., Hynes, M.J. (1991). Regulatory circuits in *Aspergillus nidulans*. In: Bennet J.W., Lasure L.L. (eds). More gene manipulations in fungi. Academic Press, San Diego, California, pp 151-189.

De Dekan, R.H. (1966). The Crabtree effect: a regulatory system in yeast. J Gen Microbiol 23: 149-156.

Dower, W.J., Miller, J.F., and Ragsdale, C.W. (1988) High efficiency transformation of *Escherichia coli* by high voltage electroporation. *Nucleic Acids Res* **16**: 6127-6145.

Forsburg, S.L. and Guarente, L. (1989a). Communication between mitochondria and the nucleus in regulation of cytochrome genes in the yeast *Saccharomyces cerevisiae*. *Ann Rev Cell Biol* **5:** 153-180.

Forsburg, S.L. and Guarente, L. (1989b). Identification and characterisation of HAP4: a third component of the CCAAT-bound HAP2/HAP3 heteromer. *Genes Dev* 3: 1166-1178.

Freire-Picos, M.A., Rodriguez Torres, A.M., Ramil, E., Cerdan, M.E., Breunig, K. D., Hollenberg, C.P., Zitomer, R. S. (1993). Sequence of a cytochrome c gene for *Kluveromyces lactis* and its upstream region. *Yeast* 9: 201-204.

Freire-Picos, M.A., Hollenberg, C.P., Breunig, K.D., Cerdan, M.E. (1995). Regulation of cytochrome c expression in the aerobic respiratory yeast *Kluyveromyces lactis*. Febs Lett 360: 39-42.

Frischauf, A., Lehrach, H., Poustka, A., and Murray, N. (1983). Lambda replacement vectors carrying polylinker sequences. *J Mol Biol* 170: 827-842.

Fytlovich, S., Gervais, M., Agrimonti, C. and Guiard, B. (1993). Evidence for an interaction between the CYP1 (HAP1) activator and a cellular factor during hemedependent transcriptional regulation in the yeast *Saccharomyces cerevisiae*. *EMBO J* 12: 1209-1218.

Glasel, J.A. (1995). Validity of nucleic acid purities monitored by 260nm/280nm absorbance ratios. *Biotechniques* 18: 62-63.

Guarente, L. and Mason, T. (1983). Heme Regulates Transcription of the CYC1 Gene of S. cerevisiae via an Upstream Activation Site. Cell 32: 1279-1286.

Guarente, L., Lalonde, B., Gifford, P., Alani, E. (1984). Distinctly regulated tandem upstream activation sites mediate catabolite repression of the *CYC1* gene of *S. cerevisiae*. *Cell* 36: 503-11

Gurr, S., Unkles, S., Kinghorn, J. (1987). The structure and organization of nuclear genes of filamentous fungi. Society of General Microbiology (special publication) Chapter 5: 93-139.

Ha, N., Hellauer, K. and Turcotte, B. (1996). Mutations in target DNA elements of yeast HAP1 modulate its transcriptional activity without affecting DNA binding. *Nucleic Acids Res* 24: 1453-1459.

Haldi, M. L. and Guarente (1995). Multiple domains mediate heme control of the yeast activator HAP1. *Mol Gen Genet* 248: 229-235.

Hamer, J.E. and Timberlake, W.E. (1987). Functional organisation of the Aspergillus nidulans trpC promoter. Mol Cell Biol 7: 2352-2359.

Hanahan, D. (1983) Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* **166**: 557-580.

Hudson, A. (1995). Fungal cytochrome c genes from plants. J Mol Evol 41:1170-1171.

Holmes, D.S. and Quigley, M. (1981). A rapid boiling method for the preparation of bacterial plasmids. *Anal Biochem* 114: 193-197.

Hooft van Huijsduijnen, R., Li, X.Y., Black, D., Matthes, H., Benoist, C. and Maithis, D. (1990). Co-evolution from yeast to mouse: cDNA cloning of the two NF-Y (CP1/CBF) subunits. *EMBO J* 9: 3119-3127.

Ish-Horowicz, D. and Burke, J.F. (1981). Rapid and efficient cosmid cloning. *Nucleic Acids Res* 9: 2989-2998.

Itakura, K., Rossi, J.J. and Wallace, R.B. (1994). Synthesis and use of synthetic oligonucleotides. *Ann Rev Biochem* 53: 323.

Jon J.P. Bruchez, Eberle, J. and Russo, V.E.A. (1993). Regulatory sequences involved in the translation of *Neurospora crassa* mRNA: Kozak sequences and stop codons. Fungal Genetics Newsletter, No. 40.

Kawasaki, E.S. (1990). Amplification of RNA. In: PCR Protocols: A Guide to Methods and Applications. Eds. Innis, M.A., Gelfand, D.H., Sninsky, J.J., White, T.J. San Diego, Academic Press, Inc. 21-27.

Karn, J., Matthes, H.W., Gait, M.J. and Brenner, S. (1984). A new selective phage cloning vector, lambda 2001, with sites for XbaI, BamHI, HindIII, EcoRI, SstI and XhoI. Gene 32: 217-224.

Kemmerer, E.C., Lei, M. and Wu, R. (1991a). Structure and molecular evolutionary analysis of a plant cytochrome c gene: Surprising implications for Arabidopsis thaliana. J Mol Evol 32: 227-237.

Kemmerer, E.C., Lei, M. and Wu, R. (1991b). Isolation and molecular evolutionary analysis of a cytochrome c gene from Oryza sativa (Rice). Mol Biol Evol 8: 212-226.

Kozak, M. (1991). Structural features in eukaryotic mRNAs that modulate the initiation of translation. *J Biol Chem* **266**: 19867-19870.

Lazowska, J., Claisse, M., Gargouri, A., Kotylak, Z., Spyridakis, A. and Slonimski, P. (1989). *J Mol Biol* 205: 275-289.

Lue, N.F., Chasman, D.I., Buchman, A.R. and Kornberg, R.D. (1987). Interaction of GAL4 and GAL80 gene regulatory proteins in vitro. Mol Cell Biol 7: 3446-3451.

Limbach, K.J. and Wu, R. (1983). Isolation and characterisation of two alleles of the chicken cytochrome c gene. *Nucleic Acids Res* 11: 8931-8950.

Limbach, K.J. and Wu, R. (1985). Characterisation of a mouse somatic cytochrome c gene and three cytochrome c pseudogenes. *Nucleic Acids Res* **13**: 617-630.

- Lodi, T. and Guiard, B. (1991). Complex transcriptional regulation of the Saccharomyces cerevisiae CYB2 gene encoding cytochrome  $b_2$ : CYP1 (HAP1) activator binds to the CYB2 upstream activation site UAS1-B2. Mol Cell Biol 11: 3762-3772.
- Maity, S.N., Vuorio, T. and de Crombrugghe, B. (1990). The B subunit of a rat heteromeric CCAAT-binding transcription factor shows striking sequence identity with the yeast HAP2 transcription factor. *Proc Natl Acad Sci* 87: 5378-5382.
- Maity, S. N., Sinha, S., Ruteshouser, E. C. and de Crombrugghe, B. (1992). Three different polypeptides are necessary for DNA binding of the mammalian heteromeric CCAAT binding factor. *J Biol Chem* **267**: 16574-16580.
- Marchuk, D., Drumm, M., Saulino, A. and Collins, F.S. (1990). Construction of T-vectors, a rapid and general system for direct cloning of unmodified PCR products. *Nucleic Acids Res* 19: 1154.
- Mathieu, M. and Felenbok, B. (1994). The Aspergillus nidulans CREA protein mediates glucose repression of the ethanol regulon at various levels through competition with the ALCR-specific transactivator. *EMBO J* 13: 4022-4027.
- McNabb, D. S., Xing, Y. and Guarente, L. (1995). Cloning of yeast *HAP5*: a novel subunit of a heterotrimeric complex required for CCAAT binding. *Genes Dev* 9: 47-58.
- Montgomery, D.L., Leung, D.W., Smith, M., Shalit, P., Faye, G. and Hall, B.D. (1980). Isolation and sequence of the gene for iso-2-cytochrome c in Saccharomyces cerevisiae. Proc Natl Acad Sci 77: 541-545.
- Mulder, W., Scholten, I.H.J.M., de Boer, R.W., Grivell, L.A. (1994). Sequence of the HAP3 transcription factor of *Kluyveromyces lactis* predicts the presence of a novel 4-cysteine zinc-finger motif. *Mol Gen Genet* **245**: 96-106.
- Mulder, W., Scholten, I.H.J.M., Grivell, L.A. (1995). Carbon catabolite regulation of transcription of nuclear genes coding for mitochondrial proteins in the yeast *Kluyveromyces lactis. Curr Genet* **28**: 267-273.
- NGuyen, C., Bolotin-Fukuhara, M., Wesolowski-Louvel, M., Fukuhara, H. (1995). The respiratory system of *Kluyveromyces lactis* escapes from HAP2 control. *Gene* **152**: 113-115
- Oechsner, U., Hermann, H., Zollner, A., Haid, A., Bandlow, W. (1992). Expression of yeast cytochrome c1 is controlled at the transcriptional level by glucose, oxygen and haem. *Mol Gen Genet* 232: 447-59.
- Olesen, J., Hahn, S., Guarente, L. (1987). Yeast HAP2 and HAP3 activators both bind to the CYC1 upstream activation site, UAS2, in an interdependent manner. Cell 51: 953-61.
- Olesen, J.T. and Guarente, L. (1990). The HAP2 subunit of yeast CCAAT transcriptional activator contains adjacent domains for subunit association and DNA recognition: model for the HAP2/3/4 complex. Genes Dev 4: 1714-1729.
- Olesen, J. T., Fikes, J. D. and Guarente, L. (1991). The Schizosaccharomyces pombe homolog of Saccharomyces cerevisiae HAP2 reveals selective and stringent conservation of the small core protein domain. Mol Cell Biol 11: 611-619.

Papagiannopoulos, P., Andrianopoulos, A, Sharp, J., Davis, M.A. and Hynes, M.J. (1996a). The *hap* genes of *Aspergillus nidulans* play a role in the regulation of *amdS* expression. Genetics Society of Australia, 43<sup>rd</sup> Annual Conference.

Papagiannopoulos, P., Andrianopoulos, A., Sharp, J.A., Davis, M.A. and Hynes, M.J. (1996b). The *hapC* gene of *Aspergillus nidulans* is involved in the expression of CCAAT-containing promoters. *Mol Gen Genet* 251: 412-421.

Parry, S. K. (1993). Construction and utilisation of a bidirectional reporter vector in the analysis of two *nod*-boxes in *Rhizobium loti*. Masters thesis, Massey University, Palmerston North, New Zealand.

Pfeifer, K., Prezant, T. and Guarente, L. (1987). Yeast HAP1 Activator Binds to Two Upstream Activation Sites of Different Sequence. *Cell* 49: 19-27.

Pfeifer, K., Kim, K., Kogan, S. and Guarente, L. (1989). Functional dissection and sequence of yeast HAP1 activator. *Cell* 56: 291-301

Pillar, T.M. and Bradshaw, R.E. (1991). Heat shock and stationary phase induce transcription of the *Saccharomyces cerevisiae* iso-2 cytochrome c gene. Curr Genet 20: 185-188.

Prezant, T., Pfeifer, K. and Guarente, L. (1987). Organisation of the regulatory region of the yeast *CYC7* gene: Multiple factors are involved in regulation. *Mol Cell Biol* 7: 3252-3259.

Punt. P. and van den Hondel, C. (1992). Analysis of transcription control sequences of fungal genes. In: (eds) Molecular Signals in Plant-Microbe Communications. CRC Press, Florida, Chapter 1, 29-48.

Punt, P.J., Dingemanse, M.A., Kuyvenhoven, A., Soede, R.D., Pouwels, P.H. and van den Hondel, C.A. (1990). Functional elements in the promoter region of the *Aspergillus nidulans gpdA* gene encoding glyceraldehyde-3-phosphate dehydrogenase. *Gene* 93: 101-9.

Raeder, U. and Broda, P. (1985). Rapid preparation of DNA from filamentous fungi. Lett Appl Microbiol 1: 17-20.

Raitt, D.R. (1992). The cytochrome c gene of Aspergillus nidulans. Ph.D. Thesis, Leicester Biocentre, United Kingdom.

Raitt, D.R., Bradshaw, R.E. and Pillar, T.M. (1994). Cloning and characterisation of the cytochrome c gene of Aspergillus nidulans. Mol Gen Genet 242: 17-22.

Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular Cloning: A laboratory Manual*. New York: Cold Spring Harbour Laboratory Press.

Sanger, F., Nicklen, S., and Coulson, A.R. (1977). DNA sequencing with chain terminating inhibitors. *Proc Natl Acad Sci* 74: 5463-5467.

Saunders, K. (1993). The regulation of the *Aspergillus nidulans* cytochrome c gene. Honours Thesis, Massey University, Palmerston North, New Zealand.

Scarpulla, R.C., Agne, K.M. and Wu, R. (1981). Isoaltion and structure of a rat cytochrome c gene. J Biol Chem 256: 6480-6486.

Schneider, J.C. and Guarente, L. (1991). Regulation of the yeast CYTI gene encoding cytochrome  $c_1$  by HAP1 and HAP2/3/4. *Mol Cell Biol* 11: 4934-4942.

Shih, C.N. and Marth, E.H. (1974). Aflatoxin formation, lipid synthesis, and glucose metabolism by *Aspergillus parasititicus* during incubation with and without agitation. *Biochem Biophys Acta* 338: 286-296.

Shizadegan, M., Christie, P. and Seemann, J.R. (1991). An efficient method for isolation of RNA from tissue cultured plant cells. *Nucleic Acids Res* 19: 6055.

Smith, M., Lueng, D.W., Gillam, S. and Astell, C.R. (1979). Sequence of the gene for iso-1-cytochrome c in *Saccharomyces cerevisiae*. Cell 16: 753-761.

Southern, E.M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* 98: 503-517.

Streatfield, S.J., Toews, S. and Roberts, C.F. (1992). Functional analysis of the expression of the 3'-phosphoglycerate kinase pgk gene in Aspergillus nidulans. Mol Gen Genet 223: 231-240.

Turcotte, B. and Guarente, L. (1992). HAP1 positive control mutants specific for one of two binding sites. *Genes Dev* **6:** 2001-2009.

Ushinsky, S.C. and Keng, T. (1994). A novel allele of HAP1 causes uninducible expression of HEM13 in *Saccharomyces cerevisiae*. *Genetics* **136**: 819-31

Van Gorcom, R.F.M., Punt, P.J., Pouwels, P.H., van den Hondel, C.A.M.J.J. (1986). A system for the analysis of expression signals in *Aspergillus*. *Mol Gen Genet* 223: 17-23.

Van Heeswijk, R. and Hynes, M.J. (1991). The *amdR* product and a CCAAT-binding factor bind to adjacent, possibly overlapping DNA sequences in the promoter region of the *Aspergillus nidulans amdS* gene. *Nucleic Acids Res* 19: 2655-2660.

Vashee, S., Xu, H., Johnston, S.A. and Kodadek, T. (1993). How do "Zn2 cys6" proteins distinguish between similar upstream activation sites? Comparison of the DNA-binding specificity of the GAL4 protein *in vitro* and *in vivo*. *J Biol Chem* **268**: 24699-24706.

Waldron, C. and Roberts, C.F. (1973). Cytoplasmic inheritance of a cold sensitive mutant in Aspergillus nidulans. J Gen Microbiol 78: 379-381.

Xing, Y., Fikes, J.D. and Guarente, L. (1993). Mutations in yeast HAP2/3 define a hybrid CCAAT box binding domain. *EMBO J12*: 4647-4655.

Xing, Y., Zhang, S., Olesen, J. T., Rich, A. and Guarente, L. (1994). Subunit interaction in the CCAAT-binding heteromeric complex is mediated by a very short α-helix in HAP2. *Proc Natl Acad Sci* **91:** 3009-3013.

Yanisch-Perron, C., Vieira, J. and Messing, J. (1985). Improved M13 phage cloning vectors and host strains: Nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* 33: 103-119.

Zhang, L., Bermingham-McDonogh, O., Turcotte, B. and Guarente, L. (1993). Antibody-promoted dimerisation bypasses the regulation of DNA binding by the heme domain of the yeast transcriptional activator HAP1. *Proc Natl Acad Sci USA* 90: 2851-2855.

Zhang, L. and Guarente, L. (1994a). The yeast activator HAP1-a GAL4 family memberbinds DNA in a directly repeated orientation. *Genes Dev* 8: 2110-2119.

Zhang, L. and Guarente, L. (1994b). HAP1 is nuclear but is bound to a cellular factor in the absence of heme. *J Bio Chem* **269**: 14643-14647.

Zitomer, R.S. and Lowry, C.V. (1992). Regulation of gene expression by oxygen in *Saccharomyces cerevisiae*. *Microbiol Rev* **56:** 1-11.