

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

RELATIONSHIPS BETWEEN HYDROGENATION AND HYDROLYSIS

OF DIETARY FAT IN THE BOVINE RUMEN

A thesis presented in partial fulfilment of the requirements  
for the degree of Master of Agricultural Science in  
Animal Science

by

William Ross SILCOCK

Massey University,  
New Zealand.

1968

## ACKNOWLEDGEMENTS

Throughout the course of this study the author has found the knowledge and experience of his supervisor, Dr. J.C. Hawke, to be invaluable in the development of ideas and experimental techniques, and expresses his gratitude for this help, readily given.

The author extends his thanks to Mr. G. Wilson and the staff of No. 2 dairy unit, who have not hesitated to provide the animals and facilities necessary for this study.

The technical assistance and helpful advice of Miss J. Barnett is acknowledged with thanks.

The very capable and rapid typing of Miss May Soo is gratefully acknowledged, as is the expert guidance received from Miss D. Scott of the Central Photographic Unit.

The author would finally like to record his thanks to Massey University and in particular the Chemistry and Biochemistry Department for making possible the circumstances under which this study was carried out.

## TABLE OF CONTENTS

<u>Chapter</u>		<u>Page</u>
1	INTRODUCTION	
1.1.	Introduction	1
1.2.	Hydrolysis of Glycerides in Rumen Fluid	5
1.3.	Hydrogenation of Unsaturated Fatty Acids in Rumen Fluid	7
1.4.	Factors Affecting the Degree of Hydrogenation of Unsaturated Fatty Acids of the Diet which takes place in the Rumen, with respect to the Composition of Depot and Milk Fats	14
2	THE AIM OF THE PRESENT STUDY	20
3	MATERIALS AND METHODS	
3.1.	Solvents and Reagents	22
3.2.	Incubation Procedures	
3.2.1.	Method	22
3.2.2.	Sampling	23
3.2.3.	Emulsification	23
3.3.	Synthesis of Radioactive Triglyceride	
3.3.1.	Synthesis of glyceryl 1-palmitate	24
3.3.2.	Synthesis of glyceryl 1-palmitate- 2-oleate and glyceryl 1-palmitate- 3-oleate	24
3.3.3.	Isolation of the pure Diglyceride Isomers	24
3.3.4.	Synthesis of glyceryl 1-palmitate- 2-oleate-3-(1- <sup>14</sup> C)-linolenate and glyceryl 1-palmitate-2-(1- <sup>14</sup> C)- linolenate-3-oleate	25
3.4.	Extraction and Methylation of Fatty Acids	
3.4.1.	Extraction of Lipid	25
3.4.2.	Saponification and Extraction of Fatty Acids	26

<u>Chapter</u>		<u>Page</u>
3	3.4.3. Methylation of Fatty Acids	26
	3.5. Thin Layer Chromatography	
	3.5.1. Preparation of Plates	27
	3.5.2. Elution Solvents	27
	3.5.3. Chromatographic Procedure	27
	3.5.4. Scanning of Thin Layer Chromatograms containing Radioactive Components	28
	3.6. Gas-liquid Chromatography	
	3.6.1. Apparatus	28
	3.6.2. Columns	29
	3.6.3. Operating Conditions	29
	3.6.4. Identification of Methyl Esters of Fatty Acids	30
	3.7. Radiochemical Methods	31
4	RESULTS	
	4.1. <u>In vivo</u> Hydrogenation of Dietary Lipid	32
	4.2. <u>In vitro</u> Hydrogenation of Dietary Lipid	33
	4.3. <u>In vitro</u> Hydrolysis of Triglyceride	34
	4.4. Relationships Between Hydrogenation and Hydrolysis of Unsaturated Triglyceride in Rumen Fluid	36
	4.5. Fatty Acid Composition of the Total Lipid after Incubation of Synthetic Triglyceride	38
	4.6. Fatty Acid Composition of the Glyceride and Free Fatty Acid Fractions after Incubation of Synthetic Triglyceride	39
5	DISCUSSION	
	5.1. Sampling of Rumen Contents	41
	5.2. <u>In vivo</u> Hydrogenation of Dietary Lipid	42
	5.3. Formation of Palmitic Acid in the Rumen	43
	5.4. <u>In vitro</u> Hydrogenation of Dietary Lipid	44

<u>Chapter</u>		<u>Page</u>
5	5.5. Hydrolysis of Triglyceride in Rumen Fluid	45
	5.6. Relationships Between Hydrogenation and Hydrolysis of Unsaturated Triglyceride in Rumen Fluid	50
	5.7. Fatty Acid Composition of the Total Lipid Obtained after Incubation with Rumen Fluid of Synthetic Triglyceride	54
	5.8. Fatty Acid Composition of the Glyceride and Free Fatty Acid Fractions Obtained after Incubation of Synthetic Triglyceride with Rumen Fluid	55
6	SUMMARY	59
	REFERENCES	61

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	The effect of triglyceride composition on degree of hydrolysis in rumen fluid	5
2	The fatty acid composition of the free fatty acid and "residual glyceride" fractions obtained following 32% hydrolysis of linseed oil in rumen fluid	6
3	Fatty acids formed following incubation of unsaturated fatty acids with sheep-rumen contents	F.P.8
4	The radioactive fatty acids obtained during incubation of U- <sup>14</sup> C-linolenic acid in rumen fluid	9
5	The products of oxidation of the non-conjugated acids obtained after incubation of U- <sup>14</sup> C-linolenic acid with rumen fluid	10
6	The products of oxidation of the octadecenoic acids obtained after incubation of U- <sup>14</sup> C-linolenic acid with rumen fluid	11
7	The effect of diet on Iodine Number of subcutaneous, visceral, and body fats of steer calves	15
8	Fatty acid composition of samples obtained during <u>in vivo</u> hydrogenation of dietary lipid	F.P.32
9	Comparison of the fatty acid composition of the residues and the fluid samples obtained by straining rumen contents (trial J)	F.P.33
10	Fatty acid composition of samples obtained during <u>in vivo</u> hydrogenation of dietary lipid (trial H)	F.P.34
11	Percentage distribution of radioactivity during hydrolysis by rumen fluid of synthetic radioactive triglyceride	F.P.35
12	Radioactive fatty acids obtained after incubation of synthetic triglyceride	F.P.36
13	Radioactive fatty acids from free fatty acid and glyceride fractions after incubation of glyceryl 1-palmitate-2-oleate-3-(1- <sup>14</sup> C)-linolenate (experiment 5)	F.P.37
14	Radioactive fatty acids of fractions isolated after reincubation of glyceryl 1-palmitate-2-oleate-3-(1- <sup>14</sup> C)-linolenate for 240 minutes	38

TablePage

15	Fatty acid composition of the total lipid extracted after incubation in rumen fluid of synthetic triglyceride	38a
16	Fatty acid composition of the glyceride and free fatty acid fractions obtained after incubation in rumen fluid of synthetic triglyceride (experiment 5)	39a

## LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1	Octadecadienoic acids formed after incubation of linolenic acid for 20 minutes with sheep-rumen contents (Ward <u>et al</u> , 1964)	F.P. 9
2	Pathways of hydrogenation of linolenic acid by rumen microorganisms (Wilde and Dawson, 1966)	F.P.12
3	Pathways of hydrolysis of glyceryl 1-palmitate-2-(U- <sup>14</sup> C)-linolenate-3-oleate in rumen fluid (experiment 1)	F.P.46
4.	Pathways of hydrolysis in rumen fluid of glycerides recovered from experiment 2 (experiment 3)	F.P.49
5	Scan of free fatty acid and glyceride fractions after reincubation of glyceryl 1-palmitate-2-oleate-3-(1- <sup>14</sup> C)-linolenate for 240 minutes	49a
6	Scan of free fatty acid and glyceride fractions after reincubation of glyceryl 1-palmitate-2-(1- <sup>14</sup> C)-linolenate-3-oleate for 240 minutes	49b
7	Scan of free fatty acid and glyceride fractions after reincubation of glyceryl 1-palmitate-2-(1- <sup>14</sup> C)-linolenate-3-oleate for 180 minutes	49c