

Etorphine induces pathophysiology in immobilized white rhinoceros through sympathomimesis that is attenuated by butorphanol

Jordyn M. Boesch^{1,*} , Robin D. Glead¹, Peter E. Buss^{2,3}, Adrian S. W. Tordiffe^{3,4} , Gareth E. Zeiler⁵, Michele A. Miller⁶, Francois Viljoen⁷, Brian H. Harvey^{7,8,9}, Stephen A. Parry¹⁰ and Leith C. R. Meyer^{3,4}

¹Department of Clinical Sciences, College of Veterinary Medicine, Cornell University, 930 Campus Road, Ithaca, New York 14853, USA

²Veterinary Wildlife Services, South African National Parks, Kruger National Park, Private Bag X402, Skukuza 1350, South Africa

³Centre for Veterinary Wildlife Research, Faculty of Veterinary Science, University of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa

⁴Department of Paraclinical Sciences, Faculty of Veterinary Science, University of Pretoria, Pathology and Veterinary Public Health Complex, Onderstepoort 0110, South Africa

⁵Department of Companion Animal Clinical Studies, Faculty of Veterinary Science, University of Pretoria, OVAH-building, Onderstepoort Campus, Private Bag X04, Onderstepoort 0110, South Africa

⁶South African Medical Research Council (MRC) Centre for Tuberculosis Research, Division of Molecular Biology and Human Genetics, Faculty of Medicine and Health Sciences, Stellenbosch University, PO Box 241, Cape Town 8000, South Africa

⁷Centre of Excellence for Pharmaceutical Sciences, School of Pharmacy (Division of Pharmacology), North-West University (Potchefstroom Campus), Universiteit Street, Potchefstroom 2531, South Africa

⁸MRC Unit on Risk and Resilience in Mental Disorders, Department of Psychiatry and Mental Health, Neuroscience Institute, Anzio Road, 1st floor, Groote Schuur Hospital, Observatory, Cape Town 7925, South Africa

⁹The Institute for Mental and Physical Health and Clinical Translation, Health and Education Research Building, Barwon Health, 299 Ryrie St, Geelong, Victoria 3220, Australia

¹⁰Cornell University Statistical Consulting Unit, Cornell University, Savage Hall, 244 Garden Avenue, Ithaca, New York 14853, USA

***Corresponding author:** Department of Clinical Sciences, College of Veterinary Medicine, Cornell University, 930 Campus Road, Ithaca, New York 14853, USA. E-mail: jmb264@cornell.edu.

White rhinoceros are a sentinel species for important ecosystems in southern Africa. Their conservation requires active management of their population, which, in turn, requires immobilization of individuals with an ultra-potent opioid such as etorphine. Unfortunately, when immobilized with etorphine, they develop severe hypoxaemia that may contribute to morbidity and mortality. We hypothesized that (i) etorphine causes sympathetic upregulation that is responsible for physiological complications that produce hypoxaemia and (ii) butorphanol, a partial μ opioid agonist, mitigates sympathetic upregulation, thereby improving arterial oxygen content (CaO_2) and delivery (DO_2). Six subadult male white rhinoceros were administered two treatments in random order: etorphine-saline (ES) and etorphine-butorphanol (EB). After intramuscular etorphine ($\sim 2.6 \mu\text{g kg}^{-1}$), rhinoceros became recumbent (time 0 min [t0]) and were instrumented. Baseline data were collected at t30, butorphanol (0.026 mg/kg) or 0.9% saline was administered intravenously at t37, and data were collected again at t40 and t50. At baseline, plasma noradrenaline concentration was $> 40 \text{ ng ml}^{-1}$, approximately twice that of non-immobilized rhinoceros (*t* test, $P < 0.05$); cardiac output (Q_t , by thermodilution) and metabolic rate (VO_2 , by spirometry/indirect calorimetry) were greater than predicted allometrically (*t* test, $P < 0.05$), and pulmonary hypertension was present. After butorphanol, noradrenaline concentration remained greater than in non-immobilized rhinoceros; in EB, CaO_2 was greater, while Q_t , DO_2 , VO_2 , and pulmonary pressures were less than in ES (linear mixed effect model, all $P < 0.05$). Increased noradrenaline concentration with increased Q_t and hypermetabolism supports etorphine-induced sympathetic upregulation. Butorphanol partly attenuated these effects, increasing CaO_2 but reducing Q_t and, thus, DO_2 . Since plasma noradrenaline concentration

remained increased after butorphanol administration while Q_t , DO_2 , and VO_2 decreased, a pathway independent of plasma noradrenaline concentration might contribute to the cardiopulmonary and hypermetabolic effects of etorphine. Developing treatments to combat this sympathomimesis could reduce capture-related morbidity in white rhinoceros.

Lay Summary

White rhinoceros are a sentinel species for ecosystems in southern Africa. Their conservation is dependent on occasional immobilization with the ultra-potent opioid etorphine. Unfortunately, etorphine produces sympathetic nervous system upregulation and severe physiological perturbations that may cause morbidity and mortality. Attenuating its effects contributes to white rhinoceros conservation.

Key words: Butorphanol, *Ceratotherium simum*, etorphine, hypoxaemia, noradrenaline, sympathetic

Abbreviations: IUCN: International Union for the Conservation of Nature, IM: intramuscular, VO_2 : metabolic rate or oxygen consumption, PaO_2 : arterial partial pressure of oxygen, $PaCO_2$: arterial partial pressure of carbon dioxide, $PA-aO_2$: alveolar-arterial oxygen partial pressure difference, Q_t : cardiac output, SNS: sympathetic nervous system, CNS: central nervous system, IV: intravenous, f_H : heart rate, mSAP: mean systemic arterial pressure, VCO_2 : carbon dioxide production, DO_2 : oxygen delivery to tissues, CaO_2 : arterial oxygen content, FiO_2 : fraction of inspired oxygen, ARRIVE: Animal Research: Reporting of In Vivo Experiments, KNP: Kruger National Park, ES: etorphine-saline group, EB: etorphine-butorphanol group, VWS: Veterinary Wildlife Services, LC-MS: liquid chromatography-mass spectrometry, SWG: standard wire gauge, SAP: systemic arterial blood pressure, PAC: pulmonary artery catheter, PAOP: pulmonary artery occlusion pressure, PAP: pulmonary artery pressure, PCV: packed cell volume, $PECO_2$: mixed expired carbon dioxide partial pressure, FEO_2 : mixed expired oxygen fraction, $PeCO_2$: end-tidal carbon dioxide partial pressure, $VEBTPS$: minute ventilation, at body temperature and pressure saturated, f_R : breathing rate, Hb: haemoglobin, USA: United States of America, ODC: oxyhaemoglobin dissociation curve, V_t : tidal volume, IQR: interquartile range, SD: standard deviation, ID: identification number, $P\bar{v}O_2$: mixed venous oxygen partial pressure, PVR: pulmonary vascular resistance, SV: stroke volume, OER: oxygen extraction ratio, $C\bar{v}O_2$: mixed venous oxygen content, V_{DPHYS} : physiological dead space ventilation, PVN: paraventricular nucleus, CNA: central noradrenergic, LC: locus coeruleus, rVLM: rostral ventrolateral medulla, PNMT: phenylethanolamine N-methyltransferase, D_2 : dopamine 2 receptor, CRF: corticotropin releasing factor, J_v : fluid flux, P_c : capillary hydrostatic pressure, K_f : reflection coefficient, Π_c : capillary oncotic pressure, EIPH: exercise-induced pulmonary haemorrhage, $V:Q$: ventilation to perfusion ratio, P_{50} : oxygen partial pressure at which 50% of haemoglobin is saturated,

Editor: Christine Madliger

Received 18 November 2024; Revised 19 January 2025; Editorial Decision 20 January 2025; Accepted 20 January 2025

Cite as: Boesch JM, Gleed RD, Buss PE, Tordiffe ASW, Zeiler GE, Miller MA, Viljoen F, Harvey BH, Parry SA, Meyer LCR (2025) Etorphine induces pathophysiology in immobilized white rhinoceros through sympathomimesis that is attenuated by butorphanol. *Conserv Physiol* 13(1): coaf009; doi:10.1093/conphys/coaf009.

(metabolic) acidaemia ensues (Nelson *et al.*, 1987; Dodd *et al.*, 1993). If oxygen debt is severe enough, cellular death occurs. Oxygen insufflation has been used to augment inspired oxygen fraction (FiO_2) in white rhinoceros during etorphine immobilization; however, this is often impractical in the field, and it can further decrease ventilation, presumably by opposing hypoxaemia-induced ventilatory drive (Haw *et al.*, 2015). Although beneficial effects of butorphanol on PaO_2 (and by inference, CaO_2) have been documented, and wildlife veterinarians presently operate under the assumption that increased PaO_2 indicates increased DO_2 in rhinoceros, the effect of butorphanol on DO_2 is undocumented (Buss *et al.*, 2018).

The primary objectives of our study were to determine if etorphine induces SNS upregulation in white rhinoceros and how this might lead to hypoxaemia and other physiological complications. Our secondary objective was to determine if butorphanol mitigates these perturbations and increases CaO_2 and DO_2 . We hypothesized that in white rhinoceros (i) etorphine causes sympathetic upregulation, including increased plasma catecholamine concentrations, which is responsible for cardiopulmonary and metabolic complications that produce hypoxaemia, and (ii) the partial μ opioid agonist butorphanol mitigates the sympathetic upregulation, thereby improving CaO_2 and DO_2 .

Materials and Methods

Animals

This research protocol was approved by the University of Pretoria Animal Ethics Committee (project number V101-15) and the South African National Parks (SANParks) Animal Use and Care Committee (reference number 001/16). It adheres to the Animal Research: Reporting of In Vivo Experiments (ARRIVE) guidelines (Percie du Sert *et al.*, 2020).

Six sub-adult (4–5 years old), male, white rhinoceros were used in the experiments. The immobilization and subsequent management of the rhinoceros followed the SANParks Standard Operating Procedures for the Capture, Transportation and Maintenance in Holding Facilities of Wildlife. The rhinoceros were captured via aerial darting in the Kruger National Park (KNP, 23°49'60 S, 31°30'0 E; altitude: ~317 m above sea level), South Africa. After capture, they underwent a physical examination to verify their health and were subsequently housed in groups of two or three at Veterinary Wildlife Services (VWS) bomas (dimensions: 10.5 × 21 m or 220.5 m²) for an habituation period of at least one month. Their diet consisted of lucerne (*Medicago sativa*) and tef (*Eragrostis tef*) hay and *ad libitum* water. Trained staff evaluated the rhinoceros daily using a scoring system developed by VWS based on the evaluation of: (i) feed intake, (ii) volume, consistency and colour of faeces, and (iii) behaviour (Miller *et al.*, 2016).

To guide the interpretation of experimental plasma catecholamine levels, venous blood samples were collected from

11 unmedicated, conscious white rhinoceros that were habituated to handling and permitted venipuncture without immobilization (i.e. 'control' white rhinoceros, Table S1). These samples were handled in the same way as the experimental samples and assayed together with them in one batch.

Study Design

A randomized, crossover study design was used. Blinding was not possible for logistical reasons. Two treatments, etorphine followed by 0.9% (isotonic) saline (ES) and etorphine followed by butorphanol (EB), were separated by a minimum washout period of 2 weeks. *A priori*, rhinoceros were assigned to treatment order by random allocation in blocks of two by lottery, and an online randomizer (www.randomiser.org) was used to determine the order of immobilization of the rhinoceros. All experiments were conducted in the morning between 6 and 9 AM local time from September to October. Ambient temperature and pressure were measured before each immobilization (Kestrel Instruments, Boothwyn, PA, USA).

Drugs were dosed based on estimated body weight (Table S2). However, body weight was measured at the end of each experiment, and drug doses, as well as physiological variables where appropriate (below), were normalized to actual body weight for reporting and analysis. Etorphine plus hyaluronidase was delivered IM in the nuchal hump by a 3-ml plastic dart with a 60-mm uncollared needle, fired from a compressed air rifle (DAN-INJECT International S.A., Skukuza, South Africa). Hyaluronidase is used routinely to increase the rate at which etorphine is absorbed from the site of injection, and systemic effects have not been reported for hyaluronidase during capture (Kock *et al.*, 1990). As soon as a rhinoceros stood still, VWS staff experienced in the handling of white rhinoceros determined if it was safe to approach; if so, the rhinoceros was blindfolded and positioned in sternal recumbency using ropes. Sternal recumbency was designated as time 0 (t_0); this was selected as t_0 based on the assumption that there was parity between the rhinoceros in the level of CNS depression at this time (Buss *et al.*, 2018).

As soon as a rhinoceros was positioned in sternal recumbency, shortened, lubricated equine endotracheal tubes (KRUSSE Silicone Endotracheal Tube, internal diameter: 28 mm, no. 282270, Jørgen Kruuse A/S, Langeskov, Denmark) were inserted into the nares, and their cuffs inflated to produce an airtight seal. The rhinoceros was then positioned in left lateral recumbency, and instrumentation continued as described below. Experience indicated that instrumentation could reliably be completed within 30 min after t_0 , hence the first (baseline) data were collected at t_{30} . At t_{37} , either butorphanol (Butonil, Wildlife Pharmaceuticals Pty Ltd, White River, South Africa) or an equivalent volume of 0.9% saline was injected into an auricular vein for treatments EB and ES, respectively. Data were collected again at t_{40} and t_{50} . Data collection was performed at t_{40} , or 3 min after injection of butorphanol (or saline), to document the early

effects of butorphanol. At the conclusion of the experiments, rhinoceros were de-instrumented. The rhinoceros in the EB group could be stimulated to stand and guided into a crate for weighing. The rhinoceros in the ES treatment group were given butorphanol IV so that they could be stimulated to stand and moved to the crate for weighing. After weighing, naltrexone (Kyron Laboratories Pty Ltd, Johannesburg, South Africa) was administered IV to fully antagonize the CNS effects of etorphine and butorphanol, and the rhinoceros were released back into the bomas. They were monitored continuously by trained VWS staff for the first hour after recovery and then hourly thereafter until they began eating, drinking, urinating and defaecating normally.

Instrumentation and Data Collection

Plasma Catecholamine Concentrations

At each data collection time, blood was collected from an auricular vein and added to tubes containing ethylenediamine-tetra-acetic acid anticoagulant; these were immediately put on ice and spun down in a refrigerated centrifuge at 2500 revolutions minute^{-1} for 10 min. Plasma was pipetted into cryotubes (Greiner Bio-One, Frickenhausen, Baden-Württemberg, Germany), snap-frozen in liquid nitrogen and stored at -80°C . This plasma was used to measure the concentrations of adrenaline, noradrenaline and dopamine with liquid chromatography-mass spectrometry (LC-MS, Dataset S1) (Xu *et al.*, 2018; Gao *et al.*, 2021).

Systemic Arterial Catheterization

After aseptic skin preparation, a 22 standard wire gauge (SWG) and 2.5-cm over-the-needle IV catheter (Nipro Medical Corporation, Bridgewater, NJ, USA) were inserted into a medial auricular artery for measurement of SAP at each data collection time. The catheter was connected by non-compliant tubing to a pressure transducer (Deltran II pressure transducer, DPT-200, Utah Medical Products Inc., Midvale, UT, USA) zeroed to atmospheric pressure at the level of the *manubrium sterni*; the entire system was filled with 0.9% saline. The transducer was connected to an amplifier (BP Amp, FE117, ADInstruments Pty Ltd., New South Wales, Australia) and a digital acquisition system (PowerLab 8/30, ADInstruments Pty Ltd).

Pulmonary Arterial Catheterization

A catheter introducer was placed aseptically into a linguofacial vein using a technique that was described previously (Boesch *et al.*, 2018). A custom-built, 200 cm, 7 Fr (2.33 mm), steel braid-reinforced, polyurethane Swan-Ganz-type (balloon-tipped) thermodilution pulmonary arterial catheter (PAC, Gaeltec Devices, Dunvegan, Isle of Skye, Scotland) was then inserted through the introducer into the jugular vein and passed into the right atrium, right ventricle and pulmonary artery (Swan *et al.*, 1970). The PAC contained two lumens, each with a diameter of 0.75 mm. The proximal

lumen was used to inflate a balloon located 3 cm from the PAC tip; the volume of air injected was sufficient to inflate the balloon to a diameter of 2–3 cm at ambient pressure. The distal lumen, which opened at the PAC tip, was connected to an arrangement similar to that used for measuring and recording SAP (above). Real-time observation of the blood pressure waveform allowed confirmation of the location of the tip of the PAC as it was introduced (Milne *et al.*, 1975). Once a pulmonary arterial pressure (PAP) waveform was observed, the PAC was advanced until the inflated balloon wedged in a branch of the pulmonary artery and a pulmonary arterial occlusion pressure (PAOP) waveform was observed on the monitor. Mean PAP (mPAP) and mean PAOP (mPAOP, recorded within 10 s of wedging) were measured at each data collection time. The balloon was then deflated, and the PAC was withdrawn slightly and left with the tip in the pulmonary artery so that it was ready for subsequent pressure measurements. The thermistor of the PAC was located just proximal to the balloon.

Cardiac Output

The PAC thermistor was connected to a cardiac output monitor (PM-9000 Vet Veterinary Portable Multi-Parameter Patient Monitor, ShenZhen Mindray Bio-Medical Electronics Co., Nanshan, ShenZhen, China). To calculate Q_t , the software in this device applied Stewart–Hamilton principles to the temperature–time curve generated by a bolus of ice-cold 0.9% saline that had been injected into the linguofacial vein introducer and transited the pulmonary artery. The software required manual input of a thermistor-specific computation constant. By design, the thermistor in the PAC had properties identical to those used in a commercial thermodilution catheter (ARROW Balloon Thermodilution catheter, Teleflex, Morrisville, NC, USA), allowing use of the computation constant for those commercial catheters. The thermistor was also used to measure pulmonary artery temperature (in $^{\circ}\text{C}$) prior to measurement of Q_t . For each Q_t measurement, 60 ml of ice-cold 0.9% saline was hand injected by the same investigator at end-expiration. Three to five sequential measurements within 20% of each other were recorded at each data collection time, and the median value was recorded for analysis.

Blood Gas and Acid-Base Analysis and Packed Cell Volume

Arterial and mixed venous blood samples were collected simultaneously from the systemic arterial catheter and the distal port of the PAC, respectively, at each data collection time. Blood samples (1 ml each) were collected anaerobically into 2-ml heparinized syringes, put on ice, and analyzed immediately with a point-of-care analyser (epoc Blood Analysis System, Siemens Medical Solutions, Inc., Malvern, PA, USA). Values for pH and blood gas partial pressures at 37°C were recorded. A portion of each sample was centrifuged at 15,000 g for 10 minutes to obtain packed cell volume (PCV).

Spirometry and Indirect Calorimetry

The nasal tubes described above were connected to a valve system that allowed inspiration of ambient air and directed all expired gas into a mixing chamber (MLA245, ADInstruments Pty Ltd.), where its temperature was measured (Thermistor Temperature Sensor, MLT415/M, ADInstruments Pty Ltd.), and then through a pneumotachometer (Respiratory Flow Head 1000L, MLT300L, ADInstruments Pty Ltd.) (Buss *et al.*, 2018). Expired gas was released into the atmosphere except for a period of 1 minute at the beginning of each of the three data collection times, when a Douglas bag was attached to the pneumotachometer for collection of the mixed expired gas (Buss *et al.*, 2018). Mixed expired carbon dioxide partial pressure (PECO₂, mm Hg) and mixed expired oxygen fraction (FEO₂, %) were measured in the Douglas bag (Cardiacap/5 monitor, Datex-Ohmeda, GE Healthcare, Helsinki, Finland). This device was also used to measure end-tidal carbon dioxide partial pressure (PECO₂) from a sampling port in one of the nasal tubes just outside the naris.

Tremor Score

Skeletal muscle activity was scored at each data collection time by a single experienced, unblinded observer using a previously published tremor scoring system (Table S3) (Buss *et al.*, 2018).

Spirometry and Blood Pressure Waveform Acquisition and Analysis

Spirometry and blood pressure waveforms were processed using dedicated hardware (PowerLab 8/30, ML870, ADInstruments Pty Ltd., New South Wales, Australia) and acquired digitally for subsequent analysis (LabChart 7 data acquisition software with Metabolic Module, version 7.3.8, ADInstruments Pty Ltd) on a laptop computer. Expired minute ventilation, corrected to pulmonary artery temperature and pressure, saturated with water vapor (VEBTPS) and respiratory rate (f_R) were measured from the expiratory flow waveform displayed by LabChart 7 (Boesch, 2020). The mPAP, mPAOP, mSAP and f_H were measured from the pulmonary and systemic arterial pressure waveforms also displayed by LabChart 7 (Boesch, 2020).

Calculations

Since a method for measuring haemoglobin concentration ([Hb]) was unavailable, we generated an equation for calculating [Hb] from PCV in the white rhinoceros. Between 2013 and 2017, 214 sub-adult, male white rhinoceros were captured in KNP. On a sample of blood from each animal, the PCV was measured as described above, and the cyanmethaemoglobin method was used to measure the corresponding [Hb] in an automated analyser (Vet abc, scil Animal Care Company, Gurnee, IL, USA) (Prakash and Banerji, 1972; Whitehead *et al.*, 2019). A regression line (described by the equation $y = mx + b$, with m = slope and b = Y intercept) for

PCV vs. [Hb] was generated using a commercial software package (GraphPad Prism 7, GraphPad Software Inc., San Diego, CA, USA). This line was described by the equation $y = 0.2528x + 4.013$; the value for r^2 was 0.535. This equation was used to calculate [Hb] from the measured PCV of the arterial and mixed venous samples.

Arterial and mixed venous oxyhaemoglobin saturations were calculated using an equation for the white rhinoceros-specific oxyhaemoglobin dissociation curve (ODC) (Reiners *et al.*, 2019). The equations for calculating other physiological variables are listed in Table S4. Allometric estimates for normal VO₂, Qt, f_H, VEBTPS, f_R, tidal volume (VT) and were calculated for each animal (Schmidt-Nielsen, 1991).

Statistics

Sample Size Calculation

Sample size was calculated *a priori* with a two-tailed t test of the difference in PaO₂ values between treatments in a similar crossover study in white rhinoceros (Buss *et al.*, 2018). Median PaO₂ in each treatment in the latter study was used in place of mean PaO₂, and interquartile range (IQR) in each treatment was used to calculate standard deviation (SD) (Wan *et al.*, 2014). Median and IQR values were entered into commercial software (G*Power 3.1.9.2, Kiel University, Kiel, Germany), with correlation between treatments set at 0.5. This produced an effect size estimate (Cohen's d_z) of 3.408. With $\alpha = 0.05$ and power (1- β) of 0.8, a sample size of three was needed. We used six rhinoceros in anticipation of loss of data for technical reasons.

Data Analysis

Data analysis was performed using commercial software (JMP 14.0.0, SAS Institute, Inc., Cary, NC, USA). No data were excluded from analysis. After performing a Shapiro-Wilk test to confirm normality of the data, a paired t test was used to compare ambient temperature and pressure, body weight of the rhinoceros, etorphine dose and time to sternal recumbency in the two treatments; alpha was set at 0.05. Residuals were calculated, and the Shapiro-Wilk test was used to confirm that the residuals were normally distributed.

For each of the physiological variables measured or calculated at t30, t40 or t50, a linear mixed effect model was constructed, with each variable designated as the response ('dependent') variable. Time, treatment (or 'phase'), and the interaction of time and treatment ('phase') were designated as fixed effects, and rhinoceros identification number (ID) and rhinoceros ID nested within treatment were designated as random effects. *A priori*, catecholamine concentrations, mPAP, mPAOP, Qt, PaO₂, mixed venous oxygen partial pressure (P \bar{v} O₂), DO₂ and VO₂, were designated as the ten primary outcomes. The remaining variables were designated as secondary outcomes. Alpha was set at 0.05 for the primary outcomes, and 0.05/19 = 0.0025 for the 19 secondary outcomes (a Bonferroni correction for multiple comparisons).

For each variable, the 36 residuals (three data collection times \times two treatments \times six rhinoceros) were calculated, and the Shapiro–Wilk test was used to confirm that the residuals were normally distributed. Residuals for pulmonary vascular resistance (PVR), stroke volume (SV), $PA-aO_2$ and f_R were not normally distributed; thus, the data for these variables were log-transformed, and residuals were re-tested to confirm normality. Residuals for log-transformed $PA-aO_2$ data were not normally distributed; however, the fit model was retained for this single variable.

A paired t test (two-tailed) was used to measure the significance of differences between measured VO_2 , Q_t , f_H , f_R , VE_{BTPS} and V_T at t30 for treatment ES and the corresponding values calculated allometrically. An unpaired t test (two-tailed) was used to measure the significance of differences in the plasma catecholamine concentrations between the ES experimental rhinoceros at t30 and the non-immobilized white rhinoceros. Alpha was set at 0.05. Data are expressed as mean \pm SD.

Results

During treatments ES and EB, respectively, ambient temperature was 22.4 ± 2.2 and $23.0 \pm 3.8^\circ C$, atmospheric pressure was 740.7 ± 4.1 and 742.1 ± 2.3 mm Hg, and body weight of the rhinoceros was 1145 ± 73 and 1152 ± 98 kg. Etorphine dose administered was 2.6 ± 0.2 and 2.6 ± 0.1 μg kg^{-1} . Time to sternal recumbency was 12.4 ± 2.0 and 11.7 ± 2.0 min. There were no significant differences between treatments for any of the above variables. No complications arose from PAC insertion, and the catheter was long enough to permit wedging in all rhinoceros at all data collection times. All rhinoceros recovered from the procedures and were subsequently released into the wild.

Changes in physiological variables over time within each treatment and between treatments are shown in Figs 1–6 and Table S5. At t30, before saline or butorphanol was administered, no significant differences were found between treatments ES and EB for any variable except noradrenaline concentration, which was significantly greater in the former at t30.

In the group of non-immobilized white rhinoceros, the plasma concentrations of adrenaline, noradrenaline and dopamine were 0.82 ± 0.01 , 23.92 ± 4.67 and 2.15 ± 0.03 ng ml^{-1} , respectively (Table S2). In these rhinoceros, one plasma noradrenaline concentration (11.28 ng ml^{-1}) was identified as an outlier; it was not excluded from the analysis since we had no explanation to support doing so.

Effects of Etorphine (Treatment ES)

Plasma Catecholamine Concentrations

Compared to the non-immobilized rhinoceros, plasma noradrenaline concentrations were greater, plasma dopamine




Figure 1: Graphs of plasma catecholamine concentrations measured in healthy, boma-habituated, sub-adult, male white rhinoceros ($n = 6$), assigned to two treatments administered once each in random order: etorphine-saline (ES, light blue circles) and etorphine-butorphanol (EB, dark blue circles). Blood was collected 30, 40 and 50 min after the rhinoceros became recumbent (t30, t40 and t50, respectively); either 0.9% saline (treatment ES) or butorphanol (treatment EB) were injected IV at t37. The central line within each data set represents the mean, whereas the error bars represent the SD. Refer to Figs 2–6 for graphs of other variables measured or calculated in the rhinoceros. A linear mixed effect model was constructed to assess changes in each of the 29 variables over time within treatments and between treatments. For the purposes of statistical analysis, the ten variables labelled with * on the Y-axis were considered primary outcomes; the 19 others were considered secondary outcomes. For the primary outcomes, alpha was set at 0.05; for the secondary outcomes, alpha was set at $0.05/19 = 0.0026$ (a Bonferroni correction for multiple comparisons). In all figures, a solid bar connecting data sets indicates a significant difference, and a dashed bar connecting data sets indicates a significant difference before Bonferroni correction. NB: Differences that lost significance after Bonferroni correction were not considered significant for the purposes of discussion in this paper. The blue arrowhead on the X-axis indicates when saline or butorphanol was administered at t37. The light blue band on the Y-axis of certain graphs indicates the 95% confidence interval for the catecholamine values measured in the non-immobilized rhinoceros or the allometrically calculated values for the etorphine-immobilized rhinoceros.




Figure 2: Graphs of physiological variables measured or calculated in healthy, boma-habituated, sub-adult, male white rhinoceros ($n = 6$), assigned to two treatments administered once each in random order: ES (light blue circles) and EB (dark blue circles). Data were collected 30, 40 and 50 min after the rhinoceros became recumbent (t30, t40 and t50, respectively); either 0.9% saline (treatment ES) or butorphanol (treatment EB) was injected IV at t37. Refer to Fig. 1 legend for further details. VO_2 , oxygen consumption; T_{PA} , pulmonary artery temperature; T_R , rectal temperature; OER, oxygen extraction ratio; VCO_2 , carbon dioxide production.

concentrations were less, and plasma adrenaline concentrations did not differ significantly. At t30 in treatment ES, the mean plasma noradrenaline concentration (49.25 ± 10.44 ng ml⁻¹) was approximately twice that observed in non-immobilized rhinoceroses ($P = 0.0012$). The mean plasma

dopamine concentration (1.28 ± 0.03 ng ml⁻¹) was lower than in the non-immobilized group ($P < 0.0001$), while plasma adrenaline concentrations (0.73 ± 0.19 ng ml⁻¹) did not differ significantly from those of non-immobilized rhinoceros ($P = 0.3050$).

Plasma adrenaline concentration was transiently less at t40, but plasma noradrenaline and dopamine concentrations did not change between t30 and t50 (Fig. 1, Table S5). Within treatment ES, plasma adrenaline concentration decreased between t30 and t40 ($P=0.0369$), but then rebounded between t40 and t50. Plasma noradrenaline and dopamine concentrations did not change significantly over time.

Allometric Comparisons

At t30, metabolic rate (oxygen consumption) and cardiac output were greater and ventilation was less than predicted allometrically. At t30, in treatment ES, VO_2 , Q_t , f_H and V_T were 129, 65, 195 and 92% greater, respectively, than predicted allometrically; f_R and VE_{BTPS} were 34 and 21% less, respectively, than predicted allometrically (Table 1).

Physiological Variables Over Time

At t40 and t50, values for most physiological variables were not different from values at t30. The exceptions were VO_2 (Fig. 2, Table S5), which was less at t40 ($P=0.0136$) and greater at t50 ($P=0.0082$); tremor score (Fig. 2, Table S5), which was less at both t40 and t50 ($P < 0.0001$ for both); and $P\bar{O}_2$ (Fig. 4, Table S5), which was higher at t50 ($P=0.0018$).

Effects of Butorphanol (Treatment EB) Over Time and Compared With Treatment ES

Catecholamine concentrations

Within treatment EB, the plasma catecholamine concentrations did not change significantly over time (Fig. 1, Table S5). At t40, plasma adrenaline concentration was greater in treatment EB than ES ($P=0.0366$). At t30, before butorphanol was administered, plasma noradrenaline concentration was less in treatment EB than ES ($P=0.0338$).

Metabolic variables

Butorphanol attenuated oxygen consumption and tremors. Within treatment EB, VO_2 (Fig. 2, Table S5) was less at both t40 and t50 than at t30 ($P < 0.0001$ for both). At t40 and t50, VO_2 was less in treatment EB than ES ($P=0.0009$ and 0.0029 , respectively). Within treatment EB, tremor score (Fig. 2, Table S5) was less at both t40 and t50 than at t30 ($P < 0.0001$ for both). Butorphanol reduced both pulmonary artery and rectal temperatures. Within treatment EB, pulmonary artery temperature (Fig. 2, Table S5) was less at both t40 and t50 than at t30 ($P < 0.0001$ for both), whereas rectal temperature (Table S5) was less at t50 than at t30 ($P < 0.0001$).

Plasma lactate concentration, oxygen extraction ratio (OER), and VCO_2 (Fig. 2, Table S5) did not change over time and were not different between treatments.

Cardiovascular Variables

Butorphanol reduced cardiac output and heart rate. Within treatment EB, Q_t (Fig. 3, Table S5) was less at both t40

Table 1: Values [mean \pm SD] for physiological variables measured in boma-habituated, subadult, male white rhinoceros ($n=6$) immobilized with etorphine (treatment ES) 30 min after sternal recumbency (t30) and corresponding normal values (mean \pm SD) calculated allometrically from body weight

	VO_2 (ml min ⁻¹ kg ⁻¹)	Q_t (ml min ⁻¹ kg ⁻¹)	f_H (beats min ⁻¹)	f_R (breaths min ⁻¹)	VE_{BTPS} (ml kg ⁻¹)	V_T (ml kg ⁻¹)
Mean \pm SD (measured at t30 in treatment ES)	4.8 \pm 0.89	81 \pm 26	121 \pm 21	5.6 \pm 2.3	100 \pm 20	19.6 \pm 7.0
Mean \pm SD (calculated allometrically)	2.1 \pm 0.03	49 \pm 0.61	42 \pm 0.82	8.6 \pm 0.16	127 \pm 2	10.2 \pm 0.0
P value	0.0007 ^a	0.027 ^a	0.0003 ^a	0.04 ^a	0.02 ^a	0.026 ^a

^aSignificant difference ($P < 0.05$) between measured and allometrically calculated values. VO_2 , oxygen consumption, standard temperature and pressure dry (STPD); Q_t , cardiac output; f_H , heart rate; f_R , breathing rate; VE_{BTPS} , expired minute ventilation, body temperature and pressure saturated with water vapour (BTPS); V_T , tidal volume.




Figure 3: Graphs of physiological variables measured or calculated in healthy, boma-habituated, sub-adult, male white rhinoceros ($n = 6$), assigned to two treatments administered once each in random order: ES (light blue circles) and EB (dark blue circles). Data were collected 30, 40 and 50 min after the rhinoceros became recumbent (t30, t40 and t50, respectively); either 0.9% saline (treatment ES) or butorphanol (treatment EB) was injected IV at t37. Refer to Fig. 1 legend for further details. Qt, cardiac output; f_H , heart rate; SV, stroke volume; mSAP, mean systemic arterial pressure.

and t50 than at t30 ($P = 0.0004$ and 0.0001 , respectively). At t40 and t50, Qt was less in treatment EB than in ES ($P < 0.0001$ for both). Within treatment EB, f_H (Fig. 3, Table S5) was less at both t40 and t50 than at t30 ($P < 0.0001$ for both).

Butorphanol increased arterial and mixed venous oxygen partial pressure and content. Within treatment EB, PaO_2 , $P\bar{v}O_2$, CaO_2 and mixed venous oxygen partial pressure ($C\bar{v}O_2$) (Fig. 4, Table S5) were greater at both t40 and t50 than at t30 (all $P < 0.0001$). At t40 and t50, all were greater in treatment EB than ES (all $P < 0.0001$).

Butorphanol reduced oxygen delivery to tissues. Within treatment EB, DO_2 (Fig. 4, Table S5) was less at t50 than at t30 ($P = 0.0091$). At both t40 and t50, DO_2 was less in treatment EB than ES ($P = 0.0173$ and 0.0015 , respectively). Blood haemoglobin concentration [Hb] (Fig. 4, Table S5) was less at t50 than at t30 ($P = 0.0003$).

Values for SV and mSAP (Fig. 3, Table S5) did not change over time and were not different between treatments.

Pulmonary variables

Butorphanol increased breathing rate and reduced $PaCO_2$. Within treatment EB, f_R (Fig. 5, Table S5) was greater at both t40 and t50 than at t30 ($P = 0.0001$ and 0.0003 , respectively). At t40, f_R was greater in treatment EB than ES ($P = 0.0012$). Within treatment EB, $PaCO_2$ (Fig. 5, Table S5) was less at

both t40 and t50 than at t30 ($P < 0.0001$ for both). At t40, $PaCO_2$ was less in treatment EB than ES ($P = 0.0005$) (Fig. 5, Table S5).

Butorphanol reduced pressure in the pulmonary vascular system. Within treatment EB, mPAP (Fig. 6, Table S5) was less at both t40 and t50 than at t30 ($P = 0.0007$ and 0.0001 , respectively). At t50, mPAP was less in treatment EB than ES ($P = 0.0153$). Within treatment EB, mPAOP (Fig. 6, Table S5) was less at both t40 and t50 than at t30 ($P = 0.0031$ and 0.0145 , respectively). At t40, mPAOP was less in treatment EB than ES ($P = 0.0458$).

The $PA-aO_2$, physiological dead space ventilation ($V_{D_{PHYS}}$), VE_{BTPS} , V_T (all Fig. 5, Table S5) and PVR (Fig. 6, Table S5) did not change over time and were not different between treatments.

Discussion

Previous studies suggested that assessing sympathetic activity and calculating DO_2 would help elucidate the effects of the ultra-potent opioid agonist etorphine with a view to identifying ways to minimize its adverse effects in white rhinoceros. This study measured plasma catecholamines and used modifications of established techniques to measure Qt, pulmonary arterial pressures, core (pulmonary artery) temperature and $P\bar{v}O_2$ for the first time in this species. Application of the




Figure 4: Graphs of physiological variables measured or calculated in healthy, boma-habituated, sub-adult, male white rhinoceros ($n = 6$), assigned to two treatments administered once each in random order: ES (light blue circles) and EB (dark blue circles). Data were collected 30, 40 and 50 min (min) after the rhinoceros became recumbent (t30, t40 and t50, respectively); either 0.9% saline (treatment ES) or butorphanol (treatment EB) were injected IV at t37. Refer to Fig. 1 legend for further details. PaO₂, arterial oxygen partial pressure; CaO₂, arterial oxygen content; PvO₂, mixed venous oxygen partial pressure; DO₂, oxygen delivery; CvO₂, mixed venous oxygen content; [Hb], haemoglobin concentration.

species-specific ODC allowed other variables to be derived for the first time, including CaO₂, CvO₂, DO₂ and OER (Reiners *et al.*, 2019). Butorphanol, a partial μ opioid agonist and κ opioid agonist, is used to mitigate the side effects of etorphine in white rhinoceros. Thus, this study aimed to investigate the mechanisms by which butorphanol might clarify the side effects of etorphine.

Effects of Etorphine in White Rhinoceros

As observed previously, etorphine-induced hypoxaemia, hypercapnia, tachycardia, systemic arterial hypertension, tremors, increased VO₂ and hyperthermia (Buss *et al.*, 2018).

Plasma Catecholamine Concentrations

Under etorphine, plasma noradrenaline concentration was greater than that measured in non-immobilized rhinoceros, and Qt, f_H and VO₂ were substantially greater than predicted allometrically (Schmidt-Nielsen, 1991). These observations are consistent with etorphine-induced SNS upregulation in white rhinoceros. Noradrenaline is the neurotransmitter of the SNS, but when a threshold plasma concentration is exceeded (1.8 ng ml⁻¹ in humans), it also acts as a hormone with important cardiopulmonary and metabolic effects that are exhibited during the fight-or-flight and stress responses, the extent of which are related to its plasma concentration (Silverberg *et al.*, 1978). Thus, it is likely that the high plasma




Figure 5: Graphs of physiological variables measured or calculated in healthy, boma-habituated, sub-adult, male white rhinoceros ($n = 6$), assigned to two treatments administered once each in random order: ES (light blue circles) and EB (dark blue circles). Data were collected 30, 40 and 50 min after the rhinoceros became recumbent (t30, t40 and t50, respectively); either 0.9% saline (treatment ES) or butorphanol (treatment EB) was injected IV at t37. Refer to Fig. 1 legend for further details. PaCO₂, arterial carbon dioxide partial pressure; f_R, respiratory rate; PA-aO₂, alveolar-arterial oxygen partial pressure difference; V_{DPHYS}, physiological dead space ventilation; V_{EBTPS}, minute ventilation, body temperature and pressure saturated with water vapour; V_T, tidal volume.

noradrenaline concentrations measured in this study contributed to the cardiopulmonary and metabolic side effects observed in etorphine-immobilized white rhinoceros.

The mechanism responsible for the etorphine-induced plasma noradrenaline concentrations is unclear. While hypoxaemia and hypercapnia can elevate noradrenaline in other mammals, their effect is minor compared to the levels seen in these rhinoceros (Rose *et al.*, 1983). Evidence from normoxic and normocapnic rats shows that opioids can induce catecholamine release and clinical signs of SNS activation, such as tachycardia and hypertension, suggesting that mechanisms other than hypoxaemia and hypercapnia may contribute to this response (Feuerstein and Siren, 1988).

Opioid-induced mechanisms, central and peripheral, that might explain sympathetic upregulation have been described in other species (Rose *et al.*, 1983; Chen *et al.*, 1999). In rats, β endorphins—endogenous opioid peptides that act as agonists at opioid receptors – increase central sympathetic outflow to the adrenal medulla and peripheral sympathetic nerve endings, thus stimulating peripheral catecholamine release and increasing plasma catecholamine concentrations (van Loon *et al.*, 1981). In humans, IV injection of both μ (fentanyl and morphine) and κ (nalbuphine) opioid receptor agonists induce significant, dose-dependent increases in plasma noradrenaline and adrenaline concentrations in healthy male individuals (Hoehe and Duka, 1993). Opioid receptors have been identified in the hypothalamus, the primary brain centre governing neuroendocrine and autonomic functions




Figure 6: Graphs of physiological variables measured or calculated in healthy, boma-habituated, sub-adult, male white rhinoceros ($n = 6$), assigned to two treatments administered once each in random order: ES (light blue circles) and EB (dark blue circles). Data were collected 30, 40 and 50 min after the rhinoceros became recumbent (t30, t40 and t50, respectively); either 0.9% saline (treatment ES) or butorphanol (treatment EB) were injected IV at t37. Refer to Fig. 1 legend for further details.

(Quirion *et al.*, 1983). Sympathetic outflow is increased by injection of low doses of μ opioid receptor agonists into nuclei of the hypothalamus, including the paraventricular nuclei (PVN), of unanaesthetized rats (Feuerstein *et al.*, 1983; Hassen *et al.*, 1984; Kiritsy-Roy *et al.*, 1986; Siren *et al.*, 1989; Siren and Feuerstein, 1991; Bachelard *et al.*, 1997). The hypothalamic PVN project directly to central noradrenergic (CNA) sites in the brainstem, including the locus ceruleus (LC) in the pons and the rostral ventrolateral medulla (rVLM) (Koba *et al.*, 2018). Etorphine's lipid solubility would allow it to cross the blood-brain barrier and reach these sites rapidly (Jolicœur *et al.*, 1992). In turn, these CNA sites project

to the cell bodies of the preganglionic SNS neurons in the intermediolateral cell column of the thoracolumbar spinal cord; after synapsing with post-ganglionic SNS neurons in various peripheral ganglia, post-ganglionic neurons terminate in organs throughout the body, notably the heart and adrenal medulla (Ross *et al.*, 1984; Strack *et al.*, 1989; Ally, 1998; Koba *et al.*, 2018).

The adrenal medulla is the principal source of plasma catecholamines (Kopin, 1968). All three opioid receptors (μ , κ and δ) have been identified in the adrenal medulla, with the concentration of each dependent on species (Dumont and Lemaire, 1984; Mansour *et al.*, 1986). Catecholamine synthesis occurs in the chromaffin cells of the medulla, beginning with the production of dopamine from dopa (Nagatsu *et al.*, 1964; Cahill *et al.*, 1996). Noradrenaline is then synthesized from dopamine by dopamine- β -hydroxylase (Levin *et al.*, 1960). Finally, adrenaline is synthesized from noradrenaline by phenylethanolamine N-methyltransferase (PNMT) (Cahill *et al.*, 1996). Catecholamines are stored in the chromaffin cells and secreted when tone increases in the SNS (Coupland, 1965; Cahill *et al.*, 1996). In the etorphine-immobilized rhinoceros at t30, compared to concentrations in non-immobilized rhinoceros, adrenaline concentration was similar, dopamine concentration was less, and noradrenaline concentration was greater. Differential secretion of catecholamines depending on which opioid is administered has been documented in people; in healthy male humans, fentanyl and nalbuphine increased plasma adrenaline and noradrenaline concentrations, while morphine increased noradrenaline only (Hoehe and Duka, 1993). This is consistent with our observations in white rhinoceros that adrenaline concentrations were unremarkable under etorphine, while noradrenaline concentrations were apparently increased. The explanation for differential secretion under μ opioid agonists may lie in the two populations of chromaffin cells that have been identified in the adrenal medulla: one in which 81% of the catecholamine content is adrenaline, and another where 75% of the catecholamine content is noradrenaline (Cahill *et al.*, 1996). The population of cells with the greater adrenaline content also has a greater concentration of the enzyme PNMT (Cahill *et al.*, 1996). This enzyme is induced by glucocorticoids; under chronic stress, glucocorticoids are transported from the adrenal cortex to the medulla in uniquely high concentrations by an intra-adrenal portal system, thus upregulating PNMT, increasing adrenaline synthesis, and thereby making more of it available for secretion (Wurtman, 2002). When stressors, and particularly chronic stress, are minimal, as we assume was the case with these boma-habituated white rhinoceros, it is possible that PNMT was not unduly induced, thus prioritizing noradrenaline secretion over adrenaline secretion during etorphine-induced sympathomimesis. It is interesting that white rhinoceros that were darted with etorphine in the field while being pursued by a helicopter had plasma concentrations of noradrenaline that are less than reported here and adrenaline concentrations that were higher

(de Lange *et al.*, 2017). The explanation for this difference might lie in the greater stress, which those rhinoceros were experiencing while being chased but could also be due to the tranquilizer azaperone, a dopamine receptor (D_2) antagonist and α_1 adrenoceptor antagonist, which was co-administered with etorphine to those rhinoceros (de Lange *et al.*, 2017).

Although these boma-habituated rhinoceros did not appear to be under stress, it is worth noting that all these samples were collected after at least one previous immobilization/capture with etorphine; half of the data was collected several weeks after a single injection of etorphine during capture in the wild, and the other half of the data was collected after a second etorphine injection during the second phase of the crossover study. The prior darting episodes might constitute an escalating allostatic load (defined as the cumulative effect of stress on the body caused by several attempts to adapt to stressors over time) that could have sensitized the rhinoceros, exaggerating normal bio-behavioural responses and generated a self-sustaining feed-forward loop involving outpouring of corticotropin-releasing hormone (CRF), a product of the hypothalamus, and noradrenaline (Elman and Borsook, 2019). However, inspection of plasma noradrenaline concentration in the rhinoceros after the first and second immobilizations showed that it was actually greater in five of six rhinoceros after the first rather than the second immobilization at t30. This seems not to support the theory of increasing allostatic load.

Metabolic Effects

Muscle Tremors. The tremors observed in etorphine-immobilized white rhinoceros might have been triggered by noradrenaline agonism of β_2 receptors in skeletal muscle (Marsden and Meadows, 1970; Kania, 1977). However, central mechanisms cannot be excluded, such as the antidopaminergic effect of etorphine in the CNS, which produces tremors in rats (Kania, 1977). In support of the latter mechanism, it should be noted that the rhinoceros under etorphine had lower plasma dopamine concentrations than the non-immobilized, 'control' cohort. It should also be noted that tremors decreased over time in the ES group while a corresponding decrease in plasma noradrenaline was not detected; this lends further support for a central mechanism, perhaps related to etorphine metabolism and decreasing CNS concentration. In any case, the skeletal muscle activity from the tremors would have contributed to the greater than predicted VO_2 and hyperthermia under etorphine.

Oxygen Extraction Ratio. The oxygen extraction ratio (OER) (or $VO_2 \div DO_2$) in these etorphine-immobilized rhinoceros was approximately twice the normal range of 20–30% reported in resting foals and horses, likely contributing to the observed low $\bar{C}\bar{v}O_2$ (Cambier *et al.*, 2008; Wong *et al.*, 2017). Greater than normal OER results from increasing VO_2 and/or decreasing DO_2 . In these rhinoceros immobilized with etorphine, as has been established previously, VO_2 is

two times greater than expected (Schmidt-Nielsen, 1991; Buss *et al.*, 2018). Decreased DO_2 , the product of CaO_2 and Qt , can result from decreased CaO_2 and/or decreased Qt . In these etorphine-immobilized rhinoceros, CaO_2 was substantially less than normal in other mammals, including domestic horses, while Qt was substantially greater than predicted allometrically; the increased Qt tends to cancel the decreased CaO_2 (Schmidt-Nielsen, 1991; Butler *et al.*, 1993). Because DO_2 has not been measured in rhinoceros that are not immobilized, and allometric values are unavailable for this variable in this species, our data do not allow us to confidently say if DO_2 is in fact decreased abnormally. Lactic acidemia ordinarily ensues if oxygen supply (i.e. DO_2) cannot meet oxygen demand (i.e. VO_2). Although the mean plasma lactate concentration in etorphine-immobilized rhinoceros was elevated compared to resting horses (2.0 vs. 0.5 mmol l^{-1}), it was lower than concentrations observed in other species under conditions of global tissue ischemia (Bayly *et al.*, 1983; McCoy *et al.*, 2011; Haas *et al.*, 2016). This finding adumbrates that the high Qt of these rhinoceros under etorphine might be lifesaving in the face of high VO_2 and low CaO_2 . Because the high Qt is caused by increased f_H , any therapy that reduces f_H (e.g. an α_2 adrenoceptor agonist sedative like detomidine) without decreasing VO_2 might exacerbate metabolic acidosis. The clinical implication of this is that co-administration of α_2 adrenoceptor agonist sedatives that cause bradycardia, e.g. detomidine or medetomidine, might be contraindicated when etorphine is used. It is also important to note that much greater plasma lactate concentrations are reported in rhinoceros darted with etorphine during field capture, where exertion and other stressors might exacerbate tissue hypoxia and thus induce anaerobic metabolism (Haw *et al.*, 2015; de Lange *et al.*, 2017).

Cardiopulmonary Effects

Hypoxaemia. Etorphine immobilization consistently resulted in severe, persistent hypoxaemia ($PaO_2 \sim 27$ mm Hg, $CaO_2 \sim 14$ ml dl^{-1} at t30). Possible causes of the hypoxaemia include reduced FiO_2 , hypoventilation and increased pulmonary venous admixture. The atmospheric pressure where the observations were made (~ 740 mm Hg) meant that inspired oxygen pressure was ~ 155 mm Hg ($FiO_2 \cong 21\%$), suggested that reduced FiO_2 would not have made an important contribution to alveolar or arterial hypoxia. Minute ventilation, however, was significantly less than predicted allometrically, and $PaCO_2$ was >80 mm Hg throughout immobilization, approaching 100 mm Hg in some rhinoceros. This confirmed that substantial alveolar hypoventilation was present during etorphine immobilization. Hypoventilation likely contributed to the hypoxaemia by increasing alveolar carbon dioxide, which displaced the other alveolar gases, including oxygen (Riley and Cournand, 1951). The observed $PA-aO_2$ values (>30 mm Hg) were much greater than values generally seen in resting, healthy mammals (<10 mm Hg), suggesting that venous admixture was present in the rhinoceros and that it made a substantial contribution to the

hypoxaemia (Hopkins *et al.*, 1994, 1998). The data suggest possible causes of venous admixture in these rhinoceros. The values for Q_t under etorphine were 65% greater than predicted allometrically, and the values for pulmonary vascular pressures exceeded those reported in resting horses and humans (Schmidt-Nielsen, 1991; Sinha *et al.*, 1996; Naeije and Chesler, 2012). Noradrenaline agonism of β_1 adrenoceptors in the sinoatrial node could explain the increased f_H , which increased pulmonary blood flow (i.e. right-sided Q_t). The increased pulmonary blood flow could have reduced erythrocyte transit time through the pulmonary capillaries, shortening the duration of erythrocyte exposure to alveolar gas and decreasing the time for oxygen to diffuse across the alveolar-capillary membrane and bind to Hb, resulting in abnormally low capillary blood oxygen content (Riley and Cournand, 1951). Interstitial pulmonary oedema increases the diffusion path for oxygen across the alveolar-capillary membrane and can occur when fluid flux (J_v) from the capillary lumen to the interstitial space increases (Levick and Michel, 2010). Clinically, the most common reasons for an increase in J_v are increased capillary hydrostatic pressure (P_c), damage to the alveolar-capillary membrane (which increases conductivity, K_f , and decreases the Staverman reflection coefficient, σ), or decreased capillary oncotic pressure (Π_c) (Starling, 1896; Murray, 2011). Although techniques for confirming the presence of interstitial pulmonary oedema were unavailable, if it had contributed to the hypoxemia, increased P_c or damage to the alveolar-capillary membrane would be the most likely mechanisms responsible for it. The mean pulmonary vascular pressures were so great in the etorphine-immobilized white rhinoceros that the rate of extravasation of water across the pulmonary capillaries would have increased and could have even caused stress failure of pulmonary capillaries, as has been described in horses during maximal exertion (exercise-induced pulmonary haemorrhage, or EIPH); this may explain unpublished reports of sanguineous froth observed at the nares of rhinoceros during immobilization with etorphine (West *et al.*, 1993). If the rate of water extravasation exceeded the lymphatic capacity to remove the water, interstitial pulmonary oedema would have occurred; such interstitial oedema would increase the barrier to diffusion of oxygen across the alveolar-capillary membrane and contribute to venous admixture (Starling, 1896; Riley and Cournand, 1951). Parts of the lung with lesser ventilation and/or greater perfusion also contribute to venous admixture; conventionally this is expressed as parts with ventilation:perfusion (V:Q) of <1 (Riley and Cournand, 1949). Electrical impedance tomography has demonstrated that the dependent lung is poorly ventilated in etorphine-immobilized, laterally recumbent white rhinoceros (Mosing *et al.*, 2020). Our data do not allow us to comment on the possible role of V:Q inequality in the venous admixture and hypoxemia observed in etorphine-immobilized rhinoceros. It is worth noting that the $\dot{C}\bar{V}O_2$ in etorphine-immobilized rhinoceros is ~ 5 ml dl⁻¹; this is much less than considered normal in horses in which it is ordinarily ~ 15 ml dl⁻¹ (Butler *et al.*, 1993). This would amplify the effects of all the

above causes for venous admixture and therefore contribute to arterial hypoxemia (Riley *et al.*, 1951).

Hypoventilation. The metabolic and cardiopulmonary responses to etorphine described above can largely be attributed to those expected from sympathetic upregulation. On the other hand, ventilation was apparently down-regulated; both f_R and VEBTPs were less than predicted allometrically (Schmidt-Nielsen, 1991). This finding is consistent with the depressant effects of μ opioid agonists on breathing pattern generation by the medullary pre-Bötzinger and pontine parabrachial complexes that have been described in other mammals at doses equal to or greater than those used to produce analgesia (Lalley, 2006; Prkic *et al.*, 2012; Miller *et al.*, 2017; Saunders and Levitt, 2020). A consequent decrease in alveolar ventilation, along with hyper-metabolically increased carbon dioxide production, would account for the observed hypercapnia.

Effects of Butorphanol in White Rhinoceros Immobilized With Etorphine

Plasma Catecholamine Concentrations

Butorphanol administration had no effect on mean plasma noradrenaline concentration; at t40 and t50, it still exceeded concentrations in the non-immobilized rhinoceros. Given that the half-life of noradrenaline is on the order of 2.5 min, this suggests that the butorphanol was not responsible for a substantial normalization of sympathetic activity (Silverberg *et al.*, 1978). Butorphanol is a partial agonist of μ opioid receptors and opposes full agonism of μ opioid receptors by pure agonists like etorphine (Garner *et al.*, 1997). Since μ opioid receptors are likely responsible for the plasma noradrenaline concentrations seen under etorphine (above), one might expect that butorphanol would substantially decrease plasma noradrenaline concentrations as it displaces etorphine from the μ opioid receptors. However, butorphanol is also an agonist of κ opioid receptors, and κ opioid receptor agonism is also capable of activating sympathetic pathways and increasing plasma adrenaline and noradrenaline concentrations, as documented after nalbuphine administration to healthy men (Chang *et al.*, 1981; Hoehe and Duka, 1993). Kappa opioid receptor agonism could explain why a substantial decrease in plasma noradrenaline concentration was not observed after butorphanol administration. Nevertheless, the effects of butorphanol on the metabolic and cardiopulmonary complications caused by etorphine were considerable and largely beneficial, which suggests a pathway independent of plasma noradrenaline concentration; however, our data do not allow us to exclude central sympatholytic effects.

Metabolic Effects

Muscle Tremors. Butorphanol reduced tremor score via an unknown mechanism. Electromyography in standing horses has demonstrated that butorphanol relaxes skeletal muscle; however, in isolated equine skeletal muscle *in vitro*, butorphanol had no effect on contraction, suggesting a

central mechanism of muscle relaxation (Wooldridge *et al.*, 2002; Kruluc and Nemeč, 2006). Butorphanol-induced muscle relaxation could explain the observed decrease in $\dot{V}O_2$.

Cardiopulmonary Effects

Hypoxaemia. Butorphanol significantly improved PaO_2 , increasing it from ~ 27 mm Hg at baseline to ~ 47 – 49 mm Hg. However, PaO_2 is ~ 90 – 100 mm Hg in non-immobilized white rhinoceros habituated to arterial puncture; thus, butorphanol did not normalize PaO_2 completely (Citino and Bush, 2007). Despite residual hypoxaemia, the increase in PaO_2 following butorphanol administration raised CaO_2 from ~ 15 to ~ 20 ml dl^{-1} , a value considered normal in horses (Butler *et al.*, 1993). Given that [Hb] did not increase after butorphanol administration, the normalization of CaO_2 while PaO_2 remains subnormal is likely due to the characteristics of the ODC of the white rhinoceros, which has a P_{50} (oxygen partial pressure at which haemoglobin is 50% saturated) of 20.6 mm Hg, less than that of dogs, humans and horses (28.8, 26.6 and 23.8 mm Hg, respectively) (Clerbaux *et al.*, 1993; Reiners *et al.*, 2019; Shepherd *et al.*, 2019). A substitution of glutamic acid for glutamine at the β_2 position of haemoglobin appears to drive the leftward 'shift' in the ODC in white rhinoceroses. This shift causes their haemoglobin to bind substantially more oxygen at PaO_2 values in the 45–55 mm Hg range compared to other mammals (Mazur *et al.*, 1982).

Butorphanol could have improved PaO_2 in a number of ways. Butorphanol decreased $PaCO_2$; this is compatible with reduced alveolar carbon dioxide partial pressure permitting an increase in PAO_2 and, in turn, increasing PaO_2 (Fenn *et al.*, 1946; Riley and Cournand, 1949; Riley *et al.*, 1951). Butorphanol may have decreased the f_H via vagal stimulation; kappa agonists have induced bradycardia in rats (Gulati and Bhargava, 1988; Hassen and Broudy, 1988). This effect would decrease Q_t and pulmonary vascular pressures, and therefore the hydrostatic pressure driving water into the extravascular space in the pulmonary bed (Starling, 1896). The decrease in Q_t would also have increased pulmonary capillary transit time. The decrease in $\dot{V}O_2$ was associated with an increase in $\bar{C}\bar{v}O_2$ to values that approach those considered normal in horses (Butler *et al.*, 1993). In turn, the tendency to normalize $\bar{C}\bar{v}O_2$ would have reduced the influence of all the possible causes of pulmonary venous admixture (including reduced Hb transit time, interstitial oedema and $V:Q < 1$) (Riley *et al.*, 1951).

Despite CaO_2 increasing substantially after butorphanol was administered, $\dot{V}O_2$ did not increase and, in fact, decreased significantly, requiring us to reject our hypothesis that butorphanol would increase $\dot{V}O_2$ in white rhinoceros under etorphine. The immediate mechanism behind this observation is that a substantial decrease in Q_t was observed after butorphanol administration; this was largely driven by a decrease in f_H , as SV did not change significantly. The decrease

in Q_t more than balanced the increase in CaO_2 , causing $\dot{V}O_2$ to decrease.

The OER was not significantly improved by butorphanol. This finding can be explained by the proportionate decrease in both $\dot{V}O_2$ and $\dot{V}O_2$. Inspection of the data suggests that variability in the OER data might conceal a transient reduction that did not achieve statistical significance 3 min after butorphanol was administered (i.e. t40). Butorphanol did not produce a significant reduction in lactate, suggesting that anaerobic metabolism was unchanged by butorphanol. However, the increase in $\bar{C}\bar{v}O_2$ nevertheless suggests an improved oxygen supply-and-demand balance.

Hypoventilation. Three minutes after it was given, butorphanol transiently reduced $PaCO_2$ due to a transient increase in f_R . This effect is consistent with the butorphanol displacing the pure opioid agonist etorphine from μ receptors on neurons in the medulla and pons that are responsible for the rhythm generation of ventilation (Lalley, 2006; Prkic *et al.*, 2012).

Limitations

The major study limitations are the small number of white rhinoceros and the lack of normal ranges for several physiological variables in non-immobilized white rhinoceros due to the challenges posed by invasive monitoring in conscious rhinoceros. However, the use of non-immobilized rhinoceros for establishing normal plasma catecholamine concentrations represents an important step forward in providing reference values for rhinoceros that will benefit future studies.

Summary and Conclusions

In white rhinoceros, the ultra-potent opioid agonist, etorphine, causes sympathetic upregulation characterized by plasma noradrenaline concentrations approximately twice those measured in habituated, non-immobilized white rhinoceros; this sympathetic upregulation contributes to severe arterial hypoxaemia. The hypoxaemia is not only caused by hypoventilation but by increased venous admixture. Sympathetic upregulation, characterized by increased plasma noradrenaline concentration, could contribute to the venous admixture. Increased f_H due to β_1 adrenoceptor agonism increased Q_t and therefore pulmonary pressures. This effect could have produced venous admixture by decreasing erythrocyte transit time through the pulmonary capillaries and/or by increasing P_c sufficiently to cause pulmonary oedema. The high $\dot{V}O_2$ produced mixed venous hypoxaemia, which amplified the effects of pulmonary venous admixture. Despite sympathetic upregulation, etorphine depressed ventilation; this is in accord with respiratory depression ordinarily associated with μ opioid agonist activity in the brainstem.

When butorphanol, a partial μ opioid agonist and pure κ opioid agonist, was administered to white rhinoceros under

etorphine, the ventilatory depression caused by etorphine was attenuated, consistent with partial agonism of, and displacement of etorphine from, μ opioid receptors in medullary and pontine centres responsible for control of breathing. Shortly after butorphanol administration, plasma adrenaline concentration was transiently greater than contemporary values for rhinoceros under etorphine alone; otherwise, the effects of butorphanol on plasma adrenaline, noradrenaline and dopamine concentrations were not significant. Plasma noradrenaline concentration remained much higher than expected, while Q_t , pulmonary pressures and VO_2 decreased from high values under etorphine alone (although they did not reach values considered normal). Butorphanol decreased Q_t and pulmonary pressures by decreasing f_H . The increase in PaO_2 and CaO_2 can be explained not only by the attenuation of ventilatory depression but by the decrease in Q_t . However, the decrease in Q_t offset the increased CaO_2 , this so that DO_2 was actually reduced after butorphanol. Butorphanol reduced VO_2 , apparently by mitigating tremors. The effect of the decrease in VO_2 was that $P\bar{v}O_2$ and $C\bar{v}O_2$ increased, thereby reducing the impact of venous admixture on CaO_2 . The partial normalization of the cardiopulmonary and metabolic values by butorphanol cannot be explained by a decrease in plasma catecholamine concentrations. Maintenance of increased plasma noradrenaline concentration could be a result of κ opioid agonism by butorphanol. Alternatively, butorphanol could decrease DO_2 and VO_2 via mechanisms independent of plasma noradrenaline concentration. Plasma catecholamine concentrations do not necessarily reflect neurotransmitter activity within the sympathetic pathways of the nervous system; therefore butorphanol could still have caused sympatholytic effects.

Future research should aim to mitigate the sympathetic upregulation induced by etorphine. Although butorphanol did not improve DO_2 , pending a practical alternative, its continued use in etorphine-immobilized white rhinoceros is recommended because it counteracts etorphine-induced hypermetabolism, thus reducing VO_2 towards normal and improving the oxygen supply (DO_2) and demand (VO_2) balance.

Acknowledgements

The authors thank SANParks' VWS staff, especially Ms. Leana Rossouw, Mr. Guy Hausler, Ms. Tebogo Manamela and the Capture Team; Ms. Thembeke Mtetwa; Ms. Penelope Miya; Ms. Petronel Nieuwoudt; Dr. Marion Leiberich; Mr. Derek Apps and Dr. Ned Snelling.

Author Contributions

Jordyn M. Boesch (Conceptualization, Methodology, Formal Analysis, investigation, Writing—Original draft preparation, Visualization, Project administration, Funding acquisition); Robin D. Gleed (Conceptualization, Methodology, Investigation, Writing—review and editing, Funding acquisition);

Peter E. Buss (Conceptualization, Methodology, Investigation, Resources, Writing—review and editing, Project administration); Adrian S. W. Tordiffe (Conceptualization, Methodology, Investigation, Resources, Writing—review and editing); Gareth E. Zeiler (Conceptualization, Methodology, Investigation, Writing—review and editing); Michele A. Miller (Conceptualization, Methodology, Investigation, Writing—review and editing); Francois Viljoen (Methodology, Investigation, Resources, Writing—review and editing); Brian H. Harvey (Conceptualization, Methodology, Investigation, Resources, Writing—review and editing); Stephen A. Parry (Formal Analysis, Writing—review and editing); and Leith C. R. Meyer (Conceptualization, Methodology, Formal Analysis, Investigation, Resources, Writing—review and editing, Supervision, Project administration, Funding acquisition)

Conflicts of Interest

Brian H. Harvey and Leith C. R. Meyer have received funding for other projects from Wildlife Pharmaceuticals South Africa. The other authors have no competing interests to declare.

Funding

This work was supported by the John T. and Jane A. Wiederhold Foundation, Cornell University; the Department of Paraclinical Sciences, University of Pretoria (UP); South African National Parks (SANParks); the South African Veterinary Association Wildlife Group; the South African National Research Foundation; and Wildlife Pharmaceuticals South Africa. These funders played no role in study design or data analysis and reporting.

Data Availability

All relevant data and resources can be found within the article and its supplementary information.

Supplementary material

Supplementary material is available at *Conservation Physiology* online.

References

- Ally A (1998) Ventrolateral medullary control of cardiovascular activity during muscle contraction. *Neurosci Biobehav Rev* 23: 65–86. [https://doi.org/10.1016/S0149-7634\(97\)00069-9](https://doi.org/10.1016/S0149-7634(97)00069-9).
- Bachelard H, Pitre M, Lessard A (1997) Mechanisms of the regional hemodynamic effects of a μ -opioid receptor agonist microinjected into the hypothalamic paraventricular nuclei of conscious unrestrained rats. *J Pharmacol Exp Ther* 280: 460–470. [https://doi.org/10.1016/S0022-3565\(24\)36372-4](https://doi.org/10.1016/S0022-3565(24)36372-4).
- Bayly WM, Grant BD, Breeze RG, Kramer JW (1983) The effects of maximal exercise on acid-base balance and arterial blood gas tension in

- thoroughbred horses. *Equine exercise physiology. Proceedings of the first international conference*: 400.
- Boesch JM (2020) *Respiratory, Cardiovascular and Metabolic Effects of Etorphine, With and Without Butorphanol, in White Rhinoceros (Ceratotherium simum)*. Faculty of Veterinary Science, University of Pretoria, Onderstepoort, South Africa.
- Boesch JM, Gleed RD, Buss P, Hofmeyr M, Tordiffe A, Zeiler G, Meyer L (2018) Effects of a supplemental etorphine dose on pulmonary artery pressure and cardiac output in immobilized, boma-habituated white rhinoceros (*Ceratotherium simum*): A preliminary study. *Journal of Zoo and Wildlife Medicine* 49: 849–855. <https://doi.org/10.1638/2017-0120.1>.
- Bogan JA, MacKenzie G, Snow DH (1978) Evaluation of tranquilizers for use with etorphine as neuroleptanalgesic agents in horse. *Veterinary Record* 103: 471–472. <https://doi.org/10.1136/vr.103.21.471>.
- Buss P, Miller M, Fuller A, Haw A, Stout E, Olea-Popelka F, Meyer L (2018) Postinduction butorphanol administration alters oxygen consumption to improve blood gases in etorphine-immobilized white rhinoceros. *Veterinary Anaesthesia and Analgesia* 45: 57–67. <https://doi.org/10.1016/j.vaa.2017.03.008>.
- Buss P, Miller M, Fuller A, Haw A, Wanty R, Olea-Popelka F, Meyer L (2016) Cardiovascular effects of etorphine, azaperone, and butorphanol combinations in chemically immobilized captive white rhinoceros (*Ceratotherium simum*). *Journal of Zoo and Wildlife Medicine* 47: 834–843. <https://doi.org/10.1638/2015-0298.1>.
- Buss P, Olea-Popelka F, Meyer L, Hofmeyr J, Mathebula N, Kruger M, Bruns A, Martin L, Miller M (2015) Evaluation of cardiorespiratory, blood gas, and lactate values during extended immobilization of white rhinoceros (*Ceratotherium simum*). *Journal of Zoo and Wildlife Medicine* 46: 224–233. <https://doi.org/10.1638/2014-0089R.1>.
- Butler PJ, Woakes AJ, Smale K, Roberts CA, Hillidge CJ, Snow DH, Marlin DJ (1993) Respiratory and cardiovascular adjustments during exercise of increasing intensity and during recovery in thoroughbred racehorses. *Journal of Experimental Biology* 179: 159–180. <https://doi.org/10.1242/jeb.179.1.159>.
- Cahill AL, Eertmoed AL, Mangoura D, Perlman RL (1996) Differential regulation of phenylethanolamine n-methyltransferase expression in two distinct subpopulations of bovine chromaffin cells. *J Neurochem* 67: 1217–1224. <https://doi.org/10.1046/j.1471-4159.1996.67031217.x>.
- Cambier C, Wierinckx M, Grulke S, Clerbaux T, Serteyn D, Detry B, Liardet MP, Frans A, Gustin P (2008) The effect of colic on oxygen extraction in horses. *Vet J* 175: 102–107. <https://doi.org/10.1016/j.tvjl.2006.12.010>.
- Chang KJ, Hazum E, Cuatrecasas P (1981) Novel opiate binding sites selective for benzomorphan drugs. *Proc Natl Acad Sci U S A* 78: 4141–4145. <https://doi.org/10.1073/pnas.78.7.4141>.
- Chen L, Sica AL, Scharf SM (1999) Mechanisms of acute cardiovascular response to periodic apneas in sedated pigs. *J Appl Physiol* 86: 1236–1246. <https://doi.org/10.1152/jappl.1999.86.4.1236>.
- Citino SB, Bush M (2007) Reference cardiopulmonary physiologic parameters for standing, unrestrained white rhinoceroses (*Ceratotherium simum*). *Journal of Zoo and Wildlife Medicine* 38: 375–379. <https://doi.org/10.1638/2006-0007R1.1>.
- Clerbaux T, Gustin P, Detry B, Cao ML, Frans A (1993) Comparative study of the oxyhaemoglobin dissociation curve of four mammals: Man, dog, horse and cattle. *Comp Biochem Physiol Comp Physiol* 106: 687–694. [https://doi.org/10.1016/0300-9629\(93\)90382-E](https://doi.org/10.1016/0300-9629(93)90382-E).
- Commiskey S, Fan LW, Ho IK, Rockhold RW (2005) Butorphanol: Effects of a prototypical agonist-antagonist analgesic on kappa-opioid receptors. *J Pharmacol Sci* 98: 109–116. <https://doi.org/10.1254/jphs.CRJ05001X>.
- Coupland RE (1965) electron microscopic observations on the structure of the rat adrenal medulla. I. The ultrastructure and organization of chromaffin cells in the normal adrenal medulla. *J Anat* 99: 231–254.
- Daniel M, Ling CM (1972) Effect of an etorphine/acepromazine mixture on heart-rate and blood-pressure of horse. *Veterinary Record* 90: 336–339. <https://doi.org/10.1136/vr.90.12.336>.
- Dodd SL, Powers SK, Brooks E, Crawford MP (1993) Effects of reduced o2 delivery with anemia, hypoxia, or ischemia on peak vo2 and force in skeletal muscle. *J Appl Physiol* 74: 186–191. <https://doi.org/10.1152/jappl.1993.74.1.186>.
- Dumont M, Lemaire S (1984) Opioid receptors in bovine adrenal medulla. *Can J Physiol Pharmacol* 62: 1284–1291. <https://doi.org/10.1139/y84-215>.
- Elman I, Borsook D (2019) The failing cascade: Comorbid post traumatic stress- and opioid use disorders. *Neurosci Biobehav Rev* 103: 374–383. <https://doi.org/10.1016/j.neubiorev.2019.04.023>.
- Emslie R (2020) *Ceratotherium simum, White Rhino, The IUCN Red List of Threatened Species 2020: e.T4185A45813880*. International Union for the Conservation of Nature and Natural Resources
- Fenn WO, Rahn H, Otis AB (1946) A theoretical study of the composition of the alveolar air at altitude. *Am J Physiol* 146: 637–653. <https://doi.org/10.1152/ajplegacy.1946.146.5.637>.
- Feuerstein G, Siren AL (1988) Hypothalamic mu-opioid receptors in cardiovascular control: A review. *Peptides* 9 Suppl 1: 75-78, [https://doi.org/10.1016/0196-9781\(88\)90227-6](https://doi.org/10.1016/0196-9781(88)90227-6)
- Feuerstein G, Zerbe RL, Faden AI (1983) Opiate receptors and cardiovascular control in conscious shr and wky rats. *Hypertension* 5: 663–671. <https://doi.org/10.1161/01.HYP.5.5.663>.
- Gao L, Zhang Z, Feng Z, Wei W, Wu W, Zhi H, Deng Y, Long H, Lei M, Hou J et al. (2021) Fast determination of 16 circulating neurotransmitters and their metabolites in plasma samples of spontaneously hypertensive rats intervened with five different uncaria. *J Chromatogr B Analyt Technol Biomed Life Sci* 1179: 122856. <https://doi.org/10.1016/j.jchromb.2021.122856>.
- Garner HR, Burke TF, Lawhorn CD, Stoner JM, Wessinger WD (1997) Butorphanol-mediated antinociception in mice: Partial agonist effects and mu receptor involvement. *J Pharmacol Exp Ther* 282: 1253–1261. [https://doi.org/10.1016/S0022-3565\(24\)36909-5](https://doi.org/10.1016/S0022-3565(24)36909-5).

- Gharagozlou P, Hashemi E, DeLorey TM, Clark JD, Lameh J (2006) Pharmacological profiles of opioid ligands at kappa opioid receptors. *BMC Pharmacol* 6: 3. <https://doi.org/10.1186/1471-2210-6-3>.
- Greenwald MK, Stitzer ML (1998) Butorphanol agonist effects and acute physical dependence in opioid abusers: Comparison with morphine. *Drug Alcohol Depend* 53: 17–30. [https://doi.org/10.1016/S0376-8716\(98\)00104-5](https://doi.org/10.1016/S0376-8716(98)00104-5).
- Gulati A, Bhargava HN (1988) Cardiovascular responses to kappa opioid agonists in intact and adrenal demedullated rats. *Eur J Pharmacol* 156: 247–257. [https://doi.org/10.1016/0014-2999\(88\)90328-7](https://doi.org/10.1016/0014-2999(88)90328-7).
- Haas SA, Lange T, Saugel B, Petzoldt M, Fuhrmann V, Metschke M, Kluge S (2016) Severe hyperlactatemia, lactate clearance and mortality in unselected critically ill patients. *Intensive Care Med* 42: 202–210. <https://doi.org/10.1007/s00134-015-4127-0>.
- Harthoorn AM, Bligh J (1965) *The use of a new oripavine derivative with potent morphine-like activity for the restraint of hoofed wild animals* 6: 290–300. [https://doi.org/10.1016/S0034-5288\(18\)34737-4](https://doi.org/10.1016/S0034-5288(18)34737-4).
- Hassen AH, Broudy EP (1988) Selective autonomic modulation by mu- and kappa-opioid receptors in the hindbrain. *Peptides* 9 Suppl 1: 63–67. [https://doi.org/10.1016/0196-9781\(88\)90225-2](https://doi.org/10.1016/0196-9781(88)90225-2)
- Hassen AH, Feuerstein G, Faden AI (1984) Selective cardiorespiratory effects mediated by mu opioid receptors in the nucleus ambiguus. *Neuropharmacology* 23: 407–415. [https://doi.org/10.1016/0028-3908\(84\)90248-X](https://doi.org/10.1016/0028-3908(84)90248-X).
- Haw A, Hofmeyr M, Fuller A, Buss P, Miller M, Fleming G, Meyer L (2014) Butorphanol with oxygen insufflation corrects etorphine-induced hypoxaemia in chemically immobilized white rhinoceros (*Ceratotherium simum*). *Bmc Veterinary Research* 10: 9. <https://doi.org/10.1186/s12917-014-0253-0>.
- Haw A, Hofmeyr M, Fuller A, Buss P, Miller M, Fleming G, Meyer L (2015) Butorphanol with oxygen insufflation improves cardiorespiratory function in field-immobilised white rhinoceros (*Ceratotherium simum*). *Journal of the South African Veterinary Association* 86: 10. <https://doi.org/10.4102/jsava.v86i1.1276>.
- Hillidge CJ, Lees P (1971) Preliminary investigations of actions of immobilon in horse. *Veterinary Record* 89: 280–281. <https://doi.org/10.1136/vr.89.10.280-a>.
- Hoehle M, Duka T (1993) Opiates increase plasma catecholamines in humans. *Psychoneuroendocrinology* 18: 141–148. [https://doi.org/10.1016/0306-4530\(93\)90065-5](https://doi.org/10.1016/0306-4530(93)90065-5).
- Hopkins SR, Bayly WM, Slocombe RF, Wagner H, Wagner PD (1998) Effect of prolonged heavy exercise on pulmonary gas exchange in horses. *J Appl Physiol* 84: 1723–1730. <https://doi.org/10.1152/jappl.1998.84.5.1723>.
- Hopkins SR, McKenzie DC, Schoene RB, Glenny RW, Robertson HT (1994) Pulmonary gas exchange during exercise in athletes. I. Ventilation-perfusion mismatch and diffusion limitation. *J Appl Physiol* 77: 912–917. <https://doi.org/10.1152/jappl.1994.77.2.912>.
- Jolicoeur LM, Nassiri MR, Shipman C Jr, Choi HK, Flynn GL (1992) Etorphine is an opiate analgesic physicochemically suited to transdermal delivery. *Pharm Res* 09: 963–965. <https://doi.org/10.1023/A:1015877621976>.
- Kania BF (1977) The effect of etorphine on dopamine and noradrenaline concentrations in different central nervous system structures in the rat. *Acta Physiol Pol* 28: 529–540.
- Kiritsy-Roy JA, Appel NM, Bobbitt FG, Van Loon GR (1986) Effects of mu-opioid receptor stimulation in the hypothalamic paraventricular nucleus on basal and stress-induced catecholamine secretion and cardiovascular responses. *J Pharmacol Exp Ther* 239: 814–822.
- Koba S, Hanai E, Kumada N, Kataoka N, Nakamura K, Watanabe T (2018) Sympathoexcitation by hypothalamic paraventricular nucleus neurons projecting to the rostral ventrolateral medulla. *J Physiol* 596: 4581–4595. <https://doi.org/10.1113/JP276223>.
- Kock MD, Lagrange M, Dutoit R (1990) Chemical immobilization of free-ranging black rhinoceros (*diceros-bicornis*) using combinations of etorphine (m99), fentanyl, and xylazine. *Journal of Zoo and Wildlife Medicine* 21: 155.
- Kock MD, Morkel P, Atkinson M, Foggin C (1995) Chemical immobilization of free-ranging white rhinoceros (*Ceratotherium simum simum*) in hwanje and matobo national parks, zimbabwe, using combinations of etorphine (m99), fentanyl, xylazine, and detomidine. *Journal of Zoo and Wildlife Medicine* 26: 207.
- Kopin IJ (1968) Biosynthesis and metabolism of catecholamines. *Anesthesiology* 29: 654–660. <https://doi.org/10.1097/0000542-196807000-00007>.
- Kruluc P, Nemec A (2006) Electroencephalographic and electromyographic changes during the use of detomidine and detomidine-butorphanol combination in standing horses. *Acta Vet Hung* 54: 35–42. <https://doi.org/10.1556/avet.54.2006.1.4>.
- Lalley PM (2006) Opiate slowing of feline respiratory rhythm and effects on putative medullary phase-regulating neurons. *Am J Physiol Regul Integr Comp Physiol* 290: R1387–R1396. <https://doi.org/10.1152/ajpregu.00530.2005>.
- de Lange SS, Fuller A, Haw A, Hofmeyr M, Buss P, Miller M, Meyer LCR (2017) Tremors in white rhinoceroses (*Ceratotherium simum*) during etorphine-azaperone immobilisation. *Journal of the South African Veterinary Association* 88: 10. <https://doi.org/10.4102/jsava.v88i0.1466>.
- Langhout MvZ, Caraguel CGB, Raath JP, Boardman WSJ (2016) Evaluation of etorphine and midazolam anesthesia, and the effect of intravenous butorphanol on cardiopulmonary parameters in game-ranched white rhinoceroses (*ceratotherium simum*). *J Zoo Wildl Med* 47: 827–833.
- Lees P, Hillidge CJ (1975) Neuroleptanalgesia and cardiovascular function in the horse. *Equine Veterinary Journal* 7: 184–191. <https://doi.org/10.1111/j.2042-3306.1975.tb03265.x>.

- Levick JR, Michel CC (2010) Microvascular fluid exchange and the revised Starling principle. *Cardiovasc Res* 87: 198–210. <https://doi.org/10.1093/cvr/cvq062>.
- Levin EY, Levenberg B, Kaufman S (1960) The enzymatic conversion of 3,4-dihydroxyphenylethylamine to norepinephrine. *J Biol Chem* 235: 2080–2086. [https://doi.org/10.1016/S0021-9258\(18\)69366-4](https://doi.org/10.1016/S0021-9258(18)69366-4).
- van Loon GR, Appel NM, Ho D (1981) B-endorphin-induced stimulation of central sympathetic outflow: B-endorphin increases plasma concentrations of epinephrine, norepinephrine, and dopamine in rats*. *Endocrinology* 109: 46–53. <https://doi.org/10.1210/endo-109-1-46>.
- Mansour A, Lewis ME, Khachaturian H, Akil H, Watson SJ (1986) Multiple opioid receptor subtypes in the pituitary-adrenal axis: A cross-species study. *NIDA Res Monogr* 75: 311–314.
- Marsden CD, Meadows JC (1970) The effect of adrenaline on the contraction of human muscle. *J Physiol* 207: 429–448. <https://doi.org/10.1113/jphysiol.1970.sp009071>.
- Mazur G, Braunitzer G, Wright PG (1982) The primary structure of the hemoglobin from a white rhinoceros (*Ceratotherium simum*, perissodactyla) - beta-2 glu. *Hoppe-Seyler's Zeitschrift Fur Physiologische Chemie* 363: 1077–1086. <https://doi.org/10.1515/bchm2.1982.363.2.1077>.
- McCoy AM, Hackett ES, Wagner AE, Mama KR, Hendrickson DA (2011) Pulmonary gas exchange and plasma lactate in horses with gastrointestinal disease undergoing emergency exploratory laparotomy: A comparison with an elective surgery horse population. *Vet Surg* 40: 601–609. <https://doi.org/10.1111/j.1532-950X.2011.00840.x>.
- McCue ME, Bannasch DL, Petersen JL, Gurr J, Bailey E, Binns MM, Distl O, Guerin G, Hasegawa T, Hill EW *et al.* (2012) A high density SNP array for the domestic horse and extant perissodactyla: Utility for association mapping, genetic diversity, and phylogeny studies. *Plos Genetics* 8: 14. <https://doi.org/10.1371/journal.pgen.1002451>.
- Miller JR, Zuperku EJ, Stuth EAE, Banerjee A, Hopp FA, Stucke AG (2017) A subregion of the parabrachial nucleus partially mediates respiratory rate depression from intravenous remifentanyl in young and adult rabbits. *Anesthesiology* 127: 502–514. <https://doi.org/10.1097/ALN.0000000000001719>.
- Miller M, Kruger M, Olea-Popelka F, Buss P (2016) A scoring system to improve decision making and outcomes in the adaptation of recently captured white rhinoceroses (*Ceratotherium simum*) to captivity. *Journal of wildlife diseases* 52: S78–S85. <https://doi.org/10.7589/52.2.S85>.
- Milne DW, Muir WW, Skarda RT (1975) Pulmonary arterial wedge pressures: Blood gas tensions and pH in the resting horse. *Am J Vet Res* 36: 1431–1434.
- Mosing M, Waldmann AD, Sacks M, Buss P, Boesch JM, Zeiler GE, Hosgood G, Gleed RD, Miller M, Meyer LCR *et al.* (2020) What hinders pulmonary gas exchange and changes distribution of ventilation in immobilized white rhinoceroses (*Ceratotherium simum*) in lateral recumbency? *J Appl Physiol* 129: 1140–1149. <https://doi.org/10.1152/jappphysiol.00359.2020>.
- Murray JF (2011) Pulmonary edema: Pathophysiology and diagnosis. *Int J Tuberc Lung Dis* 15: 155–160.
- MvZ L, Caraguel CGB, Raath JP, Boardman WSJ (2016) Evaluation of etorphine and midazolam anesthesia, and the effect of intravenous butorphanol on cardiopulmonary parameters in game-ranched white rhinoceroses (*Ceratotherium simum*). *Journal of Zoo and Wildlife Medicine* 47: 827–833. <https://doi.org/10.1638/2015-0167.1>.
- Naeije R, Chesler N (2012) Pulmonary circulation at exercise. *Compr Physiol* 2: 711–741.
- Nagatsu T, Levitt M, Udenfriend S (1964) Tyrosine hydroxylase. The initial step in norepinephrine biosynthesis. *J Biol Chem* 239: 2910–2917. [https://doi.org/10.1016/S0021-9258\(18\)93832-9](https://doi.org/10.1016/S0021-9258(18)93832-9).
- Nelson DP, King CE, Dodd SL, Schumacker PT, Cain SM (1987) Systemic and intestinal limits of O₂ extraction in the dog. *J Appl Physiol* (1985) 63: 387–394. <https://doi.org/10.1152/jappl.1987.63.1.387>.
- Percie du Sert N, Hurst V, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, Clark A, Cuthill IC, Dirnagl U *et al.* (2020) The ARRIVE guidelines 2.0: Updated guidelines for reporting animal research. *PLoS Biol* 18: e3000410. <https://doi.org/10.1371/journal.pbio.3000410>.
- Prakash N, Banerji HN (1972) Evaluation of cyanmethaemoglobin method for haemoglobin estimation. *Indian J Chest Dis* 14: 102–105.
- Prkic I, Mustapic S, Radocaj T, Stucke AG, Stuth EA, Hopp FA, Dean C, Zuperku EJ (2012) μ -opioid receptors mediate bradypnea caused by intravenous remifentanyl infusions at clinically relevant concentrations in dogs. *J Neurophysiol* 108: 2430–2441. <https://doi.org/10.1152/jn.00185.2012>.
- Quirion R, Zajac JM, Morgat JL, Roques BP (1983) Autoradiographic distribution of mu and delta opiate receptors in rat brain using highly selective ligands. *Life Sci* 33 Suppl 1: 227–230. [https://doi.org/10.1016/0024-3205\(83\)90484-8](https://doi.org/10.1016/0024-3205(83)90484-8).
- Reiners JK, Hellmann N, Schmidt J, Kastner SBR (2019) Odd haemoglobins in odd-toed ungulates: Impact of selected haemoglobin characteristics of the white rhinoceros (*Ceratotherium simum*) on the monitoring of the arterial oxygen saturation of haemoglobin. *Plos One* 14: e0226851. <https://doi.org/10.1371/journal.pone.0226851>.
- Riley RL, Cournand A (1949) Ideal alveolar air and the analysis of ventilation-perfusion relationships in the lungs. *J Appl Physiol* 1: 825–847. <https://doi.org/10.1152/jappl.1949.1.12.825>.
- Riley RL, Cournand A (1951) Analysis of factors affecting partial pressures of oxygen and carbon dioxide in gas and blood of lungs; theory. *J Appl Physiol* 4: 77–101. <https://doi.org/10.1152/jappl.1951.4.2.77>.
- Riley RL, Cournand A, Donald KW (1951) Analysis of factors affecting partial pressures of oxygen and carbon dioxide in gas and blood of lungs; methods. *J Appl Physiol* 4: 102–120. <https://doi.org/10.1152/jappl.1951.4.2.102>.
- Rose CE Jr, Althaus JA, Kaiser DL, Miller ED, Carey RM (1983) Acute hypoxemia and hypercapnia: Increase in plasma catecholamines in conscious dogs. *Am J Physiol* 245: H924–H929. <https://doi.org/10.1152/ajpheart.1983.245.6.H924>.

- Ross CA, Ruggiero DA, Joh TH, Park DH, Reis DJ (1984) Rostral ventrolateral medulla: Selective projections to the thoracic autonomic cell column from the region containing c1 adrenaline neurons. *J Comp Neurol* 228: 168–185. <https://doi.org/10.1002/cne.902280204>.
- Saunders SE, Levitt ES (2020) Kölliker-fuse/parabrachial complex mu opioid receptors contribute to fentanyl-induced apnea and respiratory rate depression. *Respir Physiol Neurobiol* 275: 103388. <https://doi.org/10.1016/j.resp.2020.103388>.
- Schlarmann B, Gorlitz BD, Wintzer HJ, Frey HH (1973) Clinical pharmacology of an etorphine-acepromazine preparation - experiments in dogs and horses. *American J Veterinary Res* 34: 411–415.
- Schmidt-Nielsen K (1991) *Scaling, Why Is Animal Size So Important?* Cambridge University Press, Cambridge, New York
- Shepherd JRA, Dominelli PB, Roy TK, Secomb TW, Hoyer JD, Oliveira JL, Joyner MJ (2019) Modelling the relationships between haemoglobin oxygen affinity and the oxygen cascade in humans. *J Physiol* 597: 4193–4202. <https://doi.org/10.1113/JP277591>.
- Silverberg AB, Shah SD, Haymond MW, Cryer PE (1978) Norepinephrine: Hormone and neurotransmitter in man. *Am J Physiol* 234: E252–E256. <https://doi.org/10.1152/ajpendo.1978.234.3.E252>.
- Sinha AK, Glead RD, Hakim TS, Dobson A, Shannon KJ (1996) Pulmonary capillary pressure during exercise in horses. *J Appl Physiol* 80: 1792–1798. <https://doi.org/10.1152/jap.1996.80.5.1792>.
- Siren AL, Feuerstein G (1991) Hypothalamic opioid mu-receptors regulate discrete hemodynamic functions in the conscious rat. *Neuropharmacology* 30: 143–152. [https://doi.org/10.1016/0028-3908\(91\)90197-J](https://doi.org/10.1016/0028-3908(91)90197-J).
- Siren AL, Paakkari P, Goldstein DS, Feuerstein G (1989) Mechanisms of central hemodynamic and sympathetic regulation by mu opioid receptors: Effects of dermorphin in the conscious rat. *J Pharmacol Exp Ther* 248: 596–604.
- Starling EH (1896) On the absorption of fluids from the connective tissue spaces. *J Physiol* 19: 312–326. <https://doi.org/10.1113/jphysiol.1896.sp000596>.
- Strack AM, Sawyer WB, Hughes JH, Platt KB, Loewy AD (1989) A general pattern of cns innervation of the sympathetic outflow demonstrated by transneuronal pseudorabies viral infections. *Brain Res* 491: 156–162. [https://doi.org/10.1016/0006-8993\(89\)90098-X](https://doi.org/10.1016/0006-8993(89)90098-X).
- Swan HJ, Ganz W, Forrester J, Marcus H, Diamond G, Chonette D (1970) Catheterization of the heart in man with use of a flow-directed balloon-tipped catheter. *N Engl J Med* 283: 447–451. <https://doi.org/10.1056/NEJM197008272830902>.
- Tolkovsky AM (1982) Etorphine binds to multiple opiate receptors of the caudate nucleus with equal affinity but with different kinetics. *Mol Pharmacol* 22: 648–656.
- Vivian JA, DeYoung MB, Sumpter TL, Traynor JR, Lewis JW, Woods JH (1999) Kappa-opioid receptor effects of butorphanol in rhesus monkeys. *J Pharmacol Exp Ther* 290: 259–265. [https://doi.org/10.1016/S0022-3565\(24\)34893-1](https://doi.org/10.1016/S0022-3565(24)34893-1).
- Wan X, Wang W, Liu J, Tong T (2014) Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 14: 135. <https://doi.org/10.1186/1471-2288-14-135>.
- West JB, Mathieu-Costello O, Jones JH, Birks EK, Logemann RB, Pascoe JR, Tyler WS (1993) Stress failure of pulmonary capillaries in racehorses with exercise-induced pulmonary hemorrhage. *Journal Appl Physiol* 75: 1097–1109. <https://doi.org/10.1152/jap.1993.75.3.1097>.
- Whitehead RD Jr, Mei Z, Mapango C, Jeffers MED (2019) Methods and analyzers for hemoglobin measurement in clinical laboratories and field settings. *Ann N Y Acad Sci* 1450: 147–171. <https://doi.org/10.1111/nyas.14124>.
- Wong DM, Hepworth-Warren KL, Sponseller BT, Howard JM, Wang C (2017) Measured and calculated variables of global oxygenation in healthy neonatal foals. *Am J Vet Res* 78: 230–238. <https://doi.org/10.2460/ajvr.78.2.230>.
- Wooldridge AA, Eades SC, Hosgood GL, Moore RM (2002) Effects of treatment with oxytocin, xylazine butorphanol, guaifenesin, acepromazine, and detomidine on esophageal manometric pressure in conscious horses. *Am J Vet Res* 63: 1738–1744. <https://doi.org/10.2460/ajvr.2002.63.1738>.
- Wurtman RJ (2002) Stress and the adrenocortical control of epinephrine synthesis. *Metabolism* 51: 11–14. <https://doi.org/10.1053/meta.2002.33185>.
- Xu H, Wang Z, Zhu L, Sui Z, Bi W, Liu R, Bi K, Li Q (2018) Targeted neurotransmitters profiling identifies metabolic signatures in rat brain by lc-ms/ms: Application in insomnia, depression and alzheimer's disease. *Molecules* 23. <https://doi.org/10.3390/molecules23092375>.