Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Cognitive Change and Assessment during Electroconvulsive Therapy
A thesis presented in partial fulfillment of the requirements for the Degree of Doctor of
Clinical Psychology at Massey University, Wellington, New Zealand.
Anneke Annette Thornton 2014

Abstract

Electroconvulsive therapy (ECT) is the most effective treatment available for depression; however, cognitive side effects limit its use. If detected early in treatment, cognitive decline can be reduced by modifying the way the treatment is administered. Currently, no gold standard measures exist to assess ECT related cognitive change. The studies within this thesis aimed to improve the standard in which cognition is assessed during a course of ECT.

Study One surveyed health professionals across New Zealand on their current practice of cognitive assessment during ECT. Study Two collected normative data for a neuropsychological measure of motor speed which had potential to be included in a cognitive screening measure for use with people receiving ECT. Study Three prospectively investigated objective cognitive changes in 13 people receiving electroconvulsive therapy for a mood disorder. Study Four investigated the subjective cognitive changes by qualitatively analysing patients' reports of cognitive function throughout treatment. Finally, a brief but comprehensive cognitive screening measure was proposed for repeated use with patients receiving ECT. This measure was largely informed by findings from the aforementioned studies.

Time, resources and a lack of sensitive tests restricted professionals from conducting more frequent and thorough cognitive assessments. ECT resulted in both cognitive decline and improvement across a range of cognitive domains. Alleviation in depression was associated with improvement in cognitive function from baseline. Six weeks post ECT, cognitive decline most often resolved back to or was superior to the baseline functioning. Domains sensitive to decline during a course of treatment included retrograde memory, anterograde memory, verbal and visual learning, attention, verbal fluency and information processing speed. Retrograde amnesia was more likely for memories formed closer in time to ECT treatment. Subjective reports of cognitive change were broad, and varied as a function of treatment phase and severity of depression. It is proposed that future research determine the psychometric properties of the suggested screening measure.

Preface

As I was entering the Doctor of Clinical Psychology programme at Massey University, a predecessor in the programme, Dr Kiri Luther, also working under the supervision of Professor Janet Leathem, had just completed her doctoral research in the field of cognitive change associated with electroconvulsive therapy (ECT). Kiri conducted cognitive assessments with 19 people who had received ECT at least two years previously and found that some areas of cognitive deficit due to ECT endured into the long term. This finding highlighted the importance of minimising cognitive dysfunction early on in treatment by frequently monitoring and assessing cognitive function and thereby preventing long term cognitive deficits. Kiri and Janet's passion for the topic was contagious, and I was soon convinced that further research in this area was warranted.

A discussion with the ECT administering psychiatrist at Capital and Coast District Health Board (C&C DHB) Dr Nisar Contractor revealed longstanding frustration with the paucity of assessment measures available to sensitively detect cognitive change during ECT. Current measures were reportedly insensitive to detecting cognitive change reported by patients, and lengthier neuropsychological assessments, though more sensitive, were impractical due to time constraints. It was decided that further research into the neuropsychological assessment of patients receiving ECT, and the subsequent development of a brief screening measure tailored to detect cognitive change during ECT would be beneficial to both patients receiving ECT and health professionals responsible for their care.

At the outset of planning the research, it was hoped that a large enough number of participants would be recruited in order to analyse the results collectively using parametric statistics. After conversations with Professors Richard Porter and Bob Knight, two New Zealand researchers who have both contributed extensively to research in this field, the issue of recruitment was raised. It soon became apparent the greatest challenge in conducting this type of research would be in recruiting participants. This concern was also expressed by Susan Vella, ECT nurse at C&C DHB.

Recruitment challenges would be due to a number of factors: the small number of people who are prescribed ECT, the large physical distances between District Health Boards (DHBs) and thus the travel involved to recruit participants from across DHBs, and due to the characteristics of the population under investigation. For example, as most people who receive ECT suffer from severe depression, many would not be well enough to undergo cognitive assessments, to consent to taking part in research or be motivated to partake in research. In addition, there was a strict timeframe for completion of the research; the entirety of the research needed to be designed, receive ethics approval, be conducted, analysed, written up and submitted within 24 months.

Although early on in planning the research it was understood and acknowledged that recruiting participants would be difficult, it was agreed that a meaningful contribution to the existing pool of knowledge on cognitive effects of ECT could still be made even with a small participant sample. The implication, however, was that the way in which the results would be analysed would need to differ. Presenting the results as a series of case studies would abolish the need for a large participant pool required for hypothesis testing, and instead would represent what cognitive changes occurred for each individual as a function of how many treatments they had had, and how their mood changed over their course of treatment. The method of analysis was appropriate as each individual was variable in terms of ECT administration, seizure duration, threshold, number of treatments, psychiatric illness, co-morbidities, socio economic status, educational history, age, sex, and ethnicity to name a few. As analysing the results as a series of case studies would still provide rich and meaningful data, it was decided to go ahead with conducting the research. Despite receiving

support and referrals from four DHBs in the Lower North Island¹, the number of participants recruited was, as anticipated, low. However, meticulous efforts were made to ensure that baseline assessments, follow-up assessments and reassessments after every three treatments of ECT were made, which resulted in a total of 87 cognitive assessments conducted.

This thesis presents and discusses current research into the short term cognitive effects of electroconvulsive therapy and the assessment thereof. The research was conducted in order to better understand the nature and severity of the cognitive effects endured by 13 patients receiving ECT and to determine precisely, what cognitive changes occurred during their courses of ECT. In addition to adding to the existing literature on the short term cognitive effects of ECT, the current study also endeavoured to determine which measures were sensitive and insensitive to detecting the effects of ECT, and which cognitive changes people were reporting that were not being detected by the objective measures of assessment. The findings of this study later informed the suggestion of a screening measure, to assess and monitor cognitive function during a course of ECT.

¹ Capital and Coast DHB, Hutt Valley DHB, Mid Central DHB and Hawke's Bay DHB

Acknowledgements

Conducting such research involves the help, support and collaboration of many individuals, and could never be achieved in isolation. First and foremost, I would like to acknowledge the people undergoing ECT who took part in the research, and the whānau of many of these people who supported their participation. It was a privilege to get to know each and every one of you and to work with you during this particularly difficult time in your lives. It takes great motivation and altruism to take part in voluntary research, particularly when suffering such debilitating illness. I have been touched by your courage and persistence, and I hope you have also benefitted from being a part of this research.

Thank you to the ongoing support and encouragement from Dr Nisar Contractor and Susan Vella psychiatrist and ECT nurse from C&C DHB who played a large part in the initiation of the research, provided ongoing support and encouragement throughout the duration of the research, and always kept me up to date with new referrals and treatment progress of participants.

Thank you to Erwin Sonnendecker, Clinical Psychologist at Hawke's Bay DHB for showing an avid interest in the research, and sharing a passion for improving the way in which cognition is assessed during a course of ECT; a passion which resulted in Erwin contributing some assessment data to the research.

Thank you to my parents for encouraging me in my first year of university to pursue my interests and to take papers which I enjoy, and not worry about whether or not they would someday lead to a career. Advice which has serendipitously led to me writing this doctorate and pursuing a career in clinical psychology. Thank you to my partner, Ben, whose endless support and numerous cups of teas and coffees (and at times dinner brought to my office) has helped me complete this thesis. I would also like to acknowledge my office peers in T4 who have now all become close friends.

Last but by no means least; I would like to extend my sincere gratitude to my wonderful supervisors, Professor Janet Leathem and Dr Ross Flett. Janet's clinical and research expertise has been indispensible throughout the duration of the research and I have been fortunate to have learnt from her. No problem was ever too big or too complicated for her to solve. If she didn't have the answers, she always knew the right person to contact. To my second supervisor Ross, thank you for your guidance, sense of humour and encouragement you generously provided.

Table of Contents

THESIS OVERVIEW	9
CHAPTER ONE	13
The History and Evolution of Electroconvulsive Therapy	13
Modern-day Electroconvulsive Therapy	17
CHAPTER TWO	27
Variables Affecting the Cognitive and Clinical Outcome during Electroconvulsive Therapy	27
Part I: Person Related Variables	28
Part II - Treatment Related Variables	33
CHAPTER THREE	42
Short Term Cognitive Side Effects of Electroconvulsive Therapy	42
Recommendations and Guidelines for the Assessment of Cognitive Change during ECT	51
CHAPTER FOUR	58
Cognitive Assessment during a Course of Electroconvulsive Therapy - A National Questionnaire	e
Survey of Current Practice in Aotearoa, New Zealand	58
Abstract	59
Method	64
Results	65
Discussion	68
References	70
CHAPTER FIVE	76

Selecting and Refining the Assessment Measures for the Investigation into the Short Term	
Cognitive Effects of Electroconvulsive Therapy	76
Refining the Selected Measures and Procedure for use with Patients receiving Electroconvulsive	e
Therapy	92
CHAPTER SIX	98
Coin Rotation Task – The Development of Norms for New Zealand and the United States	98
Abstract	99
Method	104
Results	107
Discussion	111
References	114
CHAPTER SEVEN	117
Short Term Cognitive Change during Electroconvulsive Therapy for Unipolar and Bipolar	
Depression: A Prospective Study	117
Method	123
Results	129
Group Analysis	129
Single Case Analysis	138
Case Study One	139
Case Study Two	141
Case Study Three	145
Case Study Four	147

Case Study Five	149
Case Study Six	151
Case Study Seven	153
Case Study Eight	155
Case Study Nine	159
Case Study Ten	161
Case Study Eleven	163
Case Study Twelve	166
Case Study Thirteen	168
Discussion	169
References	178
CHAPTER EIGHT	183
Patients' Perspectives on the Cognitive Effects of Electroconvulsive Therapy	183
Abstract	184
Method	191
Results	194
Discussion	211
References	216
CHAPTER NINE	224
The Development of the CASDECT – a Cognitive Screening Measure for use during	
Electroconvulsive Therapy	224
Abstract	225

Results and Discussion	229
References	240
Cognitive Assessment Screen for use during ECT (CASDECT)	247
	259
CHAPTER TEN	260
General Discussion and Reflections on the Research	260
Thesis Reference List	267
List of Figures	
THESIS OVERVIEW – CHPTER THREE	
Figure 1. Flow diagram of thesis structure	9
Figure 2. Age distribution of individuals who received ECT in 2011	25
Figure 3. How the depression and ECT related cognitive effects can overlap across time	29
Figure 4. Physical differences between sine, brief-pulse and ultrabrief-pulse waves	35
Figure 5. Various electrode placements used for ECT.	37
CHAPTER FOUR	
Figure 1. Frequency of assessment tools used to assess cognitive functioning in New	
Zealand.	66
Figure 2. Professional groups involved in assessing cognitive functioning during ECT in New	7
Zealand.	67
Figure 3. Reported time delays between ECT and cognitive assessment in New Zealand	68
CHAPTER SIX	
Figure 1. Finger positioning for the coin rotation task.	106
Figure 2. Latin square demonstrating the control of practice effects.	106
CHAPTER SEVEN	

Figure 1. Standardised changes in depression and cognition for Participant One14	40
Figure 2. Standardised changes in depression and cognition for Participant Two	12
Figure 3. Standardised changes in depression and cognition for Participant Three14	45
Figure 4. Standardised changes in depression and cognition for Participant Four	17
Figure 5. Standardised changes in depression and cognition for Participant Fiv149	9
Figure 6. Standardised changes in depression and cognition for Participant Six	51
Figure 7. Standardised changes in depression and cognition for Participant Seven	53
Figures 8 and 9. Standardised changes in depression and cognition for Participant Eight155,15	57
Figure 10. Standardised changes in depression and cognition for Participant Nine	59
Figure 11. Standardised changes in depression and cognition for Participant Ten	1
Figure 12. Standardised changes in depression and cognition for Participant Eleven16	63
Figure 13. Standardised changes in depression and cognition for Participant Twelve16	66
Figure 14. Standardised changes in depression and cognition for Participant Thirteen16	58
CHAPTER EIGHT	
Figure 1. Subjective reports of cognitive function prior to ECT.	97
Figure 2. Subjective reports of cognitive function during a course of ECT	99
Figure 3. Subjective reports of cognitive function six weeks after a course of ECT	01
Figure 4. Subjective scores of depression and cognitive over course of treatment	07

List of Tables

CHAPTERS ONE - FIVE

Table 1. Domains of Functioning Most Commonly Adversely Affected during a Course of ECT44
Table 2. Battery Selected for Assessment of Cognition and Mood for this Current Study91
Table 3. Domains of Cognitive Function Assessed by Each Test
CHAPTER SIX
Table 1. Mean Dominant Hand Performance for Males and Females for the Coin Rotation
Task
Table 2. Mean Non-Dominant Hand Performance for Males and Females for the Coin Rotation
Task
Table 3. Mean Performance for the Coin Rotation task in Dominant and Non-Dominant Hands for
Males and Females Combined
Table 4. Number of Coin Rotations for Non-Dominant and Dominant Hands for the US Nickel and
the NZ 20 Cent Coin
CHAPTER SEVEN
Table 1. Cognitive Domains Assessed by the Selected Measures
Table 2. Summary of Participant Characteristics
Table 3. Effect Sizes of Cognitive Change which occurred form Baseline to Average Performance
during the Course of ECT.
Table 4. Change in items Assessing Autobiographical and Semantic Memory from Baseline to
during the Course of ECT. 132
Table 5. Effect Sizes of Cognitive Change which occurred from During the Course to Follow-
Up
Table 6. Effect Sizes of Cognitive Change which occurred from Baseline to Follow-Up136
Table 7. Correlations between Subjective Memory Scores and Objective Cognitive Outcome at
Baseline, During the Course and at a Six Week Follow-Up

Tables 8-24. Cognitive and Clinical Changes of Participants 1-13 over a course of ECT and at a Six
Week Follow-Up
CHAPTER EIGHT
Table 1. Effect Sizes of Change in SSMQ Items which occurred form Baseline to during the course
of ECT
Table 2. Effect Sizes of change in SSMQ items which occurred during the course of ECT to the Six
Week Follow-Up
Table 3. Effect Sizes of change in SSMQ items which occurred from Baseline to Follow-Up206
Table 4. Change in Subjective Depression from Baseline to during the course of ECT209
Table 5. Scores on the Beck Depression Inventory-II from Baseline to Follow-Up
CHAPTER NINE
Table 1. Objective Cognitive Changes from Baseline to During the Course of
ECT
Table 2. Change in Items Assessing Autobiographical and Semantic Memory from Baseline to
During the Course of ECT. 231
Table 3. Effect sizes of Change in SSMQ items which occurred from Baseline to During the Course
of ECT
Table 4. Subjective Changes in Mood from Baseline to During the Course of ECT as measured by
the Beck Depression Inventory

List of Appendices

Appendix A. Electronic Questionnaire of Current Practice regarding Cognitive Assessment in	New
Zealand	294
Appendix B. Adapted Squire Ssubjective Memory Questionnaire	299
Appendix C. Summary of Patient Characteristics	300
Appendix D. Information Pack for Participants.	301
Appendix E. Case Report – Effects of Bilateral ECT for a 56 year old Man with Schizoaffecti	ve
Disorder	306
Appendix F. Content Analysis of Subjective Reports at Baseline.	319
Appendix G. Content Analysis of Subjective Reports during a course of ECT	323
Appendix H. Content Analysis of Subjective Reports at the Six Week Follow-Up Assessment	337
Appendix I. Ethics Approval.	341
Appendix J. Conference Presentations.	342