

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Recombinant protein immobilisation and display by alginate

A thesis presented in partial fulfilment of the
requirements of the degree of
Master of Science
in
Microbiology
at Massey University, Palmerston North,
New Zealand



MASSEY UNIVERSITY
TE KUNENGA KI PŪREHUROA

UNIVERSITY OF NEW ZEALAND

Andrew Jameson

2016

ABSTRACT

Biopolymers are a diverse group of organic materials with important applications in a number of industries. Their ability to adsorb and encapsulate compounds has been widely utilised in both biotechnologies and pharmaceuticals. In the last decade, biopolymers have been given new and enhanced functionality, including the separation and purification of compounds. This field is of increasing relevance as advances in the bacterial cell culture process have improved productivity in the biomanufacturing industry, with the establishment of several bacterial host cell lines and optimised protein production systems. This increase in upstream productivity is leading to bottlenecks in downstream processing as current technology platforms reach their limits of throughput and scalability. While previous studies have generated functionalised protein biopolymers using polyhydroxyalkanoate (PHA) biopolyester beads, very few studies have examined the commercially significant biopolymer alginate. Alginate is an exopolysaccharide produced by algae and some bacteria, and is widely utilised in food, pharmaceutical, and biomedical industries because of its stabilising, haemostatic, biocompatible properties and its modifiable structure. In this study, a partially functional alginate-binding recombinant protein was produced, which contained an α -amylase domain from *Bacillus licheniformis* (BLA) translationally fused to the alginate-binding domain of *Pseudomonas aeruginosa* AlgX – an alginate acetyltransferase. An Ssp DnaB mini-intein was included between BLA and AlgX to facilitate recovery of BLA, following immobilisation and display on the surface of alginate. However, aberrant activity of the intein caused total cleavage of the recombinant protein between its BLA and AlgX domains before it could be recovered from the protein production system. Additionally, the absence of a key cysteine residue in the alginate-binding domain prevented the formation of a disulfide bond, which is an essential structural element for the folding and functionality of this region. While this study was unable to overcome intein hyperactivity, functional analysis of the BLA domain showed consistent and significant levels of α -amylase activity, leading to a positive outlook for the functionality of a full-length recombinant protein if proper intein activity can be restored and the necessary cysteine included. In this way, alginate could be specifically functionalised with a desired protein, and in turn, alginate beads could be used for the separation and enrichment of target proteins.

ACKNOWLEDGEMENTS

First and foremost, my gratitude goes to my supervisor, Dr. Bernd Rehm. Thank you for the opportunity, the challenges, the exceptional work environment, and for the understanding and patience when I floundered. I wasn't always the best student, sometimes not even a decent one, but you never gave up on me, and for that I'm in your debt.

My deepest appreciation to my co-supervisor Dr. Zoe Jordens for always believing in me. It was your enthusiasm and confidence that convinced me to pursue postgrad, and without you I would not be writing this today at the conclusion of my Masters of Science.

Thank you, Dr. Fata Moradali. Not content with just completing your PhD and multiple publications, you had to help a struggling MSc student find their footing in the lab and be their mentor throughout their project. Your efforts, assistance and good spirits enriched my years at Massey University. My one regret is that my research was unable to generate a paper, in which all your hard work could be properly recognised as a co-author. However, this thesis is a testament to your guidance, and I am forever grateful.

I would like to thank my family. To Dad, you were a bastion of patience and calm on those bad days when nothing seemed to go right, and you were unquestioning in your support and advice. To Mom, your endless supply of love and study aids made our home a warm and welcoming place after a long day (week) at Massey. And for my sisters, I will miss the breakfasts and coffees together, your love and encouragement were a constant. I love you all.

A heartfelt thank you to the Rehm lab and PolyBatics groups. Too many to name, too great to leave out. Each of you made the lab my home-away-from-home. From the reagents being cooked up in the lab, the discussions around the table at lunch, or the comradery from working well into the night, every one of you enlivened my postgraduate experience, and I will sincerely miss you all.

To my friends, thank you for enduring the endless rants about my project. You saw me at my worst, and still gave me your best.

My appreciation goes out to Massey University and the Institute of Fundamental Sciences. You provided a supportive and collegial workplace for my postgrad; and finally fixed the WiFi connection in the reading room, God bless. Importantly, there are people of IFS that I wish to personally thank. To Ann Truter, you provided the greatest assistance to all my bureaucratic fumbblings. I offer my gratitude for your patience when I took an eternity to respond to emails. To Dr. Kathryn Stowell, your care and diligence to the postgraduate students of IFS is inspiring, and I do not know if I would have finished postgrad without your compassion and support. To Trevor Loo, I will always appreciate your guidance and technical expertise with gel filtration chromatography; you made one of the most confusing and frustrating experiences look effortless. And finally to the people of the Massey Genome Service, thank you all for your exceptional work sequencing my twice-damned plasmids and genes.

Speaking of cursed idols, thank you to the University of Otago and the Centre for Protein Research for your tireless efforts to ruin my day with peptidase S8. I wish I could have given you my actual recombinant protein to analyse, but I do sincerely appreciate your work on those damnable proteases.

Finally, to my partner Javier, thank you for your unconditional love and support these last two-and-a-half years. You picked me up and dusted me off after each stumble, and stood by my side and faced down every problem and obstacle. Without you, I wouldn't have made it. You deserve the world.

TABLE OF CONTENTS

ABSTRACT.....	i
ACKNOWLEDGEMENTS.....	ii
LIST OF ABBREVIATIONS.....	viii
LIST OF FIGURES.....	x
LIST OF TABLES.....	xii
CHAPTER ONE INTRODUCTION.....	1
1.1 Chromatographic protein purification and a potential separation alternative.....	3
1.1.1 Liquid chromatography.....	3
1.1.2 One-step production of immobilised recombinant protein.....	4
1.2 Alginate structure, biosynthesis and applications.....	5
1.3 AlgX, an alginate-binding protein essential to alginate production.....	13
1.4 Alpha-amylase as a potential fusion partner.....	17
1.5 Intein, the protein intron.....	19
1.6 <i>Bacillus megaterium</i> protein production system.....	22
1.7 Aim and objectives of this study.....	24
1.8 Hypothesis statement.....	24
CHAPTER TWO MATERIALS AND METHODS.....	25
2.1 Strains, plasmids, oligonucleotides and expression vectors.....	25
2.2 Media.....	27
2.2.1 Luria-Bertani (LB) medium.....	27
2.2.2 Trace metal mix A5 medium.....	27
2.2.3 A5-custom medium.....	28
2.2.4 Terrific Broth (TB) medium.....	29
2.2.5 CR5-top agar medium.....	29
2.2.6 Starch agar medium.....	31
2.3 Antibiotic stock solutions and concentrations.....	31
2.4 Media pH buffers.....	31

2.4.1 Phosphate-buffered saline (PBS) pH 7.4.....	31
2.4.2 Tris-HCl pH 8.4.....	32
2.4.3 Borate buffer pH 9.0.....	32
2.4.4 Glycine sodium hydroxide buffer pH 10.0.....	32
2.5 Cultivation conditions.....	32
2.6 Long term storage of strains	32
2.7 Competent cell preparation and plasmid DNA uptake.....	33
2.7.1 Preparation and transformation of competent <i>E. coli</i>	33
2.7.2 Preparation and transformation of competent <i>Bacillus megaterium</i> protoplasts	34
2.8 DNA isolation, analysis and manipulation	35
2.8.1 Isolation of plasmid DNA	35
2.8.2 Isolation of linear dsDNA.....	36
2.8.3 Determination of DNA concentration.....	36
2.8.4 Agarose gel electrophoresis	36
2.8.5 Polymerase chain reaction	37
2.8.6 DNA hydrolysis with restriction endonucleases	38
2.8.7 DNA ligation.....	38
2.8.8 DNA sequencing	38
2.8.9 Addition of a polyhistidine tag.....	38
2.8.9.1 Primer dimer synthesis of a polyhistidine tag.....	38
2.8.9.2 Insertion of a polyhistidine-encoding gene fragment.....	40
2.8.10 Excision of a mini-intein encoding gene (<i>dnaB</i>).....	41
2.9 Production of recombinant protein	41
2.9.1 Protein production in <i>Bacillus megaterium</i> strains	41
2.9.2 Extracellular protein collection.....	42
2.9.2.1 Ammonium sulfate precipitation.....	42
2.9.2.2 Membrane ultrafiltration.....	43
2.9.3 Determination of protein concentration.....	43
2.10 Detection of recombinant protein	43
2.10.1 Sodium dodecyl sulfate (SDS) gel electrophoresis.....	43
2.10.1.1 Preparation of protein samples for SDS-PAGE	45
2.10.1.2 Protein molecular weight standard for SDS-PAGE.....	45
2.10.2 Immunoblot analysis	45

2.10.2.1	Transfer to a nitrocellulose membrane	45
2.10.2.2	Blocking, antibody application, and visualisation	46
2.10.3	Native gel electrophoresis.....	47
2.10.3.1	Preparation of protein samples for native PAGE	48
2.10.3.2	Protein molecular weight standard for native PAGE	48
2.10.4	MALDI-TOF peptide mass analysis.....	48
2.11	Purification of recombinant protein.....	49
2.11.1	Protein solubility fractionation by ammonium sulfate	49
2.11.2	Size exclusion chromatography.....	49
2.11.3	Affinity purification of polyhistidine-tagged protein	49
2.11.4	Affinity purification of alginate-binding protein	50
2.12	Functional analysis of the recombinant protein.....	50
2.12.1	Starch degradation screen for α -amylase activity	50
2.12.2	Maltose assay for α -amylase activity.....	50
CHAPTER THREE	RESULTS	52
3.1	Molecular Cloning	52
3.1.2	Construction of plasmid p1623amySdnaBBDx	52
3.1.3	Transformation of <i>Bacillus megaterium</i>	53
3.2	Production of AmyS:DnaB:ABDx.....	54
3.3	Functional analysis of AmyS:DnaB:ABDx	57
3.3.1	AmyS functional analysis	57
3.3.2	ABDx functional analysis.....	58
3.4	Identification of AmyS:DnaB:ABDx	60
3.5	Detection and isolation of AmyS:DnaB:ABDx	60
3.5.1	Anti-AlgX immunoblot	60
3.5.2	Native protein detection and isolation.....	62
3.5.2.1	Native PAGE analysis	62
3.5.2.2	SDS-PAGE protein renaturation.....	63
3.5.3	Fractionation	64
3.5.3.1	Protein fractionation by solubility	65
3.5.3.2	Size exclusion chromatography	66
3.6	Construction of amySdnaBBDxhis	68

3.6.1 Primer dimer His-tagging.....	69
3.6.2 BsrGI-KpnI His-tag insertion.....	69
3.7 Transformation and production of AmyS:DnaB:ABDx:His	70
3.7.1 Analysis of AmyS:DnaB:ABDx:His.....	71
3.7.2 Ssp DnaB intein activity inhibition	72
3.8 Functional analysis of AmyS:DnaB:ABDx:His	74
3.8.1 Functional analysis of AmyS	75
3.8.2 Functional analysis of ABDx.....	76
3.9 Construction of <i>amySABDxhis</i> and <i>dnaB</i> excision	77
3.10 Transformation and production of AmyS:ABDx:His	79
CHAPTER FOUR DISCUSSION	82
4.1 Production of AmyS:DnaB:ABDx.....	82
4.2 Detection and isolation of AmyS:DnaB:ABDx	84
4.2.1 Anti-AlgX immunoblot	84
4.2.2 Native PAGE analysis and protein renaturation	86
4.2.3 Fractionation.....	87
4.3 Generation of AmyS:DnaB:ABDx with a detectable marker and discovery of aberrant intein activity.....	91
4.4 Intein removal.....	96
4.5 Partial functionality of AmyS:DnaB:ABDx:His	98
4.6 Conclusion.....	99
4.7 Future directions.....	100
APPENDIX I EXPRESSION VECTOR MAPS	104
APPENDIX II AMMONIUM SULFATE SATURATION TABLE.....	109
APPENDIX III MALDI-TOF PEPTIDE MASS ANALYSIS.....	110
APPENDIX IV SIZE EXCLUSION CHROMATOGRAPHY.....	112
References	114

LIST OF ABBREVIATIONS

A full list of abbreviations.

°C	Degree Celsius
A	Absorbance
AGE	Agarose gel electrophoresis
Ap	Ampicillin
APS	Ammonium persulfate
BLA	<i>Bacillus licheniformis</i> α -amylase
bp	Base pairs
BSA	Bovine serum albumin
Δ	Delta (deleted)
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
dNTPs	Deoxyribonucleotide triphosphates
dsDNA	Double-stranded deoxyribonucleic acid
DTT	Dithiothreitol
ETOH	Ethanol
EDTA	Ethylenediaminetetraacetic acid
g	Gram/gravity
GDP	Guanine diphosphate
His-tag	Polyhistidine-tag
HRP	Horse radish peroxidase
kbp	Kilo base pairs
kDa	Kilodaltons
λ	Lambda phage
LB	Luria-Bertani (broth)
MOPS	3-(N-morpholino)propanesulfonic acid
MW	Molecular weight
OD	Optical density

ORF	Open reading frame
PBS	Phosphate-buffered saline
PCR	Polymerase chain reaction
PEG	Polyethylene glycol
REase	Restriction endonuclease
RNAase	Ribonuclease
Rpm	Rotations/revolutions per minute
SDS	Sodium dodecyl sulfate
SDS-PAGE	Sodium dodecyl sulfate polyacrylamide gel electrophoresis
TB	Terrific Broth
TBE	Tris-borate-EDTA buffer
TBST	Tris-buffered saline and Tween 20
Tet	Tetracycline
TEMED	Tetramethylethylenediamine
T_m	Primer melting temperature
Tris	Trishydroxymethylaminomethane
vol	Volume
v/v	Volume per volume
w/v	Weight per volume

LIST OF FIGURES

Figure 1	Chemical structure of alginate.....	6
Figure 2	Bacterial alginate biosynthesis model.....	8
Figure 3	Applications of alginate.....	10
Figure 4	Structure of AlgX.....	15
Figure 5	Structure of α -amylase.....	18
Figure 6	Protein trans-splicing by intein.....	20
Figure 7	Hybrid gene design.....	27
Figure 8	Primer dimer His-tagging.....	39
Figure 9	His-tagged gene fragment.....	40
Figure 10	Excision of intein-encoding <i>dnaB</i>	41
Figure 11	Confirmation of <i>E. coli</i> transformed with <i>p1623amySdnaBABDx</i>	53
Figure 12	Confirmation of <i>B. megaterium</i> transformed with <i>1623amySdnaBABDx</i>	54
Figure 13	Protein analysis of <i>B. megaterium</i> transformed with <i>p1623amySdnaBABDx</i>	56
Figure 14	Confirmation of α -amylase activity by strains transformed with <i>p1623amySdnaBABDx</i>	58
Figure 15	Analysis of alginate protein purification from strains transformed with <i>p1623amySdnaBABDx</i>	59
Figure 16	Detection of AlgX alginate-binding domain.....	61
Figure 17	Detection of α -amylase activity by protein isolated by native PAGE.....	63
Figure 18	Renaturation of protein isolated by SDS-PAGE.....	64
Figure 19	Detection of α -amylase from fractionated protein.....	66
Figure 20	Protein fractionation by size exclusion chromatography.....	68
Figure 21	Confirmation of <i>E. coli</i> transformed with <i>p1623amySdnaBABDxhis</i>	70
Figure 22	Confirmation of recombinant protein from <i>B. megaterium</i> transformed with <i>p1623amySdnaBABDxhis</i>	72
Figure 23	Intein cleavage is DTT-independent.....	74

Figure 24	Confirmation of α -amylase activity by <i>B. megaterium</i> transformed with p1623amySdnaBBDxhis.....	75
Figure 25	Amplification of <i>amyS</i> and <i>ABDxhis</i> gene fragments.....	77
Figure 26	Confirmation of <i>dnaB</i> excision.....	78
Figure 27	Confirmation of <i>B. megaterium</i> transformed with p1623amySABDxhis.....	80
Figure 28	Protein analysis of <i>B. megaterium</i> transformed with p1623amySABDxhis.....	81
Figure 29	Structure of the C-terminal alginate-binding domain of AlgX.....	93
Figure 30	Sequence maps of expression vectors used in this study.....	104
Figure 31	Protein fractionation by gel filtration chromatography.....	112

LIST OF TABLES

Table 1	Bacterial strains used in this study.....	25
Table 2	Plasmids used in this study.....	26
Table 3	Oligonucleotide primers used in this study.....	26
Table 4	Antibiotic stock solutions and concentrations.....	31
Table 5	Maltose assay for α -amylase activity.....	51
Table 6	Results of a maltose assay for α -amylase activity.....	76
Table 7	Ammonium sulfate saturation table.....	109