Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author. The Investigation of Parenteral Nutrition-Aotearoa (IPNA) – setting up the 1st phase of a clinical audit of the delivery of parenteral nutrition (PN) in New Zealand (NZ)

Sue Larsen 2012

The Investigation of Parenteral Nutrition-Aotearoa (IPNA) – setting up the 1st phase of a clinical audit of the delivery of parenteral nutrition (PN) in New Zealand (NZ)

A thesis presented in partial fulfilment of the requirements for the degree of

**Master of Philosophy** 

At Massey University, Albany, New Zealand

Sue Larsen 2012

#### Abstract

Nutrition support administered as Parenteral Nutrition (PN) is given to patients that have a non-functioning gut. Parenteral nutrition is the administration of nutrients and fluids into the venous system and is potentially associated with life-threatening complications. It is therefore essential that the care and management of PN is co-ordinated by clinicians that have the specialist knowledge and expertise to ensure it is given safely and appropriately.

This is a Phase one regional pilot study which aims to examine the current standard of PN care in hospitals in New Zealand using a clinical audit process. A secondary aim is to identify if any remediable factors are found in the care of patients receiving PN which can then be used to improve patient care, focussing on the following themes:

- Indication for PN
- Type of PN
- Prescribing PN
- Catheter choice, insertion and care
- PN associated complications
- Nutrition teams

Six local hospitals from four large district health boards covering a population of 1.64 million were enrolled. Included were adult, paediatric (<16yrs), and neonates (<1yr) patients receiving PN in hospital during the period of Jan 1st to June 31st 2011. Patients receiving PN in the home were excluded, even if they were admitted into hospital within the study period.

620 cases of PN use (288 adult, 68 paediatric, 264 neonates) were identified within the study period. 151 cases (70 adult, 17 paediatric, 64 neonates) were purposely selected for expert peer review. There were, 66 adults (94%), 7 paediatric (41%), 49 neonates (76%) questionnaires returned, of these, de-Identified clinical records were also available for 100% of the adult and 41% of the neonate cases for expert review.

Data for 66 adults (34 male: 32 female) were returned and peer reviewed by advisor assessors however only 65 completed advisor assessor questionnaires were returned.

The results of the adult cases examined showed that only 12.7% of cases were deemed to involve Good Practice- defined as the standard for which advisors would consider being acceptable and in accordance to the recommended guidelines. Sixty five per cent of cases demonstrated that there was room for improvement in the care provided. Nineteen per cent of cases examined were considered to be of a less than satisfactory standard.

A limitation of this study included lack of sufficient paediatric/neonate experts available for peer review.

### Acknowledgements

This thesis would never have been possible without the help and support given to me, I would like to give thanks and acknowledgment to the following;

To my employer, Waitemata District Health Board, thank you for your assistance and financial contribution towards completing this thesis. Also the WDHB Awhina Health Campus, for all of the research and thesis writing guidance given.

To all of the clinicians involved in this audit, thank you so much for taking the time to participate, and for being motivated to spend so much time collecting, reviewing and analysing data.

To Dr Russell Walmsley, Co-Investigator of this audit, thank you for keeping me focused, on time, and for your patience.

To HBF, Thank you for your suggestions on how I could improve things along the way, both in doing the audit and in writing this thesis. Also thank you for your continuous encouragement, you know how much it means to me.

To my supervisors, Dr Stephen Neville, thank you for your guidance and encouragement in completing this journey. Also Lorraine Neave, Clinical Research Nurse Specialist, WDHB, thank you so much for all the time you have given me, your advice, support and encouragement is appreciated so much.

Finally, to Alun, thank you so much for believing I was capable of achieving this, even though I never believed it myself. Dylan & Megs, thank you for letting me get this finished and for putting up with a grumpy mum!

### Contents

Abstract iii
Acknowledgementsv
List of Figuresxiii
List of Tablesxv
List of Abbreviationsxvii
1. Introduction 1 -
Background 2 -
Normal Function of the Gastrointestinal Tract and Consequences of Malnutrition- 2
Options for Feeding the Malnourished Patient
Venous Access Options for Parenteral Nutrition
Complications of PN 6 -
1. Central Venous Catheter (CVC) complications 6 -
2. Infections 7 -

3. Re-feeding syndrome (RFS)	9 -
Significance of this Study	12 -
Study Rationale	13 -
Research Question	14 -
Organisation of the Thesis – Chapter Overview	15 -
Summary	16 -
2. Literature Review	17 -
Introduction	17 -
Literature Search Strategy	17 -
Guidelines	19 -
Review of the Literature	21 -
Indication for PN	21 -
Type of PN prescribed	23 -
Prescribing PN	24 -
Catheter Choice, Insertion and Care	25 -

PN associated Complications2	27 -
Availability and role of Nutrition Support Teams	28 -
The Mixed Bag Report 3	80 -
Significance of the Literature for this Study	82 -
Summary 3	82 -
3. Methodology and methods 3	4 -
Introduction 3	84 -
Research Methodological Framework 3	34 -
Methods3	89 -
Planning 3	39 -
Ethical Issues & Cultural Responsibilities 4	42 -
Data Collection	44 -
Local Reporter 4	44 -
Case Reviews 4	46 -
Reliability, Validity & Bias	46 -

Data Analysis 48
Summary 49
4. Results 50 -
Introduction 50
Analysis of Data 51
Adult Data 52 -
Overall Assessment of PN Care 53
Administration of PN 55
Indication for PN 56
Type of PN prescribed 60
Prescribing of PN 60
PN Associated Complications 62
Catheter Choice, Insertion and Care 64
Availability and role of Nutrition Support Teams
Organisational Data 67 -

Adult Outcome Data 73	-
Summary 77	-
5. Discussion 78	-
Introduction 78	-
The Investigation of Parenteral Nutrition – Aotearoa	-
Indication for PN78	-
Type of PN prescribed 80	-
Prescribing PN 81	-
Catheter Choice, Insertion and Care82	-
PN Associated Complications84	-
Availability and role of Nutrition Support Teams	-
Validation of study tools87	-
Adherence to guidelines 88	-
Limitations and Weaknesses of the study 89	-
Documentation 89	-

Grading 89
Recommendations from this study for phase two – IPNA 90 ·
Suggested further research activity 91 ·
Concluding statement 92
References 93 -
Appendices 105 -
Appendix 1 – Questionnaires 105 ·
Appendix 2 - Consent Form 106 -
Appendix 3 - Pediatric/Neonatal Data 108 -
Appendix 4 - Ethical Approval 122 -

# List of Figures

Figure 3.1: Audit cycle	- 37 -
Figure 4.1 Age distribution of the study population	52 -
Figure 4.2 Ward level under which PN was administered (Patient care questionnaire	55 -
Figure 4.3 Number of days for which the patient received PN (Patient care questionnaire	- 56
Figure 4.4 Day of the week on which PN was administered, and advisor's opinion or appropriateness of PN being commenced	
Figure 4.5 Deficiencies in monitoring (answers may be multiple)	62 -
Figure 4.6 Types of metabolic complications that occurred (answers may be multiple)	63 -
Figure 4.7 Length of time the initial PN catheter remained in place	65 -
Figure 4.8 Types of wards in hospital	69 -
Figure 4.9 Compositions of NSTs	72 -
Figure 5.1 Likert Scale	91 -
Figure 1.2.1 Age distribution of the paediatric/neonatal study population (at time of PN) -	108 -
Figure 1.2.2 Specialty of doctor making the decision to commence PN 1	112 -
Figure 1.2.3 Grade of doctor determining the patients' nutritional requirements 1	.14 -

Figure 1.2.4 Type of PN first administered	116 -
Figure 1.2.5 Number of days for which PN was received	117 -
Figure 1.2.6 Types of metabolic complication (answers may be multiple)	119 -
Figure 1.2.7 CVC complications	121 -

## List of Tables

Table 1.1: NICE scoring system to identify patients at risk of Re-feeding Syndrome11-
Table 3.1 Defining activities of research, audit and service review
Table 4.1 Overall of assessment of PN care- Advisor assessors
Table 4.2 Documented Indications for PN 57 –
Table 4.3 presents cases where there was evidence of both CVC and/or metabolic complications occurring 66 -
Table 4.4 Designation of the person responsible for deciding the PN composition 67 -
Table 4.5 Designation of the person prescribing PN
Table 4.6 NST involvement in the decision to give PN, and overall assessment on PN care (Adult data only)71 –
Table 4.7 Adult patient outcomes74-
Table 4.8 PN indication and outcome 75 -
Table 4.9 PN duration and outcome76 –
Table 1.2.1 Enteral feeding prior to commencing PN
Table 1.2.2 Assessment made prior to commencing PN

Table 1.2.3 Type of ward where PN was administered
Table 1.2.4 Level of ward on which PN was administered
Table 1.2.5 Indication for PN (answers may be multiple)
Table 1.2.6 Designation of the person responsible for making the decision to start PN 111 –
Table 1.2.7 Grade of doctor making the decision to commence PN
Table 1.2.8 Time between decision to start PN and its commencement
Table 1.2.9 Specialty of doctor determining the patient's nutritional requirements113-
Table 1.2.10 Specialty of doctor signing the prescription
Table 1.2.11 Grade of doctor signing the prescription
Table 1.2.12 Patient outcome 117-
Table 1.2.13 Evidence of metabolic complication118-
Table 1.2.14 Evidence of additional types of fluids
Table 1.2.15 Initial mode of PN
Table 1.2.16 Initial type of PN catheter 121 -

### **List of Abbreviations**

ADHB – Auckland District Health Board

AuSPEN - Australasian Society of Parenteral and Enteral Nutrition

BAPEN – British Society of Parenteral and Enteral Nutrition

BMI – Body Mass Index

CMDHB – Counties Manakau District Health Board

**CPG-** Clinical Practice Guidelines

CRP - C-reactive protein

CT – Computed Tomography

CVC - Central venous catheters

ESPEN – European Society of Parenteral and Enteral Nutrition

ESPGHAN - European Society of Paediatric Gastroenterology, Hepatology and Nutrition

GI – Gastro-Intestinal

ICU/HDU – Intensive Care Unit/High Dependency Unit

IPNA – Investigation of Parenteral Nutrition-Aotearoa

NCEPOD - National Confidential Enquiry into Patient Outcome and Deaths

NDHB – Northlands District Health Board

NHS – National Health System

NICE - National Institute of Clinical Excellence

NSN – Nutrition Support Nurse

NST – Nutrition Support Team

NZ – New Zealand

NZD – New Zealand Dollars

NZNO- New Zealand Nursing Organisation

NZSG - New Zealand Society of Gastroenterologist

Osm – Osmolarity

PDSA – Plan-Do-Study-Act

PICC – Peripherally Inserted Central Catheter

PN – Parenteral Nutrition

PPN- Peripheral Parenteral Nutrition

RCS – Royal College of Surgeons

RFS – Re-feeding syndrome

SIGN – Scottish Intercollegiate Guidelines Network

SIRS – Systemic Inflammatory Response Syndrome

UK – United Kingdom

USD – United State Dollars

WDHB – Waitemata District Health Board