

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

APPLICATION OF DNA HYBRIDISATION
TO THE TAXONOMY OF
RHIZOBIUM TRIFOLII AND RELATED SPECIES

A thesis presented in partial fulfilment
for the requirements for the degree of
Masterate of Science in Microbiology at
Massey University.

Allan George Dick
1979

ABSTRACT

Rhizobia were grown in Yeast Mannitol broth in the presence and absence of K_2HPO_4 . Increasing the concentration of yeast extract in the medium resulted in an increase in growth up to a certain concentration of yeast extract. Growth of rhizobia was inhibited by further increases in yeast extract concentration, especially in the presence of K_2HPO_4 . The inhibition is due to glycine present in the yeast extract and is increased by the presence of monovalent cations.

The molecular weight of DNA extracted from rhizobia was determined by centrifugation in alkaline sucrose gradients. It was found that ^{32}P -labelled DNA was more fragile under shear than unlabelled DNA. Unlabelled DNA required 75 seconds and ^{32}P -labelled DNA required 56 seconds sonication to reduce the fragment size to 230,000 Daltons.

Labelled DNA prepared by a phenol-chloroform method was contaminated with polysaccharide and only low homologous hybridisation could be obtained. A hydroxyapatite-urea method of purifying DNA was developed which produced polysaccharide-free DNA. When labelled DNA was prepared by this method homologous hybridisation averaged 72%. Polysaccharide contamination of unlabelled DNA preparations did not effect the % relative reassociation.

Complete reassociation of heterologous DNA required a longer incubation time than did homologous DNA. The homologous reaction was complete after 32 hours (Cot 200) whereas heterologous DNA required an incubation of 40 hours (Cot 250) to achieve maximum reassociation.

Deoxyribonucleic acid homologies were determined among 27 strains of Rhizobium trifolii, 4 strains of R. leguminosarum and 4 of R. phaseoli. Results from

related strains indicate that DNA homologies correlate with serological relationships and ability to form nodules on legume roots can be lost without detectable change in homology with an independent reference strain. All rhizobia which nodulated effectively on T. repens, T. subterraneum, T. ambiguum, and Vicia hirsuta formed one population with an average relatedness of 70% (range 49-94%) and $\Delta T_m(e)$ of 0.2-7.5°C with respect to reference strains capable of nodulating the first two clover species. Two strains from African Trifolium species and one from a Japanese species were less closely related. The average relatedness of strains from Phaseolus vulgaris with clover rhizobia was 46% (range 37-50%) and $\Delta T_m(e)$ 6.5-10.0°C. Taxonomic revisions consistent with these observations are discussed. It is proposed that R. trifolii and R. leguminosarum should be combined and called Rhizobium leguminosarum Frank. Within this species various biotypes should be designated according to their plant specificity R. phaseoli should be retained as separate species and examined in more detail. The results are discussed in relation to proposed genetic basis for plant specificity.

ACKNOWLEDGEMENTS

I am indebted to the Department of Microbiology and Genetics, Massey University, for providing the opportunity and facilities for this investigation. In particular I would like to thank my supervisor, Dr Jarvis, for constant encouragement and enthusiasm.

I would also like to thank:

the entire academic and technical staff of the Department of Microbiology and Genetics for general help and providing friendly surroundings in which to work;

Mr R.M. Greenwood, D.S.I.R., for supplying the cultures and helpful advice;

Dr W.D. Sutton, D.S.I.R., for helpful advice on DNA hybridisation;

Mr J. McLean for being a steady source of information;

Mr R. Tucker for technical assistance and enthusiasm in the face of adversity;

Miss B. Hunter for the typing;

Miss J. Trow for the excellent diagrams;

Ms C.J. Macdonald, Mr J.B. Smart, and Miss Y. Sprey for helping to get it in on time.

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ACKNOWLEDGEMENTS	iv
LIST OF FIGURES	x
LIST OF TABLES	xii

INTRODUCTION

1. Significance of the Genus <u>Rhizobium</u>	1
2. Classification By Cross-Inoculation Groups . . .	2
3. Application of Numerical Taxonomy and Molecular Genetics to the Classification of <u>Rhizobium</u> ...	5
4. Determination of Base Sequence Homology	11
5. Aim of the Investigation	15

MATERIALS AND METHODS

	Page
1. Microbiological Methods	16
1.1 Media	16
1.2 Microorganisms	16
1.3 Cultivation	17
1.4 Growth Curves	17
1.5 Cell Yield Experiments	21
2. DNA Preparation	22
2.1 Preparation of Unlabelled DNA by a Modification of the Method of Brenner <u>et al.</u> (1969) .	22
2.1.1 Lysis of Cells	22
2.1.2 Extraction of DNA	23
2.1.3 Purification of DNA	23
2.1.4 Routine Measurement of Purity of DNA .	24
2.1.5 Concentration of DNA	24
2.1.6 Shearing of DNA	24
2.2 Preparation of Labelled DNA	24
2.2.1 Lysis and Extraction of DNA	24
2.2.2 Preparation of Hydroxyapatite	24
2.2.3 Hydroxyapatite Purification of DNA ..	25
2.2.4 Shearing of DNA	25
2.2.5 Stripping of DNA	26
3. Methods of Assessing the Suitability of DNA for Hybridisation Experiments	26
3.1 Determination of Melting Temperature of Double-Stranded DNA	26
3.2 Molecular Weight Determination	28
3.3 Colourimetric Determination of DNA Concentration	29
3.4 Colourimetric Determination of Protein Concentration	30
3.4.1 The Method of Bradford (1976)	30
3.4.2 The Method of Lowry <u>et al.</u> (1951) ..	30
3.5 Colourimetric Determination of Polysaccharide Concentration	33
3.5.1 The Method of Trevelyn and Harrison (1952)	32
3.5.2 The Method of Bailey (1967)	32

3.6	Colourimetric Determination of Ribonucleic Acid Concentration	33
4.	Basic Sequence Analysis	33
4.1	Reassociation of DNA	33
4.2	Separation of Single and Double-stranded DNA on Hydroxyapatite	37
4.3	Thermal Stability of Reassociated Duplexes	38
4.4	Statistical Analysis of Reassociation Data	38
4.5	Regeneration of Hydroxyapatite	39

RESULTS

1.	Growth of Rhizobium	40
1.1	Growth Curves	40
1.2	Improvements in Cell Yield	40
2.	Molecular Weight Determination of DNA	45
2.1	Molecular Weight of Unsonicated DNA	45
2.2	Molecular Weight of Sonicated DNA	45
3.	Difficulties Encountered in Hybridisation	45
3.1	First Preparation of labelled DNA	45
3.1.1	Attempted Hybridisation	45
3.1.2	Absorbance Spectrum	48
3.1.3	Thermal Melting Curve of Reassociated DNA	48
3.1.4	Separation of DNA	48
3.1.5	Assay of Potential Contaminants in Labelled and Unlabelled DNA	49
3.2	Second Preparation of Labelled DNA	49
3.2.1	Absorbance spectra and Attempted Hybridisation	49
3.2.2	Assay of Potential Contaminants in the Second Preparation of labelled DNA	51
3.2.3	Stripping of unlabelled DNA	51
3.2.4	Ultra centrifugation of Sonicated and Stripped ^{32}P -DNA	51

3.3	Purification of DNA by the Hydroxyapatite-Urea Procedure	53
3.3.1	Purification of DNA Prepared by the Phenol-Chloroform Procedure	53
3.3.2	Preparation of DNA by the Hydroxyapatite-Urea Procedure.	56
3.3.3	Assay of Contaminants in DNA Prepared by the Hydroxyapatite-Urea Procedure .	56
3.3.4	Suitability for DNA for Hybridisation .	57
3.4.	Hybridisation Conditions	57
3.4.1	The Effect of Phosphate Buffer Molarity on the Separation of Single and Double-Stranded DNA with Hydroxyapatite .. .	57
3.4.2	Effect of Cot on DNA Reassociation ..	59
3.4.3	Experiments to Decrease the Apparent Self-Association of labelled DNA .. .	62
4.	DNA Homology Among Strains of <u>R. trifolii</u> and Related Species	63
4.1	Labelled DNA Preparations	63
4.1.1	Specific Activities	63
4.1.2	Hybridisation-Self Association of Labelled DNA, Homologous Reaction, and Preparation of Unlabelled DNA	64
4.2	Thermal Stability of DNA Reassociated at 65°C.	
4.2.1	Thermal elution profiles	64
4.2.2	Thermal Binding Indices	67
4.3	Homologies Among Related Strains	67
4.3.1	Serologically Related Strains	67
4.3.2	Non-Modulating Variants	71
4.3.3	Different Histories of Reisolates .. .	71
4.3.4	Same Host Origin	71
4.4	DNA Homology and Plant Specificity	76
4.4.1	Homologies Among the European Clover Rhizobia	76
4.4.2	Homologies Between <u>R. leguminosarum</u> and the European Clover Rhizobia .. .	76
4.4.3	Homologies Between American Clover Rhizobia and the European Clover Rhizobia	76
4.4.4	Homologies Between <u>Trifolium ambiguum</u> Rhizobia and European Clover Rhizobia	77
4.4.5	Homologies Between African Clover Rhizobia and European Clover Rhizobia	77
4.4.6	Homologies Between Japanese Clover Rhizobium and European Clover Rhizobia	81
4.4.7	Homologies Between <u>Rhizobium Phaseoli</u> and European Clover Rhizobia	81
4.4.8	Homologies Between <u>Rhizobium Phaseoli</u> and other groups of Rhizobia	82

DISCUSSION

	Page
1. Growth of <i>Rhizobium</i>	85
1.1 Growth Curves	85
1.2 Improvements in Cell Yield	86
2. Molecular Weight Determination of DNA	87
3. Difficulties Encountered in Hybridisation	88
3.1 Preparation of DNA	88
3.1.1 Phenol-chloroform Method	88
3.1.2 Hydroxyapatite-Urea Method	88
3.1.1 Effect of Polysaccharide in the Labelled DNA Preparation on Hybridisation	89
3.2 Optimisation of Hybridisation	90
3.2.1 Effective Phosphate Buffer Molarity on the Separation of Single and Double Stranded DNA with Hydroxyapatite	90
3.2.2 Effect of Cot on the Reassociation of DNA	91
3.2.3 Zero-time Binding of Denatured Labelled DNA to Hydroxyapatite	92
4. DNA Homology Among Strains of <i>Rhizobium trifolii</i> and related species	93
4.1 Thermal Stability of DNA Reassociated at 65°C	93
4.2 Relationships Between Specific Pairs of Strains	94
4.2.1 Non-nodulating variants	94
4.2.2 Serologically Related Strains	95
4.2.3 Historical Aspects	95
4.3 Homology and Plant Specificity	96
4.3.1 Relationships Within Sub-groups of <u><i>R. trifolii</i></u>	96
4.3.2 Relationship Between <u><i>R. Leguminosarum</i></u> and the European Clover Rhizobia	97
4.3.3 Relationships Between <u><i>R. Phaseoli</i></u> and European Clover Rhizobia	98
4.3.4 Proposed Classification of the Fast- Growing Rhizobia	99
4.3.5 Evolutionary Considerations	101
4.3.6 Plasmids in <u><i>Rhizobium</i></u>	103
BIBLIOGRAPHY	104

LIST OF FIGURES

	Page
1. Apparatus for the preparation of linear sucrose gradients	27
2. Colourimetric determination of DNA concentration by the method of Burton (1956)	31
3. Colourimetric determination of protein,	
a) by the method of Bradford (1976)	34
b) by the method of Lowry <u>et al.</u> (1976)	34
4. Colourimetric determination of polysaccharide concentration,	
a) by the Method of Trevelyn and Harrison (1952)	35
b) by the method of Bailey (1967)	35
5. Colourimetric determination of RNA concentration by the method of Ashwell (1957)	36
6. Growth of <u>Rhizobium</u> in yeast mannitol broth with and without dipotassium Hydrogen Phosphate	
a) <u>Rhizobium trifolii</u> strain TA1	41
b) <u>Rhizobium trifolii</u> strain WU290.	41
c) <u>Rhizobium phaseoli</u> strain TA101	41
7. Effect of increasing yeast extract concentration on the growth of <u>Rhizobium trifolii</u> strains 1/6, WU290, and <u>Rhizobium phaseoli</u> strain TA101	43
8. Effect of different batches of yeast extract on the growth of <u>Rhizobium trifolii</u> strain TA1:	
a) as measured by cellular protein	44
b) as measured by cell dry weight.	44
9. The effect of probe diameter on the sonication of DNA.	46
10. The effect of ^{32}P -labelling on the molecular weight of sonicated DNA.	47
11. Melting curve of incubated homologous DNA from <u>Rhizobium trifolii</u> strain cc275e.	50
12. Ultracentrifugation of sonicated and stripped ^{32}P -DNA from <u>Rhizobium trifolii</u> strain cc275e.	54
13. Effect of molar concentration of sodium phosphate buffer (pH 6.8) containing 0.4% sodium lauryl sulphate on the separation of homologous reassociated and labelled unreassociated DNA as determined by the hydroxyapatite batch centrifugation method.	60

14. Effect of equivalent Cot value on the reassociation of homologous and heterologous DNA with reference DNA extracted from Rhizobium trifolii strain TA1. 61
15. Thermal elution profiles of reassociated DNA duplexes formed at 65°C between DNA from Rhizobium trifolii strain TA1 and DNA from other strains of Rhizobium trifolii. 66
16. Thermal elution profiles of reassociated DNA duplexes, formed at 65°C, between DNA from Rhizobium phaseoli strain cc511 and DNA from strains of R. phaseoli and R. trifolii 68
17. The relationship between hybridisation at 65°C and the difference in melting point ($\Delta T_m(e)$) of reassociated homologous and heterologous*Rhizobium trifolii strain TA1 and R. phaseoli strain cc511 and DNA from R. Phaseoli, R. leguminosarum and R. trifolii 69
 * duplexes formed between reference DNAs from
18. Relationship between thermal binding indices (TBI) and the difference in melting point ($\Delta T_m(e)$) of reassociated homologous and heterologous duplexes formed between reference DNA from* strains of R. phaseoli, R. leguminosarum and R. trifolii. 70
 * Rhizobium phaseoli strain cc511 and DNA from
19. Relationships between the relative hybridisation (%) at 65°C of DNA from strains of rhizobia and reference DNAs from Rhizobium trifolii strains cc275e and TA1 78
20. Relationship between the relative hybridisation (%) at 65°C of DNA from strains of rhizobia and reference DNAs from Rhizobium trifolii strain cc275e and Rhizobium phaseoli strain cc511 79
21. Relationship between the relative hybridisation (%) at 65°C of DNA from strains of Rhizobia and reference DNAs from Rhizobium Trifolii strain TA1 and Rhizobium phaseoli strain cc511 80
22. Genetic relationship between strains of Rhizobium from plants in the genera Trifolium, Vicia, Pisum and from Phaseolus vulgaris 100

LIST OF TABLES

	Page
I, Species of <u>Rhizobium</u> and their characteristics (After Jordan and Allen, 1974)	3
II. Species of <u>Rhizobium</u> and their characteristics (obtained from Graham, 1976; as proposed by De Ley, 1968)	9
III. Strains of <u>Rhizobium</u> included in this study and their ability to nodulate and fix atmospheric nitrogen (effectiveness) on two species of <u>Trifolium</u>	18
IV. Assays of labelled and unlabelled DNA prepared by the phenol-chloroform method.	52
V. Effect of hydroxyapatite-urea purification on the absorbance spectra of two samples of DNA prepared by the phenol-chloroform procedure ...	55
VI. Composition of fractions obtained in preliminary trials of the hydroxyapatite-urea procedure for DNA preparation	58
VII. Relative hybridisation of TA1 reference DNA with unlabelled DNA prepared by the phenol-chloroform and the hydroxyapatite-urea procedures	65
VIII. Relative hybridisation of DNA strains of <u>Rhizobium trifolii</u> , <u>R. phaseoli</u> , <u>R. leguminosarum</u> and <u>Escherichia coli</u> with reference DNA from <u>R. trifolii</u> cc275e at 65°C and Cot 100... .. .	72
IX Relative hybridisation of DNA from strains of <u>Rhizobium Trifolii</u> , <u>R. phaseoli</u> , <u>R. leguminosarum</u> and <u>Escherichia Coli</u> with labelled reference DNA from <u>R. trifolii</u> TA1 after reassociation at 65°C and Cot 250	74
X. Relative hybridisation of DNAs from strains of <u>Rhizobium trifolii</u> , <u>R. phaseoli</u> , <u>R. leguminosarum</u> , and <u>Escherichia Coli</u> with reference DNA from <u>R. phaseoli</u> cc511 after reassociation at 65°C or 80°C at Cot 250	83