

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

TOOTH DETERIORATION IN SHEEP

A STUDY OF FARMS IN SOUTHERN NORTH ISLAND OF NEW ZEALAND

A thesis presented in partial fulfilment of

the requirements for the degree of

Master of Veterinary Science

at Massey University

GERALD DOUGLAS MITCHUM

December, 1985

## ACKNOWLEDGEMENTS

I am grateful to the people and the government of New Zealand for the opportunity to live and study in this country.

Further I would like to thank those who have been directly involved in the completion of this thesis.

Firstly to my supervisor, Professor A.N.Bruere, Department of Veterinary Clinical Sciences, Massey University, to whom I am indebted for the use of facilities and access to farms provided. Without his keen interest and critical approach to veterinary problems, the need for continued research in the area of excessive tooth wear in sheep may well have remained unrecognized for many years.

Secondly my thanks to Dr.B.C.Farquharson for his encouragement and valuable assistance in both the formulation of ideas and the technical areas of analytical chemistry, statistics and computer operation and to Dr.D.M.West for his knowledgeable advice and genuine concern in the early stages of this project.

Thirdly my sincere appreciation to Dr.M.Baxter, Dr.A.N.MacGregor, Mr.J.V.Pauli and Mr.C.B.Christie for their enthusiastic contributions in the areas of microbiology, soil science, clinical pathology and horticulture respectively. It was a great encouragement to be able to utilize the expertise of these men.

I also wish to thank Elizabeth Davies, Linley Denby, Jan Schrama, Pam Slack and Peter Winter for their willing assistance in their specific areas and Rose Law for her cheerful aid in organizing materials. My thanks also to Tom Law for his very capable help in the area of photography.

A special thank you to the farmers and their employees of the six farms included in this study. Their interest in this problem and the total sheep industry helped to make this project a very fulfilling endeavor.

Finally I wish to recognize my wife, Frances, for her help in proof reading and typing as well as her encouragement throughout this entire project.

## TABLE OF CONTENTS

	Page
Acknowledgements	
Table of Contents	
List of Tables	
List of Figures	
CHAPTER 1: INTRODUCTION	1
CHAPTER 2: A REVIEW OF MAJOR DENTAL PROBLEMS IN NEW ZEALAND SHEEP	4
CHAPTER 3: A FIELD INVESTIGATION INTO EXTREME DENTAL DETERIORATION: A FARM STUDY NEAR WAIPUKURAU	
Introduction	21
Farm Profile	22
History	22
The Initial Problem	23
Clinical Observations of Teeth on Various Age Groups of Sheep	25
1. Lambs	25
2. Hoggets	25
3. Two-Tooth Ewes	27
4. Mixed Age Ewes	30

Methods Used to Investigate Some of the Clinical Observations Made	32
1. The Percentage of Faecal Ash	32
2. The Correlation Between Percentage of Faecal Ash-Percentage of Acid Insoluble Residue	34
3. Observations on the Eruption Times of the Central Permanent Incisors	34
4. Wear of the Central Permanent Incisors	34
Results	35
1. The Percentage of Faecal Ash	35
2. The Correlation Between Percentage of Faecal Ash-Percentage of Acid Insoluble Residue	38
3. Observations on the Eruption Times of the Central Permanent Incisors	38
4. Wear of the Central Permanent Incisors	40
Discussion	41
1. Discussion of Management	41
a. Change of Breed	41
b. Change in the Grazing System	41
c. Intensification of Sheep Farming	46
d. Increase in Stocking Rate	46
2. Discussion of Clinical Observations	47
a. Sensitivity of Deciduous Incisors	47
b. Enamel Defects	47
c. Demineralization of Tips of Central Permanent Incisors	49
d. Disappearance of Shallow Data Points	49
3. Discussion of the Percentage of Faecal Ash	49
4. Discussion of the Correlation Between Percentage of Faecal Ash-Percentage of Acid Insoluble Residue	50
5. Discussion of Eruption Times of the Central Permanent Incisors	51

6. Discussion of Wear of the Central Permanent Incisors	51
CHAPTER 4: OBSERVATIONS AND TRIALS ON FARMS: AN INVESTIGATION INTO TOOTH WEAR ON FIVE NORTH ISLAND FARMS	
Introduction	53
Farm Profiles	53
1. The Tangimoana Farm	53
2. The Riverside Farm	54
3. The Pohangina Farm	54
4. The Ranui Farm	55
5. The Motukai Farm	55
Clinical Observations	56
1. The Tangimoana Farm	56
2. The Riverside Farm	56
3. The Pohangina Farm	56
4. The Ranui Farm	57
5. The Motukai Farm	58
Method of Trials on the Five Farms	58
Results of the Trials of the Five Farms	59
1. The Tangimoana Farm	59
a. The Percentage of Faecal Ash	59
b. The Correlation Between Percentage of Faecal Ash—Percentage of Acid Insoluble Residue	60
c. Observations on the Eruption Times of the Central Permanent Incisors	60
d. The Wear of the Central Permanent Incisors	60

e. Comparison of Measurements	61
2. The Riverside Farm	68
a. The Percentage of Faecal Ash	68
b. The Correlation Between Percentage of Faecal Ash—Percentage of Acid Insoluble Residue	68
c. Observations on the Eruption Times of the Central Permanent Incisors	68
d. The Wear of the Central Permanent Incisors	69
3. The Pohangina Farm	75
a. The Percentage of Faecal Ash	75
b. The Correlation Between Percentage of Faecal Ash—Percentage of Acid Insoluble Residue	75
c. Observations on the Eruption Times of the Central Permanent Incisors	75
d. The Wear of the Central Permanent Incisors	75
4. The Ranui Farm	81
a. The Percentage of Faecal Ash	81
b. The Correlation Between Percentage of Faecal Ash—Percentage of Acid Insoluble Residue	81
c. Observations on the Eruption Times of the Central Permanent Incisors	81
d. The Wear of the Central Permanent Incisors	81
5. The Motukai Farm	87
a. The Percentage of Faecal Ash	87
b. The Correlation Between Percentage of Faecal Ash—Percentage of Acid Insoluble Residue	87
c. Observations on the Eruption Times of the Central Permanent Incisors	87



d. The Wear of the Central Permanent Incisors	87
Discussion of Farm Trials	93
1. The Percentage of Faecal Ash	93
2. The Correlation Between Percentage of Faecal Ash-Percentage of Acid Insoluble Residue	94
3. The Eruption Times of Central Permanent Incisors	95
4. Wear of Central Permanent Incisors	97
A Summary of the Important Farm Trial Observations	100
1. Enamel Defects	100
2. Unexplained Clinical Signs	101
3. The Correlation Between Soil Ingestion and Tooth Wear	101
4. High Soil Ingestion on Low Wear Farms	102
5. Variation in Relationship of Stocking Rate to Tooth Wear	102
CHAPTER 5: SOLUBILIZATION OF SHEEP'S TEETH: AN ALTERNATIVE HYPOTHESIS ON THE LOSS OF TOOTH SUBSTANCE	
Introduction	104
Hypothesis	105
Demonstrations	106
1. Demonstration 1	106
2. Demonstration 2	107
3. Demonstration 3	107
Summary of Solubilization Hypothesis	112

CHAPTER 6: DISCUSSION AND CONCLUSIONS ON TOOTH DETERIORATION IN SHEEP	113
--	-----

#### APPENDICES

I Dry-Ashing in an Open Vessel	120
II Acid-Insoluble Ash	121

#### REFERENCES

## LIST OF TABLES

	Page
Table I: Stock units per hectare, total number of cattle, lambing percentage and percentage of dry ewes on the Waipukurau farm are listed from 1975 to 1983.	23
Table II: Fertilizer and lime applied to the Waipukurau farm from 1973 to 1983.	24
Table III: The percentage of faecal ash of faecal samples collected from September 1983 to September 1984 on the Waipukurau farm with mean and standard deviation.	36
Table IV: The correlation between the percentage of faecal ash and percentage of acid insoluble residue on the Waipukurau farm.	39
Table V: The eruption of central permanent incisors of 100 hoggets on the Waipukurau farm with the first day of lambing day one.	40
Table VI: Tooth wear in cm on the Waipukurau farm with the mean wear and mean daily wear calculated and the eruption date and enamel defects recorded.	42-45
Table VII: The percentage of faecal ash of faecal samples collected from October 1983 to September 1984 on the Tangimoana farm with mean and standard deviation.	62

Table VIII:	The correlation between the percentage of faecal ash and percentage of acid insoluble residue on the Tangimoana farm.	64
Table IX:	The eruption of central permanent incisors of 100 hoggets on the Tangimoana farm with the first day of lambing day one.	65
Table X:	Tooth wear in cm on the Tangimoana farm with the mean wear and mean daily wear calculated.	66
Table XI:	Correlation of two sets of measurements of central permanent incisors in sheep on the Tangimoana farm re-measured and recorded on the same day.	67
Table XII:	The percentage of faecal ash of faecal samples collected from October 1983 to May 1984 on the Riverside farm with mean and standard deviation.	70
Table XIII:	The correlation between the percentage of faecal ash and percentage of acid insoluble residue on the Riverside farm.	72
Table XIV:	The eruption of central permanent incisors of 100 hoggets on the Riverside farm with the first day of lambing day one.	73
Table XV:	Tooth wear in cm on the Riverside farm with the mean wear and mean daily wear calculated.	74

Table XVI:	The percentage of faecal ash of faecal samples collected from October 1983 to September 1984 on the Pohangina farm with mean and standard deviation.	76
Table XVII:	The correlation between the percentage of faecal ash and percentage of acid insoluble residue on the Pohangina farm.	78
Table XVIII:	The eruption of central permanent incisors of 100 hoggets on the Pohangina farm with the first day of lambing day one.	79
Table XIX:	Tooth wear in cm on the Pohangina farm with the mean wear and mean daily wear calculated.	80
Table XX:	The percentage of faecal ash of faecal samples collected from October 1983 to August 1984 on the Ranui farm with mean and standard deviation.	82
Table XXI:	The correlation between the percentage of faecal ash and percentage of acid insoluble residue on the Ranui farm.	84
Table XXII:	The eruption of central permanent incisors of 100 hoggets on the Ranui farm with the first day of lambing day one.	85
Table XXIII:	Tooth wear in cm on the Ranui farm with the mean wear and mean daily wear calculated.	86
Table XXIV:	The percentage of faecal ash of faecal samples collected from October 1983 to August 1984 on the Motukai farm with mean and standard deviation.	88

Table XXV:	The correlation between the percentage of faecal ash and percentage of acid insoluble residue on the Motukai farm.	90
Table XXVI:	The eruption of central permanent incisors of 100 hoggets on the Motukai farm with the first day of lambing day one.	91
Table XXVII:	Tooth wear in cm on the Motukai farm with the mean wear and mean daily wear calculated.	92
Table XXVIII:	A comparison of the mean percentage of faecal ash for the six farms included in this investigation with relative tooth wear status.	95
Table XXIX:	A comparison of the mean percentage difference between faecal ash and acid insoluble residue from the six farms in this investigation.	96
Table XXX:	The mean eruption time of the central permanent incisors in hoggets on the six farms involved in this investigation.	97
Table XXXI:	Calculated daily wear in cm of central permanent incisors in two-tooth ewes on the six farms involved in this investigation.	98
Table XXXII:	The numbers of colonies from soil at progressive dilutions from the Tangimoana farm and the Pohangina farm.	108

Table XXXIII: Measurements and weights of tomato plants after 82 days. Pots I and II contained powdered sheep's teeth and dextrose. Pots III and IV contained dextrose and no powdered sheep's teeth. Pots V and VI contained powdered sheep's teeth and no dextrose and pot VI had autoclaved soilless potting media.

## LIST OF FIGURES

	Page
Fig. 1: Severely deteriorated deciduous incisors in a 14 month old ewe hogget on the Waipukurau farm.	26
Fig. 2: Severely deteriorated deciduous incisors in a 14 month old ewe hogget on the Waipukurau farm with obvious involvement of the entire exposed tooth surface.	26
Fig. 3: Deciduous incisors in a 14 month old ewe hogget from the Tangimoana farm where tooth deterioration is not a problem.	28
Fig. 4: Enamel defects on central permanent incisors in a two-tooth ewe on the Waipukurau farm.	28
Fig. 5: Enamel defects on central permanent incisors in a two-tooth ewe on the Waipukurau farm.	29
Fig. 6: Linear enamel defects on the central permanent incisors of a four-tooth ewe on the Waipukurau farm.	29
Fig. 7: A data point nearly missing on a central permanent incisor with considerable tooth exposed below the point.	31
Fig. 8: Deciduous incisors which have been soaked in concentrated citric acid for 24 hours.	31
Fig. 9: The incisors of a four year old ewe on the Waipukurau farm representative of the older sheep in this flock.	33



Fig. 10:	The incisors of a four-tooth ewe on the Waipukurau farm with extensive deterioration of the central permanent incisors.	33
Fig. 11:	Data point on central permanent incisor with a visible transparent line on the tips.	35
Fig. 12:	A graph showing the mean percentage faecal ash with standard deviation for the period from September 1983 to September 1984 on the Waipukurau farm.	37
Fig. 13:	A graph showing the mean percentage faecal ash with standard deviation for the period from October 1983 to September 1984 on the Tangimoana farm.	63
Fig. 14:	A graph showing the mean percentage faecal ash with standard deviation for the period from October 1983 to May 1984 on the Riverside farm.	71
Fig. 15:	A graph showing the mean percentage faecal ash with standard deviation for the period from October 1983 to August 1984 on the Pohangina farm.	77
Fig. 16:	A graph showing the mean percentage faecal ash with standard deviation for the period from October 1983 to August 1984 on the Ranui farm.	83
Fig. 17:	A graph showing the mean percentage faecal ash with standard deviation for the period from October 1983 to August 1984 on the Motukai farm.	89

- Fig. 18: Photograph of solubilized  $\text{CaHPO}_4$  precipitate. 109
- Fig. 19: A. Photograph of culture plates of soil at varying dilutions from the Pohangina farm.  
B. Photograph of culture plates of soil from the Tangimoana farm at the same dilutions as the Pohangina plates. 109
- Fig. 20: Photograph showing relative size of the tomato plants receiving the various treatments. 111