

1 **Detecting Genes Associated with Pathogenicity and Antimicrobial**

2 **Resistance in Three New Zealand Waterways.**

3

4

5 *Meredith T. Davis,^{1,2} Anne C. Midwinter,² Russell G. Death,¹ Richard Cosgrove,³ and*

6 *Richard C. Winkworth⁴*

7

8 ¹ School of Agriculture and the Environment - Ecology, Massey University, Palmerston

9 North, New Zealand

10 ² Molecular Epidemiology and Veterinary Public Health Laboratory - Hopkirk Research

11 Institute, School of Veterinary Science, Massey University, Palmerston North, New Zealand

12 ³ Fish and Game New Zealand - 32 Richard Pearse Drive, Temuka, New Zealand

13 ⁴ School of Fundamental Sciences, Massey University, Palmerston North, New Zealand

14

15

16

17 Corresponding Author:

18 Meredith T. Davis

19 E-mail: M.Davis@massey.ac.nz

20 School of Agriculture and the Environment, Massey University, Private Bag 11 222,

21 Palmerston North, 4442, New Zealand.

22

23

24 Competing Financial Interests:

25 The authors declare they have no actual or potential competing financial interests.

26 **BACKGROUND:** More than 100 zoonoses may be transmitted via water, among them enteric
27 diseases are leading causes of human mortality. Traditional monitoring for zoonoses relies on
28 culturing of indicator species, but environmental DNA (eDNA) provides an alternative,
29 allowing direct testing for genetic loci associated with pathogenicity and/or antimicrobial
30 resistance in zoonotic bacteria.

31 **OBJECTIVES:** To evaluate whether genes associated with Shiga toxin producing *Escherichia*
32 *coli* (STEC) and antimicrobial resistance can be monitored in waterways using culture-free
33 sampling of eDNA combined with PCR-based testing.

34 **METHODS:** Water and sediment samples were collected from two sites on each of three rivers
35 in Canterbury, New Zealand; sample sites were situated above and below reaches bordered
36 by intensive dairy farming. Samples from each site were tested for genes typically associated
37 with *Escherichia coli*, STEC serogroups O26 and O157, human pathogenic strains of STEC,
38 and resistance to a broad range of antibiotics.

39 **RESULTS:** Both culturing and genetic testing confirmed the presence of *E. coli* in all samples.
40 In contrast, presence of genes associated with STEC and antibiotic resistance varied by
41 season and substrate. The O157 serogroup was identified at more than twice as many sites as
42 O26, with the latter more common in autumn samples. In autumn, genes associated with
43 pathogenic STEC were detected in one Ashley and both Rangitata River samples but were
44 present in all spring samples, except one Ashley and one Selwyn River collection. The
45 antibiotic resistance gene was only identified in spring, predominantly at sites downstream of
46 intensive dairying.

47 **DISCUSSION:** While our sample is small this study indicates that genetic testing of eDNA can
48 be a useful tool for monitoring the presence and persistence of zoonoses in waterways. How
49 the presence of these genetic elements is related to that of pathogenic STEC and incidence of
50 disease in humans now needs to be examined.

52 **Introduction**

53 Globally, the degradation of freshwater ecosystems due to anthropogenic land use changes
54 has been linked to outbreaks of waterborne disease in humans (OECD 2017).
55 Industrialisation (Closs et al. 2016; Ratha 2019), urbanisation (Merritt et al. 2010; Power et
56 al. 2018), and intensive agriculture (Clapcott et al. 2012; Gluckman 2017; Joy 2015) all have
57 the potential to negatively affect freshwater ecosystems through removal of riparian
58 vegetation (Hickey and Doran 2004; Sweeney and Newbold 2014), heavy metal
59 accumulation (Hickey and Clements 1998), increased sediment and nutrient inputs (Waters
60 1995; Yan et al. 2016), and contamination with faecal effluent (Dangendorf 2004). In New
61 Zealand rapid changes in land use (Julian et al. 2017), the degradation of freshwater
62 ecosystems (Clapcott et al. 2012), and steady increases in outbreaks of notifiable bacterial
63 zoonoses are all well documented (Ministry for the Environment and Statistics New Zealand
64 2017).

65 Enteric diseases, which may be waterborne, are a leading cause of human deaths
66 worldwide, especially in children under five years of age in lower income countries (World
67 Health Organization 2019). Currently more than 100 zoonoses are recognised as being
68 transmitted in aquatic ecosystems. These include members of *Leptospira*, *Campylobacter*,
69 *Escherichia*, and *Salmonella* (Fang 2014; Gluckman 2017; Shaw et al. 2016). Human and
70 animal faecal effluent are primary sources of the microorganisms responsible for waterborne
71 zoonoses (Reddy et al. 1981). In broad terms, the risk of contracting bacterial zoonoses from
72 effluent contaminated waterways increases for those swimming in (McBride et al. 2002),
73 gathering food from (Perkins et al. 2016; Rose et al. 2001) or eating undercooked produce
74 irrigated with (Adator et al. 2018; King et al. 2012; Solomon et al. 2002) effluent
75 contaminated water. However, in specific terms, the risks associated with using effluent

76 contaminated recreational waterways remain poorly understood because they are complex,
77 dynamic and multi-factorial (Colford Jr et al. 2007; Prieto et al. 2001).

78 Shiga toxin producing *Escherichia coli* (STEC) is an emerging group of zoonoses
79 linked to ruminants (Colford Jr et al. 2007; Oporto et al. 2019) that may occur in in
80 recreational waters and to be transmitted via the oral-faecal route (Swaggerty et al. 2018),
81 STEC is the fourth most commonly reported zoonosis in the EU (Severi et al. 2016) and USA
82 (European Center for Disease Prevention and Control 2018) and an emerging issue in New
83 Zealand (ESR 2019). In humans STEC infections typically present as enteric disease that in
84 severe cases may progress to haemolytic uraemic syndrome (HUS) or kidney failure
85 (Karmali 2018). Strains of *E. coli* recognised as belonging to the STEC group are generally
86 characterised by the presence of one or a pair of Shiga toxin producing genes (*stx1* and *stx2*)
87 (Donnenberg et al. 1997; Perna et al. 1998). However, these strains are otherwise diverse;
88 belonging to 129 serogroups and with more than 260 antigen combinations (Valilis et al.
89 2018). Typically, human pathogenic STEC strains contain the locus of enterocyte and
90 effacement (LEE) island (Paton and Paton 1998) and, until recently, were most likely to be
91 identified as belonging to the O157 serogroup (Centers for Disease Control and Prevention
92 2018; European Center for Disease Prevention and Control 2018). In New Zealand human
93 cases of STEC disease tripled between 2014 and 2017 (Health and Environment Group ESR
94 2019). However, over this same period the proportion of cases linked to the O157 serogroup
95 fell from 88.1% to 37.1% (Health and Environment Group ESR 2019). Non-O157 STECs
96 (e.g., O26, O45, O111) are increasingly being linked to disease in humans (Germinario et al.
97 2016; Gill et al. 2019; Luna-Gierke et al. 2014; Severi et al. 2016). This observation may
98 simply reflect improved screening (Health and Environment Group ESR 2019) or may be due
99 to changes in serotype prevalence.

100 Culture for human pathogenic STEC involves the isolation of candidate organisms
101 based on their ability to metabolise different carbohydrates (Amézquita-López et al. 2018)
102 which is then followed by molecular characterisation. Molecular characterisation of STEC
103 typically focuses on genes associated with virulence (i.e., *stx1* and *stx2*, *eae* located on the
104 LEE island) and the serotypes most commonly found in human cases (e.g., *rfbE* for O157,
105 *wzy* for O26) (Anklam et al. 2012; Franz et al. 2007). However, in most studies of
106 environmental samples, the microbial communities are enriched, using growth media, for 18-
107 24 hours prior to plating and identification (De Boer and Heuvelink 2000; Irshad et al. 2016).
108 Enriching samples in this way is not ideal as it alters the composition of the original
109 microbial community; effects may be due to contrasting media preferences (Amagliani et al.
110 2018) or interactions between community members (Chekabab et al. 2013; Mauro et al.
111 2013). At worst such changes may result in failure to detect potentially pathogenic
112 community members.

113 Use of traditional culturing is further complicated in the case of STEC as strains may
114 lose genes associated with virulence during incubation for enrichment or culturing
115 (Senthakumaran et al. 2018; Tarr et al. 2019). An alternative to the traditional approach is to
116 test DNA extracted directly from environmental samples for the presence of genes associated
117 with virulence or antibiotic resistance (Bélanger et al. 2002; Werber et al. 2002). Testing of
118 environmental DNA (eDNA) can provide a rapid and inexpensive survey of human
119 pathogens at a site. This approach could be used to examine whether genes typically
120 associated with pathogenic STEC (e.g., *stx1*, *stx2* and *eae*) are present at a given site.
121 Although this approach does not identify pathogenic isolates, the presence of virulence genes
122 in a given sample should raise concerns about the potential level of faecal contamination at
123 that site. This approach allows us to identify sites where further investigation, including the
124 isolation and testing of individual microbes, is warranted.

125 In this study we use a culture free, eDNA method to test for genes associated with
126 STEC from sites on three Canterbury, New Zealand rivers. Samples were collected to
127 coincide with seasonal peaks in human STEC cases during autumn and spring (Health and
128 Environment Group ESR 2019). We tested both benthic sediments and water column samples
129 for five genes frequently associated with human pathogenic STEC and the recurrently co-
130 morbid, group 1 CTX-M β -lactamases which confer resistance to multiple antibiotics (Ishii et
131 al. 2005; Valat et al. 2012). For each sample we also performed traditional *E. coli* counts to
132 determine whether there was any relationship between colony numbers and the prevalence of
133 these genes.

134

135 **Materials and Methods**

136 ***Sample Collection***

137 Water and sediment samples were collected from the Ashley, Rangitata, and Selwyn rivers
138 once in austral autumn and once in the spring of 2018. Collections were made at two sites
139 along each river; these sites were 10-15 km apart with one above and the other below reaches
140 bordered by high densities of intensive dairying (Fig.1). At each site, water and benthic
141 sediment samples were collected into separate sterile containers. Three 1 L water and three
142 25 g sediment samples were collected at each site. Samples were packed on ice, transported
143 to the laboratory, and processed within 24 hours of collection.

144

145 ***Sample Processing for Bacterial Culturing***

146 Water column sample aliquots were diluted 1:10, 1:5, and 1:2.5 with sterile MilliQ H₂O to a
147 final volume of 50 ml. At sites where the level of suspended sediments was high, an
148 additional 1:50 dilution was also prepared to ensure ease of enumeration. For each sample we
149 prepared three technical replicates at each dilution (i.e., 9-12 dilutions per sample).

150 For sediment samples, 2 g of wet sediment was first transferred to a 5 ml microtube, 3
151 ml MilliQ H₂O added, and the mixture then agitated vigorously for 30 seconds. Aliquots of
152 the resulting supernatant were diluted 1:500, 1:200, 1:100, and 1:50 with MilliQ H₂O to a
153 final volume of 50 ml. For each sample we prepared three technical replicates at each dilution
154 (i.e., 12 dilutions per sample).

155 For each dilution the total 50 ml volume was vacuum filtered through a single sterile
156 0.45 µm cellulose ester membrane filter (Merck KGaA, Darmstadt, Germany).

157

158 ***Bacterial Culturing***

159 Bacterial culturing followed United States Environmental Protection Agency method 1603
160 (EPA 2002). Each filter was placed onto a Difco Modified mTEC Agar (VWR, Radnor, PA)
161 plate, incubated at 37.5 °C for two hours, and then incubated at 45 °C for 18-20 hours.
162 Following incubation colonies resembling *E. coli* (red/magenta colonies) were counted.

163

164 ***Sample Processing for Molecular Testing***

165 Three 500 ml aliquots of water from each site were vacuum filtered through separate 0.45 µm
166 cellulose ester membrane filters. Environmental DNA was extracted from half of each filter
167 using the NucleoSpin® soil kit (Machery-Nagel GmbH and Co. KG, Düren, Germany)
168 following the manufacturer's instructions. For each sediment sample eDNA was extracted
169 from three 0.5 g aliquots of wet sediment, using the NucleoSpin® soil kit.

170

171 ***Molecular Testing for Specific Genes***

172 The presence of five genes most commonly associated with human pathogenic STEC was
173 evaluated using a polymerase chain reaction (PCR). Specifically, we targeted genes
174 associated with serogroup specific antigen biosynthesis, *rfbE* for O157 and *wzy* for O26, the

175 *stx*₁ and *stx*₂ toxin genes, and the intimate attachment and effacing gene, *eae*, using the
176 primers reported by Anklam et al. (2012). We also investigated the presence of antimicrobial
177 resistance by targeting the *bla* gene associated with group 1 CTX-M β -lactamases using the
178 primers reported by Lalzampaia et al. (2013). Finally, as an amplification control we targeted
179 the beta-glucuronidase gene, *uidA*, which is present in most *E. coli* (Anklam et al. 2012).

180 Amplification reactions were performed in 20 μ l reaction volumes containing 0.5 \times iQ
181 PerfeCTa® qPCR ToughMix™, ROX™ (QIAGEN, Düsseldorf, Germany), 1 pM of each
182 primer, and 2.5 μ l of DNA template. Thermocycling was performed in a T1 thermocycler
183 (Biometra GmbH, Göttingen, Germany) using standard cycling conditions including an initial
184 denaturation at 94°C for 3 mins, followed by 35 cycles of 94°C for 30 s, 60°C for 30 s and
185 72°C for 1 min, with a final extension at 72°C for 5 mins. Amplification products were
186 visualised using SYBR Safe (ThermoFisher Scientific, Waltham, MA) following
187 electrophoresis on 2% Tris-acetate-ethylenediamine tetraacetic acid agarose gels.

188

189 **Results**

190 ***Bacterial Culturing***

191 With just one exception, the spring sampling at the above intensive dairy site on the Selwyn
192 River, *E. coli* counts were consistently higher from sediment samples than water column
193 samples (Table 1). Moreover, counts were also higher for all but one sample taken at below
194 intensive dairy sites. Only the spring sediment sample from the above intensive dairy site on
195 the Rangitata River had a higher *E. coli* count than the corresponding sample from the below
196 intensive dairy site.

197

198 ***Molecular Testing for Specific Genes***

199 The *uidA* locus was successfully amplified from every sample in this study (Table 2). In
200 contrast, detection of the three gene loci associated with pathogenic STEC (*stx1*, *stx2* and
201 *eae*) and the gene associated with antibiotic resistance (*bla* CTX-M) varied with location,
202 time, and substrate. The *stx1*, *stx2* and *eae* genes were more frequently detected in water
203 samples in the autumn but in sediment samples in the spring. At both sampling times these
204 three virulence genes were found together with the O26 (*wzy*) and O157 (*rfbE*) genes. The
205 O26 marker was more frequently recovered from water, while that for O157 was evenly
206 distributed between the substrates. The antibiotic resistance gene was detected in both
207 substrates in samples from the Selwyn and Rangitata rivers.

208 In autumn, the genes associated with pathogenic STEC O157 (i.e., *stx1*, *stx2*, *eae* and
209 *rfbE*) were present at 33% of sample sites. All three virulence and both serogroup genes were
210 detected in Rangitata River water and Ashley River sediment samples from below intensive
211 dairy. Additionally, the *stx1*, *eae*, and *rfbE* genes were present in water from the site below
212 intensive dairy on the Ashley River. No other virulence or serogroup genes were detected in
213 the autumn samples.

214 In spring *stx1*, *stx2* and *eae* genes were detected in all rivers. These three STEC
215 virulence genes were detected in sediment samples from five of the six sites (i.e., all except
216 the Ashley below intensive dairy) but only from below intensive dairy in the water samples;
217 only the *eae* gene was detected in water samples from above intensive dairy. In spring
218 samples, the gene associated with the O157 serogroup (*rfbE*) was detected four times more
219 often than that associated with the O26 serogroup (*wzy*) and the *bla* CTX-M gene was more
220 frequent below intensive dairy. In three samples *stx1*, *stx2* and *eae* were detected together; in
221 one of these the gene associated with the O157 serogroup was present but the genes
222 associated with O157 and O26 were not detected in the remaining two. The *stx1* and *eae*

223 genes were detected in two water samples from below intensive dairy whereas *stx2* and *eae*
224 were detected in three sediment samples, two from below and one above intensive dairy.

225

226 **Discussion**

227 The presence of human pathogenic STEC in New Zealand waterways is poorly documented.
228 In part, this is because STEC strains have diverse metabolic requirements; since strains may
229 co-occur, identifying individual organisms present in a sample would require multiple culture
230 methods (Kerangart et al. 2018; Possé et al. 2007). In studies on STEC strains from ruminant
231 faeces and environmental samples the microbial communities are generally enriched prior to
232 DNA-based testing (Browne et al. 2018; Irshad et al. 2016; Jaros et al. 2013). The potential
233 impacts of such enrichment on the microbial communities and on inferences concerning the
234 detection of zoonotic disease are difficult to quantify. In the present study, we tested samples
235 for genes associated with virulence and antibiotic resistance in STEC without an initial
236 enrichment step.

237 In our analyses the genetic markers indicative of *E. coli* (*uidA*) and STEC virulence
238 (*stx1*, *stx2* and *eae*) genes were detected in all three of the rivers sampled. That all samples
239 tested positive for the *uidA* gene is consistent with *E. coli* being isolated at all six sites, from
240 both substrates, and at both sampling times using traditional plating. Using relatively small
241 sample volumes we detected genes associated with both human pathogenic STEC and
242 antibiotic resistance without first enriching the microbial communities. Although all the
243 STEC virulence and O serotype genes were detected in all rivers, and in some cases all these
244 components were detected from a single sample, their presence varied by location, substrate,
245 and season. These results suggest that explanations for the distribution of these genes across
246 the landscape are likely to be complex and involve a range of factors.

247 *Escherichia coli* levels were higher and STEC associated genes more commonly
248 detected at sites below intensive dairy. For example, the *stx1* gene was detected in six of 12
249 (50%) samples, and at least once at every site, below intensive dairy but in only two of 12
250 (17%) samples taken above intensive dairy. These results are consistent with previous reports
251 suggesting that agricultural effluent is a primary source of faecal contamination in New
252 Zealand waterways (Gluckman 2017). Studies have shown that faecal bacteria are transferred
253 from pastures to waterways via run-off; carried either directly by the flow or indirectly as a
254 result of adsorption to soil particles (Byappanahalli and Ishii 2011; Muirhead et al. 2004;
255 Palmateer et al. 1993). However, studies have suggested that paddock-feeding waterfowl may
256 also transmit ruminant hosted faecal bacteria when environmental contamination is high
257 (Yang et al. 2019; Zou et al. 2019). In the current study we sampled adjacent to intensive
258 dairy operations but expected microbial communities to vary along each river due to the
259 diversity of land uses within each catchment. Microbial communities are likely to be most
260 strongly influenced by adjoining land use but also impacted by inputs from upstream of the
261 sampling location. Moreover, microbial communities reflect the prevailing environmental
262 conditions and primary land use within the catchment. For example, the *stx1* and *stx2* genes
263 were only detected at above intensive dairy sites during spring. One explanation for this
264 observation may be inputs from smaller farms upstream of the reach bordered by the high
265 densities of intensive dairying.

266 The STEC associated genes occurred at greater frequency in the spring sampling than
267 the autumn sampling. Specifically, they were detected in eight of 12 samples (66%) in the
268 spring and in three of 12 samples (25%) in the autumn. This result reflects a combination of
269 increased detection below intensive dairy (i.e., five in September and three in May) as well as
270 detection above intensive dairy in the spring. Increased detection of STEC and antibiotic
271 resistance during the spring is likely a response to several factors. Spring calving may

272 increase the load of pathogenic bacteria on land neighbouring these rivers. As calves have
273 poorly developed intestinal biomes, are stressed by weaning, or are removed from their
274 mother prior to receiving colostrum they are prone to colonization by and heavy shedding of
275 bacteria, including STEC (Browne et al. 2018). In addition to the increased faecal loading on
276 pastures, spring rainfall patterns may lead to higher faecal inputs reaching rivers. Other
277 possible explanations for fewer detections in the autumn include lower base flow rates or
278 drying out of rivers during the summer (e.g., the Selwyn), removal of stock from flood prone
279 areas for winter grazing, and geomorphology that encourages adsorption of bacteria to
280 sediments (Reddy et al. 1981). That STEC associated genes were more frequently detected
281 during the spring is in contrast to the seasonal cycle of human STEC disease cases in New
282 Zealand. Although there is a peak in notified human cases during spring, the autumn peak is
283 typically larger. For example, in 2017 there were almost twice as many cases of STEC in
284 autumn than during spring (Health and Environment Group ESR 2019). Moreover, although
285 rates of notified human cases of STEC do differ between regions the differences are not
286 always consistent with the prevalence of dairy farming. For example, in 2017 the Auckland
287 and Waikato districts reported 7.4 and 8.6 cases per 100,000 individuals despite dairy
288 farming being far more common in the Waikato region (Health and Environment Group ESR
289 2019). Taken together these observations suggest that livestock are unlikely to be the sole
290 source of human STEC infections in New Zealand and that we need to be cautious when
291 interpreting data on the distribution of STEC, or the genetic components linked to it, in the
292 context of public health.

293 In all three of the sampled rivers *E. coli* levels were consistently higher in sediments
294 than in the water column. The STEC associated genes were also more commonly detected in
295 sediment samples. These results are consistent with previous studies that indicate aquatic
296 sediments may act as a store for *E. coli* (Muirhead et al. 2004; Perkins et al. 2014; Wilkinson

297 et al. 1995). Given that such stores may persist for months or years (Anderson et al. 2005;
298 Garzio-Hadzick et al. 2010; Gerba and McLeod 1976) and that suspended sediments increase
299 *E. coli* levels in the water column (Davies-Colley et al. 2018; Weiskerger and Whitman
300 2018) this finding has potentially important implications for water monitoring. In New
301 Zealand, water quality testing by local government agencies is currently restricted to the
302 water column. Although recreational use of waterways is discouraged when levels of
303 suspended sediment are high (e.g., following precipitation) (Davies-Colley et al. 2018),
304 disturbance of sediments by recreational users is not generally considered. Further work is
305 needed to quantify the pathogens associated with the localised mixing of sediment into the
306 water column by recreational users and to determine whether zoonoses are being
307 underestimated by the sampling of a single substrate.

308

309 **Conclusion**

310 Microbial culturing is not an efficient tool for monitoring STEC in recreational waterways. In
311 part this can be attributed to the metabolic diversity of STEC and the apparent lack of a
312 relationship between faecal indicator bacteria (e.g., *E. coli* counts) and the presence of genes
313 associated with STEC. Understanding this relationship would require more intensive
314 sampling over a broader geographic range. In most cases monitoring is conducted on
315 samples retrieved from the water column. Such samples may not accurately reflect the
316 microbial community that recreational users of the waterway may be exposed to. Our results
317 suggest sediments may act as an important STEC reservoir and resuspension of these
318 sediments by waterway users could potentially increase exposure to STEC.

319 There is a growing appreciation of new technologies that enable the presence and
320 persistence of zoonoses to be monitored without the need for microbial culturing
321 (Byappanahalli and Ishii 2011; Rose et al. 2001). This represents a fundamental change in

322 our approach to microbial monitoring allowing us to take a holistic view of the riverine
323 environment and improving our ability to ensure environmental, animal, and human health.
324 While small, this study is the first step towards understanding zoonoses in New Zealand
325 waterways at a time when global health is under the microscope.

326

327 **Acknowledgements**

328 We thank Martin Taylor (Fish and Game New Zealand) for funding this study and providing
329 staff to collect samples.

330

331 **References**

- 332 Adator EH, Cheng M, Holley R, McAllister T, Narvaez-Bravo C. 2018. Ability of Shiga
333 toxigenic *Escherichia coli* to survive within dry-surface biofilms and transfer to fresh
334 lettuce. *Int J Food Microbiol* 269: 52–59, PMID: 29421358,
335 <https://doi.org/10.1016/j.ijfoodmicro.2018.01.014>.
- 336 Amagliani G, Rotundo L, Carloni E, Omiccioli E, Magnani M, Brandi G, et al. 2018.
337 Detection of Shiga toxin-producing *Escherichia coli* (STEC) in ground beef and bean
338 sprouts: evaluation of culture enrichment conditions. *Food Res Int* 103: 398–405, PMID:
339 29389629, <https://doi.org/10.1016/j.foodres.2017.10.059>.
- 340 Amézquita-López BA, Soto-Beltrán M, Lee BG, Yambao JC, Quiñones B. 2018. Isolation,
341 genotyping and antimicrobial resistance of Shiga toxin-producing *Escherichia coli*. *J*
342 *Microbiol Immunol Infect* 51: 425–434, PMID: 28778595,
343 <https://doi.org/10.1016/j.foodres.2017.10.059>.
- 344 Anderson KL, Whitlock JE, Harwood VJ. 2005. Persistence and differential survival of fecal
345 indicator bacteria in subtropical waters and sediments. *Appl Environ Microbiol* 71:
346 3041–3048, PMID: 15933000, <https://doi.org/10.1128/AEM.71.6.3041-3048.2005>.

- 347 Anklam KS, Kanankege KST, Gonzales TK, Kaspar CW, Döpfer D. 2012. Rapid and reliable
348 detection of Shiga toxin–producing *Escherichia coli* by real-time multiplex PCR. *J Food*
349 *Prot* 75: 643–650, PMID: 22488051, <https://doi.org/10.4315/0362-028X.JFP-11-392>.
- 350 Bélanger SD, Boissinot M, Ménard C, Picard FJ, Bergeron MG. 2002. Rapid detection of
351 Shiga toxin-producing bacteria in feces by multiplex PCR with molecular beacons on
352 the smart cycler. *J Clin Microbiol* 40: 1436–1440, PMID: 11923369,
353 <https://doi.org/10.1128/JCM.40.4.1436-1440.2002>.
- 354 Browne AS, Midwinter AC, Withers H, Cookson AL, Biggs PJ, Marshall JC, et al. 2018.
355 Molecular epidemiology of Shiga toxin-producing *Escherichia coli* (STEC) on New
356 Zealand dairy farms: application of a culture-independent assay and whole-genome
357 sequencing. *Appl Environ Microbiol* 84: e00481-18, PMID:29752274,
358 <https://doi.org/10.1128/AEM.00481-18>
- 359 Byappanahalli MN, Ishii S. 2011. Environmental sources of fecal bacteria. In: *The fecal*
360 *bacteria*, 93-110. Sadowsky MJ, Whiman RL, eds. Washington D.C., USA: ASM
361 Press,.
- 362 CDC (Centers for Disease Control and Prevention). 2018. *National enteric disease*
363 *surveillance: Shiga toxin-producing Escherichia coli (STEC) annual report, 2016*.
364 Atlanta: Centers for Disease Control and Prevention.
365 <https://www.cdc.gov/ncezid/dfwed/PDFs/national-stec-surveillance-overview-508c.pdf>
366 [accessed 6 April 2020].
- 367 Chekabab SM, Daigle F, Charette SJ, Dozois CM, Harel J. 2013. Shiga toxins decrease
368 enterohaemorrhagic *Escherichia coli* survival within *Acanthamoeba castellanii*. *FEMS*
369 *Microbiol Lett* 344: 86–93, PMID: 23581502, <https://doi.org/10.1111/1574-6968.12158>.
- 370 Clapcott JE, Collier KJ, Death RG, Goodwin EO, Harding JS, Kelly D, et al. 2012.
371 Quantifying relationships between land-use gradients and structural and functional

- 372 indicators of stream ecological integrity. *Freshw Biol* 57:74–90,
373 <https://doi.org/10.1111/j.1365-2427.2011.02696.x>.
- 374 Closs GP, Angermeier PL, Darwall WRT, Balcombe SR. 2016. Why are freshwater fish so
375 threatened. In: *Conservation of freshwater fishes*, 37-75 Closs GP, Kirkosek M, Olden
376 JD. Cambridge, England: Cambridge University Press.
- 377 Colford Jr JM, Wade TJ, Schiff KC, Wright CC, Griffith JF, Sandhu SK, et al. 2007. Water
378 quality indicators and the risk of illness at beaches with nonpoint sources of fecal
379 contamination. *Epidemiology* 18: 27–35, PMID: 17149140,
380 <https://doi.org/10.1097/01.ede.0000249425.32990.b9>.
- 381 Dangendorf F. 2004. Impacts of anthropogenic and environmental factors on the distribution
382 of zoonoses. In: *Waterborne zoonoses identification, causes, control*, 46–65. Cotruvo
383 JA, Dufour A, Rees G, Bartram J, Carr R, Cliver DO, et al. eds. London, UK: IWA
384 Publishing.
- 385 Davies-Colley R, Valois A, Milne J. 2018. Faecal contamination and visual clarity in New
386 Zealand rivers: correlation of key variables affecting swimming suitability. *J Water*
387 *Health* 16: 329–339, PMID: 29952322, <https://doi.org/10.2166/wh.2018.214>.
- 388 De Boer E, Heuvelink AE. 2000. Methods for the detection and isolation of Shiga
389 toxin-producing *Escherichia coli*. *Symp Ser Soc Appl Microbiol*. 29: 133S-143S,
390 PMID:10880188, <https://doi.org/10.1111/j.1365-2672.2000.tb05341.x>
- 391 Donnenberg MS, Kaper JB, Finlay BB. 1997. Interactions between enteropathogenic
392 *Escherichia coli* and host epithelial cells. *Trends Microbiol* 5: 109–114, PMID:
393 9080609, [https://doi.org/10.1016/S0966-842X\(97\)01000-7](https://doi.org/10.1016/S0966-842X(97)01000-7).
- 394 ECDC (European Center for Disease Prevention and Control). 2018. Annual epidemiological
395 report for 2016 Shiga-toxin/verocytotoxin-producing *Escherichia coli* (STEC/VTEC)
396 infection. Stockholm: European Center for Disease Prevention and Control

- 397 https://www.ecdc.europa.eu/sites/default/files/documents/AER_for_2016-STEC-
398 [VTEC_0.pdf](https://www.ecdc.europa.eu/sites/default/files/documents/AER_for_2016-STEC-VTEC_0.pdf) [accessed 6 April 2020].
- 399 EPA (Environmental Protection Agency). 2002. Method 1603: Escherichia coli (E. coli) in
400 water by membrane filtration using modified membrane-thermotolerant Escherichia coli
401 agar (modified mTEC). Washington, D.C.: Environmental Protection Agency
402 https://www.epa.gov/sites/production/files/2015-08/documents/method_1603_2009.pdf
403 [accessed 6 April 2020].
- 404 ESR (Institute of Environmental Science and Research). 2019. *Surveillance report: notifiable*
405 *diseases in New Zealand 2017*. Porirua: Institute of Environmental Science and
406 Research.
407 https://surv.esr.cri.nz/PDF_surveillance/AnnualRpt/AnnualSurv/2017/2017AnnualNDR
408 [eport_FINAL.pdf](https://surv.esr.cri.nz/PDF_surveillance/AnnualRpt/AnnualSurv/2017/2017AnnualNDR). [accessed 6 April 2020].
- 409 Franz E, Klerks MM, De Vos OJ, Termorshuizen AJ, van Bruggen AHC. 2007. Prevalence
410 of Shiga toxin-producing Escherichia coli stx1, stx2, eaeA, and rfbE genes and survival
411 of E. coli O157: H7 in manure from organic and low-input conventional dairy farms.
412 *Appl Environ Microbiol* 73: 2180–2190, PMID: 17277204,
413 <https://doi.org/10.1128/AEM.01950-06>.
- 414 Garzio-Hadzick A, Shelton DR, Hill RL, Pachepsky YA, Guber AK, Rowland R. 2010.
415 Survival of manure-borne E. coli in streambed sediment: effects of temperature and
416 sediment properties. *Water Res* 44: 2753–2762, PMID: 20219232,
417 <https://doi.org/10.1016/j.watres.2010.02.011>.
- 418 Gerba CP, McLeod JS. 1976. Effect of sediments on the survival of Escherichia coli in
419 marine waters. *Appl Environ Microbiol* 32: 114–120, PMID: 788634.
- 420 Germinario C, Caprioli A, Giordano M, Chironna M, Gallone MS, Tafuri S, et al. 2016.
421 Community-wide outbreak of haemolytic uraemic syndrome associated with Shiga toxin

- 422 2-producing *Escherichia coli* O26: H11 in southern Italy, summer 2013.
423 *Eurosurveillance* 21, PMID: 27684204, [https://doi.org/10.2807/1560-](https://doi.org/10.2807/1560-7917.ES.2016.21.38.30343)
424 [7917.ES.2016.21.38.30343](https://doi.org/10.2807/1560-7917.ES.2016.21.38.30343).
- 425 Gill A, Carrillo C, Hadley M, Kenwell R, Chui L. 2019. Bacteriological analysis of wheat
426 flour associated with an outbreak of Shiga toxin-producing *Escherichia coli* O121. *Food*
427 *Microbiol* 82: 474–481, PMID: 31027808, <https://doi.org/10.1016/j.fm.2019.03.023>.
- 428 Gluckman P. 2017. *New Zealand's fresh waters □: values, state, trends and human impacts*.
429 Wellington: Office of the Prime Minister's Chief Science Advisor.
430 <https://www.pmcsa.org.nz/wp-content/uploads/PMCSA-Freshwater-Report.pdf>
431 [accessed 6 April 2020].
- 432 Hickey CW, Clements WH. 1998. Effects of heavy metals on benthic macroinvertebrate
433 communities in New Zealand streams. *Environ Toxicol Chem* 17: 2338–2346,
434 <https://doi.org/10.1002/etc.5620171126>
- 435 Hickey M, Doran B. 2004. A review of the efficiency of buffer strips for the maintenance and
436 enhancement of riparian ecosystems. *Water Qual Res J Canada* 39: 311–317,
437 <https://doi.org/10.2166/wqrj.2004.042>.
- 438 Irshad H, Cookson AL, Ross CM, Jaros P, Prattley DJ, Donnison A, et al. 2016. Diversity
439 and relatedness of Shiga toxin-producing *Escherichia coli* and *Campylobacter jejuni*
440 between farms in a dairy catchment. *Epidemiol Infect* 144: 1406–1417, PMID:
441 26593403, <https://doi.org/10.1017/S0950268815002782>.
- 442 Ishii Y, Kimura S, Alba J, Shiroto K, Otsuka M, Hashizume N, et al. 2005. Extended-
443 spectrum β -lactamase-producing shiga toxin gene (stx1)-positive *Escherichia coli* O26:
444 H11: a new concern. *J Clin Microbiol* 43: 1072–1075, PMID:15750063,
445 <https://doi.org/10.1128/JCM.43.3.1072-1075.2005>.
- 446 Jaros P, Cookson AL, Campbell DM, Besser TE, Shringi S, Mackereth GF, et al. 2013. A

- 447 prospective case–control and molecular epidemiological study of human cases of Shiga
448 toxin-producing *Escherichia coli* in New Zealand. *BMC Infect Dis* 13: 450, PMID:
449 24079470, <https://doi.org/10.1186/1471-2334-13-450>.
- 450 Joy M. 2015. *Polluted inheritance: New Zealand's freshwater crisis*. Wellington, New
451 Zealand: BWB Texts.
- 452 Julian JP, de Beurs KM, Owsley B, Davies-Colley RJ, Ausseil AGE. 2017. River water
453 quality changes in New Zealand over 26 years: response to land use intensity. *Hydrol*
454 *Earth Syst Sci* 21:1149–1171; doi:10.5194/hess-21-1149-2017.
- 455 Karmali MA. 2018. Factors in the emergence of serious human infections associated with
456 highly pathogenic strains of shiga toxin-producing *Escherichia coli*. *Int J Med Microbiol*
457 308: 1067–1072, PMID: 30146439, <https://doi.org/10.1016/j.ijmm.2018.08.005>.
- 458 Kerangart S, Cournoyer B, Loukiadis E. 2018. C-source metabolic profilings of foodborne
459 Shiga-toxin producing *E. coli* match serogroup differentiations and highlight functional
460 adaptations. *Int J Food Microbiol* 266: 324–336, PMID: 29217302,
461 <https://doi.org/10.1016/j.ijfoodmicro.2017.10.018>.
- 462 King LA, Nogareda F, Weill F-X, Mariani-Kurkdjian P, Loukiadis E, Gault G, et al. 2012.
463 Outbreak of Shiga toxin–producing *Escherichia coli* O104: H4 associated with organic
464 fenugreek sprouts, France, June 2011. *Clin Infect Dis* 54: 1588–1594, PMID: 22460976,
465 <https://doi.org/10.1093/cid/cis255>.
- 466 Lalzampaia H, Dutta TK, Warjri I, Chandra R. 2013. PCR-based detection of extended-
467 spectrum β -lactamases (*bla* CTX-M-1 and *bla* TEM) in *Escherichia coli*, *Salmonella*
468 spp. and *Klebsiella pneumoniae* isolated from pigs in North Eastern India (Mizoram).
469 *Indian J Microbiol* 53: 291–296, PMID: 24426125, [https://doi.org/10.1007/s12088-013-](https://doi.org/10.1007/s12088-013-0378-z)
470 0378-z.
- 471 Luna-Gierke RE, Griffin PM, Gould LH, Herman K, Bopp CA, Strockbine N, et al. 2014.

- 472 Outbreaks of non-O157 Shiga toxin-producing *Escherichia coli* infection: USA.
473 *Epidemiol Infect* 142: 2270–2280, PMID: 24398154,
474 <https://doi.org/10.1017/S0950268813003233>.
- 475 Mauro SA, Opalko H, Lindsay K, Colon MP, Koudelka GB. 2013. The microcosm mediates
476 the persistence of shiga toxin-producing *Escherichia coli* in freshwater ecosystems. *Appl*
477 *Environ Microbiol* 79: 4821–4828, PMID: 23747699,
478 <https://doi.org/10.1128/AEM.01281-13>.
- 479 McBride G, Till D, Ryan T, Ball A, Lewis G, Palmer S, et al. 2002. *Freshwater microbiology*
480 *research programme report: pathogen occurrence and human health risk assessment*
481 *analysis*. Wellington, Ministry of Health. Accessed on April 6, 2020, at
482 [http://www.mfe.govt.nz/publications/water/freshwater-microbiology-nov02/freshwater-](http://www.mfe.govt.nz/publications/water/freshwater-microbiology-nov02/freshwater-microbiology-nov02.pdf)
483 [microbiology-nov02.pdf](http://www.mfe.govt.nz/publications/water/freshwater-microbiology-nov02/freshwater-microbiology-nov02.pdf) [accessed 6 April 2020].
- 484 Merritt DM, Scott ML, Poff NL, Auble GT, Lytle DA. 2010. Theory, methods and tools for
485 determining environmental flows for riparian vegetation: riparian vegetation-flow
486 response guilds. *Freshw Biol* 55:206–225, doi:10.1111/j.1365-2427.2009.02206.x.
- 487 MfE and Stats New Zealand (Ministry for the Environment and Statistics New Zealand).
488 2017. *New Zealand's Environmental Reporting Series: Our fresh water 2017*.
489 Wellington: Ministry for the Environment and Statistics New Zealand.
490 [https://www.mfe.govt.nz/sites/default/files/media/Environmental%20reporting/our-](https://www.mfe.govt.nz/sites/default/files/media/Environmental%20reporting/our-fresh-water-2017_1.pdf)
491 [fresh-water-2017_1.pdf](https://www.mfe.govt.nz/sites/default/files/media/Environmental%20reporting/our-fresh-water-2017_1.pdf) [accessed 6 April 2020].
- 492 Muirhead RW, Davies-Colley RJ, Donnison AM, Nagels JW. 2004. Faecal bacteria yields in
493 artificial flood events: quantifying in-stream stores. *Water Res* 38: 1215–1224, PMID:
494 14975655, <https://doi.org/10.1016/j.watres.2003.12.010>.
- 495 OECD (The Organisation for Economic Co-operation and Development). 2017. *Diffuse*
496 *pollution, degraded waters: emerging policy solutions*.

- 497 <https://dx.doi.org/10.1787/9789264269064-en> [accessed 6 April 2020].
- 498 Oporto B, Ocejo M, Alkorta M, Marimón JM, Montes M, Hurtado A. 2019. Zoonotic
499 approach to Shiga toxin-producing *Escherichia coli*: integrated analysis of virulence and
500 antimicrobial resistance in ruminants and humans. *Epidemiol Infect* 147, PMID:
501 32077399, <https://doi.org/10.1017/S0950268820000357>.
- 502 Palmateer GA, McLean DE, Kutas WL, Meissner SM. 1993. Suspended particulate-bacterial
503 interaction in agricultural drains. In: *Particulate matter aquatic contaminants*, 1-40. Rao
504 SS, ed. Boca Raton, USA: CRC Press.
- 505 Paton AW, Paton JC. 1998. Detection and characterization of Shiga toxigenic *Escherichia*
506 *coli* by using multiplex PCR assays for *stx1*, *stx2*, *eaeA*, enterohemorrhagic *E. coli*
507 *hlyA*, *rfb O111*, and *rfb O157*. *J Clin Microbiol* 36: 598–602, PMID: 14632994,
508 <https://doi.org/10.1046/j.1365-2672.2003.02091.x>.
- 509 Perkins TL, Clements K, Baas JH, Jago CF, Jones DL, Malham SK, et al. 2014. Sediment
510 composition influences spatial variation in the abundance of human pathogen indicator
511 bacteria within an estuarine environment. *PLoS One* 9: e112951, PMID: 25397595,
512 <https://doi.org/10.1371/journal.pone.0112951>.
- 513 Perkins TL, Perrow K, Rajko-Nenow P, Jago CF, Jones DL, Malham SK, et al. 2016. Decay
514 rates of faecal indicator bacteria from sewage and ovine faeces in brackish and
515 freshwater microcosms with contrasting suspended particulate matter concentrations. *Sci*
516 *Total Environ* 572: 1645–1652, PMID: 27039273,
517 <https://doi.org/10.1016/j.scitotenv.2016.03.076>.
- 518 Perna NT, Mayhew GF, Pósfai G, Elliott S, Donnenberg MS, Kaper JB, et al. 1998.
519 Molecular evolution of a pathogenicity island from enterohemorrhagic *Escherichia coli*
520 O157: H7. *Infect Immun* 66: 3810–3817, PMID: 9673266,
521 <https://dx.doi.org/10.1128/IAI.66.8.3810-3817.1998>

- 522 Possé B, De Zutter L, Heyndrickx M, Herman L. 2007. Metabolic and genetic profiling of
523 clinical O157 and non-O157 Shiga-toxin-producing *Escherichia coli*. *Res Microbiol*
524 158: 591–599, PMID: 17845842, <https://doi.org/10.1016/j.resmic.2007.06.001>.
- 525 Power AL, Tennant RK, Jones RT, Tang Y, Du J, Worsley AT, et al. 2018. Monitoring
526 impacts of urbanisation and industrialisation on air quality in the Anthropocene using
527 urban pond sediments. *Front Earth Sci* 131, <https://doi.org/10.3389/feart.2018.00131>.
- 528 Prieto MD, Lopez B, Juanes JA, Revilla JA, Llorca J, Delgado-Rodriguez M. 2001.
529 Recreation in coastal waters: health risks associated with bathing in sea water. *J*
530 *Epidemiol Community Heal* 55: 442–447, PMID: 11351003,
531 <https://doi.org/10.1136/jech.55.6.442>.
- 532 Ratha KC. 2019. Growing industrialisation and its cumulative impacts: insights from the
533 Mahanadi River basin. *Soc Change* 49: 519–530,
534 <https://doi.org/10.1177%2F0049085719863904>.
- 535 Reddy KR, Khaleel R, Overcash MR. 1981. Behavior and transport of microbial pathogens
536 and indicator organisms in soils treated with organic wastes. *J Environ Qual* 10: 255–
537 266, <https://doi.org/10.2134/jeq1981.00472425001000030001x>
- 538 Rose JB, Epstein PR, Lipp EK, Sherman BH, Bernard SM, Patz JA. 2001. Climate variability
539 and change in the United States: potential impacts on water-and foodborne diseases
540 caused by microbiologic agents. *Environ Health Perspect* 109: 211–221, PMID:
541 11359688, <https://doi.org/10.1289/ehp.01109s2211>.
- 542 Senthakumaran T, Brandal LT, Lindstedt B-A, Jørgensen SB, Charnock C, Tunsjø HS. 2018.
543 Implications of stx loss for clinical diagnostics of Shiga toxin-producing *Escherichia*
544 *coli*. *Eur J Clin Microbiol Infect Dis* 37: 2361–2370, PMID: 30267169,
545 <https://doi.org/10.1007/s10096-018-3384-6>.
- 546 Severi E, Vial F, Peron E, Mardh O, Niskanen T, Takkinen J. 2016. Community-wide

547 outbreaks of haemolytic uraemic syndrome associated with Shiga toxin-producing
548 *Escherichia coli* O26 in Italy and Romania: a new challenge for the European Union.
549 *Eurosurveillance* 21, PMID: 27983509, [https://doi.org/10.2807/1560-](https://doi.org/10.2807/1560-7917.ES.2016.21.49.30420)
550 [7917.ES.2016.21.49.30420](https://doi.org/10.2807/1560-7917.ES.2016.21.49.30420).

551 Shaw A, Helterbran K, Evans MR, Currey C. 2016. Growth of *Escherichia coli* O157: H7,
552 non-O157 Shiga toxin-producing *Escherichia coli*, and *Salmonella* in water and
553 hydroponic fertilizer solutions. *J Food Prot* 79: 2179–2183, PMID: 28221949,
554 <https://doi.org/10.4315/0362-028X.JFP-16-073>.

555 Solomon EB, Yaron S, Matthews KR. 2002. Transmission of *Escherichia coli* O157: H7 from
556 contaminated manure and irrigation water to lettuce plant tissue and its subsequent
557 internalization. *Appl Environ Microbiol* 68: 397–400, PMID: 11772650,
558 <https://doi.org/10.1128/aem.68.1.397-400.2002>.

559 Swaggerty CL, Corcionivoschi N, Ricke SC, Callaway TR. 2018. The first 30 years of Shiga
560 toxin-producing *Escherichia coli* in cattle production: incidence, preharvest ecology,
561 and management. In: *Food and feed safety systems and analysis*, 117–131, Ricke S,
562 Atungulu G, Rainwater C, Park S, eds. London, UK: Academic Press.

563 Sweeney BW, Newbold JD. 2014. Streamside forest buffer width needed to protect stream
564 water quality, habitat, and organisms: a literature review. *J Am Water Resour Assoc*
565 50:560–584, <https://doi.org/10.1111/jawr.12203>.

566 Tarr GAM, Stokowski T, Shringi S, Tarr PI, Freedman SB, Oltean HN, et al. 2019.
567 Contribution and interaction of Shiga toxin genes to *Escherichia coli* O157: H7
568 virulence. *Toxins (Basel)* 11: 607, PMID: 31635282,
569 <https://doi.org/10.3390/toxins11100607>.

570 Valat C, Haenni M, Saras E, Auvray F, Forest K, Oswald E, et al. 2012. CTX-M-15
571 extended-spectrum β -lactamase in a Shiga toxin-producing *Escherichia coli* isolate of

- 572 serotype O111: H8. *Appl Environ Microbiol* 78: 1308–1309, PMID: 22156432,
573 <https://doi.org/10.1128/AEM.06997-11>.
- 574 Valilis E, Ramsey A, Sidiq S, DuPont HL. 2018. Non-O157 Shiga toxin-producing
575 *Escherichia coli*—a poorly appreciated enteric pathogen: systematic review. *Int J Infect*
576 *Dis* 76: 82–87, PMID: 30223088, <https://doi.org/10.1016/j.ijid.2018.09.002>.
- 577 Waters TF. 1995. Sediment in streams: sources, biological effects, and control. American
578 Fisheries Society Monograph 7, Bethesda, USA: American Fisheries Society.
- 579 Weiskerger CJ, Whitman RL. 2018. Monitoring *E. coli* in a changing beachscape. *Sci Total*
580 *Environ* 619: 1236–1246, PMID: 29734602,
581 <https://doi.org/10.1016/j.scitotenv.2017.11.167>.
- 582 Werber D, Fruth A, Liesegang A, Littmann M, Buchholz U, Prager R, et al. 2002. A
583 multistate outbreak of Shiga toxin–producing *Escherichia coli* O26: H11 infections in
584 Germany, detected by molecular subtyping surveillance. *J Infect Dis* 186: 419–422,
585 PMID: 12134240, <https://doi.org/10.1086/341457>.
- 586 Wilkinson J, Jenkins A, Wyer M, Kay D. 1995. Modelling faecal coliform dynamics in
587 streams and rivers. *Water Res* 29: 847–855, PMID: 17037150,
588 <https://doi.org/10.2166/wst.2006.466>.
- 589 WHO (World Health Organization). 2019. Global health estimates 2016: deaths by cause,
590 age, sex, by country and by region, 2000-2016. Geneva, Switzerland.
- 591 Yan Z, Han W, Peñuelas J, Sardans J, Elser JJ, Du E, et al. 2016. Phosphorus accumulates
592 faster than nitrogen globally in freshwater ecosystems under anthropogenic impacts.
593 *Ecol Lett* 19: 1237–46, PMID: 27501082, <https://doi.org/10.1111/ele.12658>.
- 594 Yang H, Rehman MU, Zhang S, Yang J, Li Y, Gao J, et al. 2019. High prevalence of CTX-M
595 belonging to ST410 and ST889 among ESBL producing *E. coli* isolates from waterfowl
596 birds in China’s tropical island, Hainan. *Acta Trop* 194: 30–35, PMID: 30853356,

597 <https://doi.org/10.1016/j.actatropica.2019.03.008>.

598 Zou H, Zheng B, Sun M, Ottoson J, Li Y, Berglund B, et al. 2019. Evaluating dissemination

599 mechanisms of antibiotic-resistant bacteria in rural environments in China by using

600 CTX-M-producing *Escherichia coli* as an indicator. *Microb Drug Resist* 25: 975-84,

601 PMID: 30942653, <https://doi.org/10.1089/mdr.2018.0431>.

602

Table 1. *Escherichia coli* colony counts, in CFU/100 ml, for water and sediment samples collected from the Selwyn, Rangitata, and Ashley rivers during May and September 2018.

Sampling time	Sampling sites and substrates											
	Above intensive dairy						Below intensive dairy					
	Ashley water	Ashley sediment	Rangitata water	Rangitata sediment	Selwyn water	Selwyn sediment	Ashley water	Ashley sediment	Rangitata water	Rangitata sediment	Selwyn water	Selwyn sediment
May (autumn)	45	230	20	7,100	10	200	440	730	175	273,300	160	230
September (spring)	20	400	20	5,000	505	200	2,000	3,600	40	1,600	2,250	12,200

Table 2. Presence of *Escherichia coli* control (*uidA*), STEC virulence (*stx1*, *stx2*, *eae*), serogroup (O26 *wzy* and O157 *rfbE*), and antibiotic resistance (*bla* CTX-M) genes in water and sediment samples collected from the Selwyn, Rangitata, and Ashley rivers during May and September 2018.

Gene locus	Sampling sites and substrates											
	Above intensive dairy						Below intensive dairy					
	Ashley water	Ashley sediment	Rangitata water	Rangitata sediment	Selwyn water	Selwyn sediment	Ashley water	Ashley sediment	Rangitata water	Rangitata sediment	Selwyn water	Selwyn sediment
May (autumn)												
<i>uidA</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>stx1</i>	-	-	-	-	-	-	+	+	+	-	-	-
<i>stx2</i>	-	-	-	-	-	-	-	+	+	-	-	-
<i>eae</i>	-	-	-	-	-	-	+	+	+	-	-	-
<i>wzy</i> O26	-	-	-	-	-	-	+	+	+	-	-	-
<i>rfbE</i> O157	-	-	-	-	-	+	+	+	+	-	-	+

<i>bla</i> CTX-M	-	-	-	-	-	-	-	-	-	-	-	-	-
September (spring)													
<i>uidA</i>	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>stx1</i>	-	+	-	+	-	-	-	-	+	-	+	+	+
<i>stx2</i>	-	+	-	+	-	+	+	-	-	+	-	+	+
<i>eae</i>	-	+	+	+	-	+	+	+	+	+	+	+	+
<i>wzy</i> 026	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>rfbE</i> O157	-	-	-	-	-	+	-	-	-	+	+	+	+
<i>bla</i> CTX-M	-	-	-	+	-	-	-	-	+	-	+	+	+

604 + Gene detected

605 - Gene not detected

606

607 **Figure Legends**

608

609 **Figure 1.** Map of central and southern Canterbury, New Zealand with the Ashley, Rangitata,
610 and Selwyn rivers labeled. Locations at which water and sediment were collected during May
611 and September 2018 are marked with circles; for each river the white centered dot indicates
612 the above intensive dairy site and the black centered dot the below intensive dairy site.

