Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

THE EFFECT OF LIPID ON THE DIGESTION OF CELLULOSE BY THE RUMINANT.

A thesis presented in partial fulfilment of the requirements for the degree of Master of Agricultural Science in Animal Science at Massey University.

ROBIN DAVID KIRK

1969

ACKNOWLEDGEMENTS

I wish to thank the following who gave valuable assistance in various ways:

Dr. J.C. Hawke for the guidance and helpful advice given throughout the course of this work.

Professor D.S. Flux and Mr. G.F. Wilson for their valuable discussion and criticism of this work.

The Chemistry and Biochemistry Department for the provision of apparatus and facilities during 1969.

Mr. N.A. Thomson and Miss J.M. Aldersley for technical assistance; and the staff of No. 2 Dairy Unit for the care of the fistulated cattle.

Mrs. O.J. England for typing the final copies of the thesis.

Special thanks are accorded my wife, Rosemary, for giving the encouragement and financial supportance necessary.

TABLE OF CONTENTS

	Page
REVIEW OF LITERATURE	
1. Metabolism of Lipids by Ruminants	1
2. Interconversion of Individual Volatile Fatty Acids	13
3. Propionate Metabolism	13
4. Cellulose Digestion	14
5. Techniques of Sampling from the Rumen	17
6. *In-vitro* Experimentation	19
OBJECT OF THE PRESENT WORK	24
MATERIALS AND METHODS	
Treatment of Animals	26
Clarified Rumen Liquor (Dialysate)	26
Grass Fibre	27
Lipid Substrates	27
Artificial Saliva	27
Radioactive Materials	28
Impermeable Apparatus	28
Semi-permeable Apparatus	29
Analytical Methods	
Gas Production	31
Ammonia Determination	31
Isolation and Determination of the VFA's in Rumen Fluid	32
Individual Volatile Fatty Acids	33
Gas-Liquid Chromatography using the Titrimetric Method (James & Martin)	34
Varian Aerographs	35
Separation of ^{14}C-VFA's using the Titrimetric Method (James & Martin)	36
Separation of ^{14}C-VFA's using a Varian Aerograph 1520	37
Measurement of Radioactivity	38
pH Readings	38
Statistical Methods	38

Page

EXPERIMENTAL AND RESULTS 40

1. Preliminary Experiments

1.1. Gas Production 40

1.2 Ammonia Production 42

1.3 Total and individual VFA production 42

2. Effect of different amounts of lipid substrate on total and individual VFA production and ammonia production from grass fibre incubated in rumen liquor for 24, 36, 48, or 72hr. 45

2.1 Total VFA production 46

2.2 Acetate production 47

2.3 Propionate production 48

2.4 Butyrate production 49

2.5 Ammonia production 51

2.6 pH determination of rumen liquor incubations containing grass fibre and linseed oil within impermeable apparatus 52

3. The effect of various lipids on volatile fatty acid production. 56

3.1 Total and individual VFA production from rumen liquor containing grass fibre. 56

3.2 Comparison of the effect of lipid upon the production of total and individual VFA's of rumen liquor incubated with or without grass fibre. 59

4. The use of ^{14}C-cellulose in rumen fermentations of lipid and grass fibre within impermeable apparatus. 61

4.1 Total and individual VFA production 61

4.2 Incorporation of ^{14}C into VFA and CO_2 62

4.3 Distribution of ^{14}C among individual VFA's 62

4.4 Net production of total VFA, acetic, propionic and butyric acids. 64

4.5 Incorporation of ^{14}C into VFA and CO_2 65

4.6 Distribution of ^{14}C among individual VFA's 65

Page

5. The incubation of (2-^{14}C)glycerol (0.45' μc./ml. in rumen liquor with grass fibre and cod liver oil. 68

5.1 Net production of total VFA, acetic, propionic, and butyric acids. 68

5.2 The percentage incorporation of ^{14}C from (2-^{14}C)glycerol into total and individual VFA's. 70

6. In-vitro experiments using the micro-artificial rumen apparatus. 72

6.1 Experiments a and b 72

6.2 Experiments c and d 75

7. Use of ^{14}C-cellulose in rumen fermentations of lipid and grass fibre, within the micro-artificial rumens. 83

7.1 Total VFA production 83

7.2 Total and individual ^{14}C-VFA production 84

7.3 Investigations into the adsorption and carryover of VFA's on gas-liquid chromatography columns. 86

DISCUSSION 89

Effect of Lipid on the Production of Total and Individual VFA's. 89

Use of Micro-Artificial Rumens. 91

Comparison of the Impermeable Apparatus and the Micro-Artificial Rumens. 92

Use of ^{14}C-cellulose in the Impermeable Apparatus. 93

Use of ^{14}C-cellulose in Micro-Artificial Rumens. 94

Effect of Different Lipids on VFA Production. 96

Effect of Lipid on Ammonia Concentrations of Rumen Liquor. 98

Effect of Lipid on the pH of Rumen Liquor. 98

Use of (2-^{14}C)glycerol. 99

Conclusion. 101

SUMMARY 104

REFERENCES 107

APPENDICES

LIST OF FIGURES

Figure		Page
1	The Micro-Artificial Rumen. (Dawson, Ward & Scott, 1964).	30
2	The effect of linseed oil(2.0g.) on individual VFA concentrations (m-mole/l.) when 0.5g. of grass fibre was incubated in rumen liquor (50ml.) for 48hr.	43
3	The effect of linseed oil(0, 1, 2 & 3g./100ml. rumen liquor) on total and individual VFA concentrations (m-mole/l.) when 0.5g. of grass fibre was incubated in rumen liquor (50ml.) for 24, 36, 48 or 72hr.	50
4	The effect of time of incubation on the ammonia concentration (mg.NH_3/100ml.) of rumen liquor (50ml.) containing 0.5g. of grass fibre and linseed oil (0, 1, 2 & 3g./100ml.)	53
5	The effect of linseed oil(0, 1, 2 & 3g./100ml. rumen liquor) on ammonia concentrations (mg.NH_3/100ml.) when 0.5g. of grass fibre was incubated in rumen liquor (50ml.) for 24, 36, 48 or 72hr.	54
6	Total VFA production(m-mole/l.) from rumen liquor(50ml.) incubated with (a) 0.5g. of grass fibre + 1.5g. of cod liver oil; (b) 0.5g. of grass fibre; (c) or without additional substrates, for 24hr. within the semi-permeable apparatus. (Experiment c.).	77
7	Total VFA production(m-mole/l.) from rumen liquor(50ml.) incubated with (a) 0.5g. of grass fibre + 1.5g. of cod liver oil; (b) 0.5g. of grass fibre; (c) or without additional substrates, for 24hr. within the semi-permeable apparatus. (Experiment d.).	78

Figure Page

8 Rate of total VFA production(m-mole/l./2hr.) within the semi-permeable apparatus containing 50ml. of rumen liquor incubated for 24hr. with (a) 0.5g. grass fibre + 1.5g. of cod liver oil; (b) 0.5g. of grass fibre; or (c) without additional substrates. 80

9 Rate of production of individual VFA's(m-mole/l./2hr.) within the semi-permeable apparatus containing 50ml. of rumen liquor incubated for 24hr. with (a) 0.5g. of grass fibre + 1.5g. of cod liver oil; (b) 0.5g. of grass fibre; or (c) without additional substrates. 81

10 The production of ^{14}C-VFA from ^{14}C-cellulose incubated with rumen liquor in the presence of (a) 0.5g. of grass fibre; (b) 0.5g. of grass fibre + 1.0g. of cod liver oil; or (c) 0.5g. of grass fibre + 3.0g. of cod liver oil for 24hr. within the semi-permeable apparatus. 85