

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**Can mating ewes on *Lotus
corniculatus* be used to reduce
lamb mortality between birth and
weaning?**

**A Thesis Presented in Fulfillment of the Requirements for the
Degree of Master in Animal Science at Massey University**

Victor Hugo Rojas Maldonado

2005

Acknowledgments

All my gratitude to my chief supervisor Professor Tom N. Barry, for all his support, guidance and kindness even before coming to New Zealand, and mainly for being more than a supervisor, a good friend. To my family: my Father Victor M. Rojas, my two sisters Gabriela and Monica, my niece Linda and my brother in law Francisco Juarez for all their love and support at all times and for being the root of my essence, I love you all.

Also to all my friends for cheering me and supporting me every time I needed, mainly those that I had the fortune to meet here, because more than my friends they became my family in this country: Alejandra, Natalia, Klelia, Lisa, Carolina, Andres, Dipti, Fernanda, Nacho, Himansu, Sumit, and Gonzalo. Thanks guys for all those great moments and for being there every single time I felt down.

To my co-supervisors: Nicolas Lopez-Villalobos and Timothy J. Parkinson for their time (which is so valuable), patience and guidance. To Geoff Purchas for his help and patience during the time that I was developing my experiment at Riverside Farm, but mainly for his good head with that extraordinary ability to organize things; thank you so much Geoff, without you I know everything would have been so difficult and chaotic at the farm. Also to Nathan Crombie for his help during the development of the experiment, for making the long working hours at the lab and the paddocks go faster with all those interesting, funny and weird stories, and for being such a talkative and laughing person. To Colin Morgan for his help at the farm and for sharing his experience and knowledge regards animal management.

To all the people from the 'international student support office' at Massey University, especially to Silvia Hooker and Sue Flynn; what can I say? Both of you are simply amazing and marvelous, thanks for everything.

Finally, to NZAID for giving me such a great opportunity to come to this beautiful country to study my Masters.

Table of contents

ACKNOWLEDGMENTS	I
TABLE OF CONTENTS	II
FIGURES AND TABLES.....	IV
CHAPTER 1.....	1
ABSTRACT	1
LITERATURE REVIEW	3
1.1 INTRODUCTION.....	3
1.2 SHEEP PRODUCTION SYSTEMS IN NEW ZEALAND.....	4
1.2.1 <i>The annual cycle</i>	4
1.2.2 <i>Grazing systems</i>	8
1.2.3 <i>Grazing forages</i>	10
1.3 LOTUS CORNICULATUS	11
1.3.1 <i>Nutritional characteristics</i>	11
1.3.2 <i>Lotus corniculatus as a grazing forage</i>	14
1.4 CONDENSED TANNINS.....	18
1.4.1 <i>Chemical characteristics</i>	18
1.4.2 <i>Concentration in plants</i>	22
1.4.3 <i>Effects on ruminal digestion</i>	25
1.4.4 <i>Effects on animal production</i>	29
1.5 CONCLUSIONS	38

CHAPTER 2.....	40
CAN MATING EWES ON <i>LOTUS CORNICULATUS</i> BE USED TO REDUCE LAMB MORTALITY BETWEEN BIRTH AND WEANING?.....	40
2.1 INTRODUCTION.....	40
2.2 MATERIAL AND METHODS.....	41
2.2.1 <i>Experimental design</i>	41
2.2.2. <i>Forage</i>	42
<i>Sheep grazing Lotus corniculatus in the sward.</i>	42
2.2.3 <i>Grazing Management</i>	43
2.2.4 <i>Animals</i>	43
2.2.5 <i>Laboratory analyses</i>	45
2.2.6 <i>Statistical analyses</i>	45
2.3 RESULTS	46
2.3.1 <i>Forages</i>	46
2.3.2 <i>Animals</i>	50
2.3.3 <i>Wool production</i>	57
2.4 DISCUSSION	58
2.5 CONCLUSIONS	65
2.6 REFERENCES.....	66

Figures and Tables

- Figure 1.1. Metabolisable energy (ME) concentration (MJ/Kg DM) of diet selected by lambs grazing *Lotus corniculatus* and perennial ryegrass/white clover pasture over the summer season of 2002-2003 in a New Zealand dry land pastoral system. 12
- Figure 1.2. Pregrazing dead matter in areas of *Lotus corniculatus* and perennial ryegrass/white clover pasture grazed by lambs over the spring/summer season of 2002 and 2003 in a commercial dryland system on the east coast of New Zealand..15
- Figure 1.3. Constituents of hydrolysable tannins and the structure of tannic acid. 18
- Figure 1.4. Thiolysis reaction of condensed tannins polymers with procyanidin (R=H) and/or prodelfinidin (R=OH) units. 20
- Figure 1.5. The relationship between condensed tannin concentration in the dry matter of forage species, and (a) the ratio of non-amonia-nitrogen (NAN) flowing at the abomasum or duodenum, and (b) microbial N flow at the abomasum or duodenum per unit of N eaten by sheep 26
- Figure 1.6. The effect of forage CT concentration on wool production of grazing sheep, calculated as the wool production response of CT-acting sheep relative to PEG-supplementation (CT-inactivated) or non-CT-containing pasture sheep..... 29
- Figure 1.7. Effect of intake of digestible energy (MJ/day) and digestible protein (g/day) on the percentage of ewes with multiple ovulations..... 35

Figure 2.1. Change in mean ewe live weight per treatment during the whole experimental period.	52
Figure 2.2. Change in mean ewe body condition score per treatment during the whole experimental period.	52
Figure 2.3. Rainfall per month (January to December) in three consecutive years at Riverside farm.	60
Figure 2.4a. Lamb mortality per treatment during lambing weeks.....	61
Figure 2.4b. Lambs born per week, in each treatment group, during lambing weeks. . . .	62
Figure 2.4c. Number of ewes lambing per week, in each treatment group, during lambing weeks.....	62
Figure 2.5a. Lamb mortality per birth rank during lambing weeks in the lotus group. . . .	63
Figure 2.5b. Lamb mortality per birth rank during lambing weeks in the control pasture group.....	63

Table 1.1. Daily ME and DM intake requirements for ewes at different physiological stages.....	7
Table 1.2. Mean values of metabolizable energy concentration (ME; MJ/Kg DM), total nitrogen (N; g/Kg DM) and condensed tannin concentration (CT; g/Kg DM) of the diet selected by sheep and deer grazing perennial ryegrass/white clover (<i>Lolium perenne</i> / <i>Trifolium repens</i>) dominant pasture, <i>Lotus corniculatus</i> (birds foot trefoil; cv. Grasslands Goldide), or lucerne (<i>Medicago sativa</i>) ¹ .	13
Table 1.3. Annual and seasonal dry matter production (t DM/ha) of perennial ryegrass/white clover (<i>Lolium perenne</i> / <i>Trifolium repens</i>) pasture or <i>Lotus corniculatus</i> (cv. Grasslands Goldie) averaged over three consecutive years in a commercial dryland farming system on the East Coast in New Zealand.	17
Table 1.4. Plant materials that contain tannins.	23
Table 1.5. The extractable and bound condensed tannin content of some legumes, grasses and herbs fed to ruminants in temperate grazing systems, measured by the butanol-HCl method.	24
Table 1.6. Effect of condensed tannins (CT) upon the intake and absorption of essential amino acids (EAA) from the small intestine of sheep fed on fresh <i>L. corniculatus</i> and <i>L. pedunculatus</i> with or without an intraruminal infusion of polyethylene glycol (PEG) to inactivate the CT.	27
Table 1.7. Rumen concentration of ammonia and plasma concentration of urea in ewes grazing perennial ryegrass/white clover pasture and <i>L. corniculatus</i> , with or without twice-daily oral administration of polyethylene glycol (PEG; MW 3 500).30	30

Table 1.8. Plasma concentration of amino acids (μM) in ewes grazing perennial ryegrass/white clover pasture and <i>Lotus corniculatus</i> (cv. Grasslands Goldie) <i>ad libitum</i> , with and without twice-daily oral administration of polyethylene glycol (PEG;MW 3500).	31
Table 1. 9. The effect of condensed tannins in <i>Lotus corniculatus</i> upon dry matter intake, milk yield, milk protein production and methane production in grazing dairy cows.	32
Table 1.10. Voluntary feed intake (VFI), live weight gain (LWG), carcass gain and wool growth in lambs (Experiment 1) and dry ewes (experiment 2) grazing the forage legumes <i>Lotus corniculatus</i> (27-34 g CT/Kg DM) and lucerne (0.3 g. total CT/Kg DM) during summer.	33
Table 1.11. Concentration of plasma amino acids measured in wethers at 4 levels of intake and 2 levels of protein content in diet, and ovulation rates in ewes fed the same diets.....	35
Table 1.12. The effect of grazing ewes during mating on <i>L. corniculatus</i> or perennial ryegrass/white clover (pasture) and of supplementation with polyethylene glycol (PEG) on ovulation rate (OR).	37
Table 1.13. The effect of mating on <i>Lotus corniculatus</i> in 2001 on the reproductive performance (%ewes mated) of ewes on Massey University dryland farm in the Wairarapa.	38
Table 2.1. Pre-grazing and post-grazing mass and dead matter content of forages grazed during the experiment.	48
Table 2.2. Chemical composition of the diet selected of forages grazed during the experiment (g/Kg DM)..	49

Table 2.3. Ewe live weight change (g/day) and body condition score (BCS) change during the 74-day of nutritional trial, and reproductive rate expressed as a percentage of the total number of ewes exposed to the ram	51
Table 2.4. Conception rate (at scanning), expressed as a percentage of the total number of ewes exposed to the ram, and fecundity as a percentage of lambing ewes	54
Table 2.5. Total lamb mortality per treatment and per birth rank (%) from birth to weaning (mean values with 95% confidence intervals), and mean lambing date per treatment and birth rank	55
Table 2.6. Lamb birth and weaning weights (Kg) per treatment, corrected for the small difference in the mean date of lambing	56
Table 2.7. Twelve-month wool production and staple length for each group	57
Table 2.8. Liveweight gain (LWG) in ewes grazing either control pasture or <i>L. corniculatus</i> during mating in different studies in New Zealand.....	58
Table 2.9. Nutritive value of pasture and <i>Lotus corniculatus</i> during mating at Riverside farm, and ewe lambing percentage.....	59