

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**Deterministic modelling of energy supply and demand
for foaling mares managed at pasture on commercial
Thoroughbred stud farms.**

A thesis presented in fulfilment of requirement for the degree,
Master of Science (Animal Science).

Massey University, Palmerston North, New Zealand.

Yin Ying Chin

2018

Abstract

There is an economic incentive to breed and foal mares early in the season within the New Zealand commercial Thoroughbred production industry. This has caused disparity between the natural and commercially imposed breeding season, and possibly, the period of peak pasture energy availability and the mare energy requirements. A deterministic model was developed to model the energy balance, energy intake and energy requirement of Thoroughbred mares managed at pasture under commercial conditions to assess their energy status. The response of energy intake and energy balance to changes in five variables were tested. The variables tested were dry matter intake, bodyweight, foaling date, pasture metabolisable energy, and energy requirement. For all foaling dates modelled, the mare was in energy surplus during pregnancy. Onset of lactation created a rapid and significant decrease in mare energy balance. Delay in foaling increased the magnitude of the post-partum energy deficit, and later foaling mares experienced a prolonged decrease in post-partum energy balance. The size of the deficit would, theoretically, decrease 1 body condition score, decrease circulating leptin concentration and initiate lean body mass mobilization which was previously found to negatively impact reproductive performance in the mares. The duration and size of the post-partum energy deficit could be reduced by shifting foaling date closer to the beginning of breeding season (1st September), thus synchronizing period of peak pasture energy with the mare's theoretical requirements. Energy balance and energy intake were more sensitive to pasture energy, dry matter intake and energy requirement changes with less influence from bodyweight to energy balance, suggesting pasture quality and quantity is an important influencer on nutritional status of the mare kept on pasture. Initiatives should be taken to monitor pasture quality and growth pattern, and optimize energy and dry matter intake in mares at pasture. These efforts could assist in maximizing pasture utilization, minimize energy deficit and achieve cost effective pastoral and nutritional management. When more pasture quality and growth data becomes available for equine stud farms, the model has the potential to achieve farm specific application on a wide range of equine body weights.

Acknowledgement

I would like to thank my supervisors Associate Professor Chris Rogers, Dr. Erica Gee and Dr. Penny Back for their guidance since my undergraduate studies. Their dedication and patience have allowed me to learn and work under a supportive environment. I am grateful for the opportunity to work on my topic of interest and with the teachers who inspired this endeavour. A special thanks to Dr. Javier Roca for going above and beyond in assisting with my thesis writing, constructing the nutritional model, data analysis, and providing helpful advice from time to time. Huge thanks to my family for their support, encouragement and acting as my reminder for being diligent and persistent in the face of challenges. This incredible journey would be impossible without them. Special thanks to Terry, Vicki and Cindy for sharing their knowledge and providing me with the opportunity to acquire more experience and understanding in the equine industry, which expanded my thinking and interpretation of my findings. This is dedicated to my good friend and role model, Adele: “Getting the most out of life is not about how much you keep for yourself, but how much you pour into others.”

Table of Contents

Abstract.....	i
Acknowledgement	ii
Table of Contents	iii
List of Tables.....	v
List of Figures	vii
Chapter 1: Introduction	1
Chapter 2: Literature Review.....	4
2.1 The commercial Thoroughbred breeding industry	4
2.2 Commercial breeding season versus natural breeding season in the mare	4
2.3 Seasonality of pasture	7
2.3.1 Energy content of pasture	8
2.3.2 Pasture chemical composition	10
2.3.3 Pasture availability	11
2.4 Energy supply versus energy demand of mares under pastoral system.	14
2.5 Dry matter intake	17
2.6 Energy systems	20
2.7 Published recommendation for nutrient requirements for horses.	21
2.8 Estimating energy requirements in mares	24
2.8.1 Pregnancy	24
2.8.1.1 The fetal growth models	24
2.8.1.2 Non-fetal tissue accretion.....	26
2.8.1.3 Oxygen consumption of conceptus	26
2.8.1.4 Chemical composition of conceptus	26
2.8.1.5 Pregnancy requirements.....	26
2.8.2 Lactation.....	28
2.8.2.1 Milk production.....	28
2.8.2.2 Gross energy of milk.....	30
2.8.2.3 Lactation Requirements	31
2.9 Nutritional regulation of reproduction	32
2.9.1 Body condition score and energy intake.....	32
2.9.2 Leptin	32
2.9.3 Plasma insulin growth factor (IGF-1).....	33
2.10 Reproductive performance of Thoroughbred mares in New Zealand.....	34
2.11 Concepts of modelling.....	37
2.11.1 Definition of modelling.....	37

2.11.2 Types of model	37
2.11.3 Steps in building a model	38
2.11.3.1 Purpose of modelling	38
2.11.3.2 Defining the energy balance system for pregnant and lactating Thoroughbred mares managed on pasture (Figure 10).....	40
2.11.4 Testing for robustness	40
2.12 Conclusion	42
Chapter 3: Deterministic modelling of energy supply and demand for foaling mares managed at pasture on commercial Thoroughbred stud farms.....	44
3.1 Methods.....	44
3.1.1 The model	44
3.1.2 Model simulation	50
3.2 Results	51
3.2.1 Modelling of energy balance pattern (Main analysis).....	51
3.2.2 Modelling of energy balance pattern using lower extra energy expenditure at 25% above maintenance.	53
3.2.3 Effect of mare bodyweight on modelled energy balance.....	56
3.2.4 Effect of foaling date on modelled energy balance.....	56
3.2.5 Effects of pasture metabolizable energy (ME) and dry matter intake (DMI).	59
3.2.6 Sensitivity Analysis	63
3.3 Discussion	65
3.3.1 Pre-partum energy balance.....	65
3.3.2 Post-partum energy balance.....	66
3.3.3 Energy balance and mobilization of body composition.....	66
3.3.4 Foaling date and modelled energy balance.....	67
3.3.5 Post-partum energy balance and reproductive performance.....	68
3.3.6 Modelled energy balance and leptin levels	69
3.3.7 Energy expenditure level and modelled energy balance	69
3.3.8 Limitations.....	70
3.3.9 Sensitivity analysis	71
3.4 Conclusion.....	73
Chapter 4: General discussion	74
4.1 Summary of findings	74
4.2 Limitations	74
4.3 Future studies	75
4.4 Implications and conclusion.....	76
References.....	78

List of Tables

Table 1. Average pasture energy content for equine and dairy pasture.	9
Table 2. Seasonal variation in chemical composition of equine pasture in New Zealand from Hirst (2011).	10
Table 3. Estimated dry matter intake (DMI) of horses fed different types of forages and supplements.	19
Table 4. Daily requirement of a 500 kg mare at her 11th month of pregnancy and 1 st month of lactation estimated based on the German, French (INRA) and NRC systems.	28
Table 5. Reference value for milk production and composition for estimating lactation requirements. Figure from INRA (2015).	32
Table 6. First cycle pregnancy rate (FCPR), end-of-season pregnancy rate (SPR) and foaling rate of five Thoroughbred stud farms in the Waikato region of New Zealand over three consecutive breeding season (2006-2008) from Hanlon <i>et al.</i> (2012b).	35
Table 7. Pregnancy loss rates and multiple pregnancy rate by reproductive status for Thoroughbred mares from the Waikato region of New Zealand over three consecutive breeding season (2006-2008) from Hanlon <i>et al.</i> (2012b).	35
Table 8. The organisational hierarchy of biology as described by Thornley and France (2007).	37
Table 9. Equations used to model pregnancy tissue accretion, milk yield, digestible energy (DE) requirements, energy intake and energy balance for Thoroughbred mares during pregnancy and lactation.	46
Table 10. Estimated voluntary dry matter intake (DMI) of horses fed different types of forages and supplements.	49
Table 11. Efficiency of digestible energy (MJ DE) utilization in pregnant and lactating mares, and effects of diet.	50
Table 12. Default values for dry matter intake, pasture metabolizable energy (ME) and mare body weight (BWT) in the sensitivity analysis of the energy balance model for foaling and lactating Thoroughbred mares.	51
Table 13. Total energy balance (MJ ME), total energy requirements (MJ ME) and total energy balance expressed as percentage of requirements (%) of Thoroughbred mares managed at pasture under commercial conditions with different initial bodyweight (BWT) (kg) and relative foaling date (FD) during last three months of pregnancy and during the first 150 days of lactation.	55
Table 14. Maximum positive energy balance, maximum negative energy balance at any point during the pre and post-foaling period modelled (day -90 to day 150) for Thoroughbred mares managed at pasture under commercial conditions with different initial bodyweight (BWT) (kg)	

and relative expected foaling date (FD), presented in percentage of requirement and total relative change between two points..... 59

Table 15. Sensitivity of energy balance (MJME/day), energy intake (MJME/day) and energy requirements (MJME/day) of Thoroughbred mares managed at pasture under commercial conditions to changes in pasture metabolizable energy (ME), body weight (kg), relative expected foaling date, and dry matter intake (% initial body weight) from default values²..... 64

List of Figures

Figure 1. Proportion of mares ovulating each month over a 12 month period in South Eastern Australia. Data obtained from Sydney horse abattoir. Hatched area represents commercial breeding season. Figure from Hanlon (2012).	6
Figure 2. Perennial dairy pasture yield in different regions of New Zealand a) Waikato, b)Wairarapa, c) Canterbury-dry, d) Canterbury-irrigated, e) Southland, f) Central Otago (Valentine and Kemp 2007).....	12
Figure 3. Four year actual pasture yield and seven year mean pasture yield (thick bold line) on a Manawatu dairy farm, showing year to year variation in seasonal and annual pasture production (Litherland and Lambert 2007).....	13
Figure 4. a) Energy requirement of a 500 kg mare foaling at different times of the year (MJ DE/day) versus the pasture metabolisable energy available (MJ ME/kg DM); b) Annual pasture growth rate on Dairy 1, Massey University, Palmerston North in 2015-2016 (Correa 2015 - 2017, unpublished); Energy requirement obtained from NRC (2007); Pasture metabolizable energy obtained from Litherland & Lambert (2007).	15
Figure 5. Feed energy partitioning. Information adapted from Waghorn <i>et al.</i> (2007) and NRC (2007).....	20
Figure 6. Lactation curves for nursing or milked mares and milked or nursed cows calculated using Wood's model (Doreau and Martuzzi 2006). Figure from INRA (2015).	29
Figure 7. Daily milk yield of mares statistically transformed to gram per kg body weight ($x =$ day of lactation) (Coenen <i>et al.</i> 2011).....	29
Figure 8. Mean daily milk production in mares according to stage of lactation and breed. Figure from Chavatte (1997).....	29
Figure 9. Variation in composition of mare's milk gross energy with stage of lactation. Figure from INRA (2015).	30
Figure 10. Steps in building a biological model using the Energy Balance system for pregnant and lactating Thoroughbred mares as an example. Information adapted from Wastney (1999), Gold (1977).....	39
Figure 11. Pasture metabolisable energy (ME) of typical New Zealand pasture used in the energy balance model adapted from (Litherland and Lambert 2007), and growth rate of pasture collected from Dairy 1, Massey University, Palmerston North (Correa 2015 - 2017, unpublished).....	48
Figure 12. Pre and post-partum changes in energy balance (MJME/day) of a Thoroughbred mare (BWT=500kg) managed at pasture under commercial conditions, foaling at 0 days (FD0), 15 days (FD15), 50 days (FD50), 90 days (FD90) after 1 st September when extra energy expenditure is 45% above maintenance.	52

Figure 13 Pre and post-partum changes in energy balance (MJME/day) of a Thoroughbred mare (BWT=500kg) managed at pasture under commercial conditions, foaling at 0 days (FD0), 15 days (FD15), 50 days (FD50), 90 days (FD90) after 1st September when extra energy expenditure is 25% above maintenance.....	54
Figure 14. Effect of bodyweight (kg) on modelled pre-partum energy balance (MJME/day) in a Thoroughbred mare (at differing bodyweights) managed at pasture under commercial conditions and foaling at 0 days (FD0), 15 days (FD15), 50 days (FD50), 90 days (90) after 1 st September.	57
Figure 15. Effect of bodyweight (kg) on modelled post-partum energy balance (MJME/day) in a Thoroughbred mare (at differing bodyweights) managed at pasture under commercial conditions and foaling at 0 days (FD0), 15 days (FD15), 50 days (FD50), 90 days (FD90) after 1 st September.....	58
Figure 16. Relationship between average energy balance (MJME/day) and relative expected foaling date (0 to 90 days after 1st September) during (a) pregnancy and (b) lactation in Thoroughbred mares (500 kg BWT).	60
Figure 17. Pasture energy available (MJME/kg DM) and pasture energy intake (MJME/day) of a Thoroughbred mare (500 kg BWT) managed at pasture under commercial conditions, foaling on different relative expected foaling date (FD) (days after 1 st September), with dry matter intake at 2% during pregnancy and 2.5% body weight during lactation.	61
Figure 18. Energy intake and energy requirements of Thoroughbred mare (500 kg BWT) managed at pasture under commercial conditions, foaling on different relative expected foaling date (FD) (days after 1 st September), consuming dry matter at (a) 2% body weight during pregnancy and (b) 2.5% body weight during lactation.....	62