

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

INTERACTIONS BETWEEN
FLAVOUR COMPOUNDS AND
MILK PROTEINS

A THESIS
PRESENTED IN PARTIAL FULFILMENT OF THE
REQUIREMENTS FOR THE DEGREE OF
MASTER OF PHILOSOPHY
IN FOOD TECHNOLOGY
AT MASSEY UNIVERSITY, PALMERSTON NORTH,
NEW ZEALAND

XIANG QIAN ZHU
2003

ACKNOWLEDGEMENTS

I would like to thank my chief supervisor, Professor Harjinder Singh, for the many hours of discussion and guidance throughout the length of this project. I am also thankful for the support of my co-supervisors, Dr Rogerio Pereira and Professor Peter Munro. I would also like to thank Dr Owen Mills (Fonterra Research Center, Palmerston North) for kindly providing flavour compounds and making time to discuss aspects of this project. Thanks also to Dr Mike Taylor for him kindly taking time to discuss the solubility of sodium palmitate.

I would also like to thank Mr. John Sykes for his kindly help in setting up GC machine in the beginning of my project. I am grateful for the help and support the staff and students in dairy cluster have given me throughout this project, especially Dr Aiqian Ye, Mr. Warwick Johnson, Mr. Steve Glasgow, Ms Karen Pickering, Maya Sugiarto, Jian Cui, Yiling Tan and Kelvin Goh.

I would like to sincerely thank the staff in First Aid Office and Reception of Institute of Food, Nutrition and Human Health for their kindly help during my project.

Finally, I would like to express my sincere gratitude to my family- my wife, Yanli, for her love, supporting and encouragement throughout this thesis; my parents and my son.

TABLE OF CONTENTS

CHAPTER ONE: INTRODUCTION	1
CHAPTER TWO: LITERATURE REVIEW	3
2.1 Flavour-matrix interaction	3
2.2 Interactions of volatile flavours in aqueous solution	3
2.3 Mechanism of binding of ligands to β -lactoglobulin	5
2.3.1 The structure of β -lactoglobulin and the binding sites	5
2.3.2 Effect of pH on ligand binding	7
2.4 Equilibrium binding phenomena	9
2.5 Published results on flavour-protein binding	11
2.6 Analysis methods	14
2.6.1 Fluorescence	14
2.6.2 Affinity chromatography	15
2.6.3 Headspace techniques	17
2.6.4 Equilibrium dialysis	17
2.7 Solid-phase microextraction technique	18
2.7.1 Equilibrium sampling	19
2.7.2 Dynamic sampling (rapid sampling)	21
2.8 Conclusions	26
2.9 Objectives	27
CHAPTER THREE: MATERIALS AND METHODS	28
3.1 General materials and equipment	28

3.2 Gas chromatography	29
3.3 Headspace solid-phase microextraction	29
3.4 Standard solutions	31
3.4.1 Flavour compounds stock solution	31
3.4.2 External standard method	31
3.4.3 Protein solutions	31
3.4.4 Flavour-protein mixed solutions	32
3.5 Determination of analyte retention times	32
CHAPTER FOUR: OPTIMISATION OF SOLID-PHASE MICROEXTRACTION CONDITIONS FOR FLAVOUR BINDING ANALYSIS	35
4.1 Introduction	35
4.2 Materials and methods	35
4.3 Results and discussion	35
4.3.1 The size of containers	35
4.3.2 The type of fibre and extraction time	37
4.4 Conclusions	43
CHAPTER FIVE: BINDING OF FLAVOUR COMPOUNDS TO MILK PROTEINS USING SOLID-PHASE MICROEXTRACTION	45
5.1 Introduction	45
5.2 Materials and methods	46
5.3 Results and discussion	47
5.3.1 The effect of ethanol addition	47

5.3.2 The equilibrium binding time	50
5.3.3 Flavour binding to sodium caseinate	52
5.3.4 Flavour binding to whey protein isolate	55
5.4 Conclusions	64
CHAPTER SIX: EFFECT OF pH AND SODIUM PALMITATE ADDITION ON BINDING OF 2-NONANONE TO WPI	66
6.1 Introduction	66
6.2 Materials and methods	66
6.2.1 Preparation of solutions	66
6.3 Results and discussion	67
6.3.1 Effect of pH on WPI-2-nonanone binding	67
6.3.2 Effect of sodium palmitate on WPI-2-nonanone binding	74
6.4 Conclusions	80
CHAPTER SEVEN: CONCLUSIONS AND FURTHER DIRECTIONS	82
REFERENCE	84
APPENDIX A: CALCULATION RESULTS FOR KLOTZ PLOT	96
APPENDIX B: SIMULATION OF DISSOCIATION OF β-LACTOGLOBULIN	99

LIST OF FIGURES

Figure	Title	Page
2.1	Schematic representation of the tertiary structure of bovine β -lactoglobulin	6
2.2	The effect of pH on the quaternary structure of β -lactoglobulin	7
2.3	Modification of aroma compound retention according to structural modifications of β -lactoglobulin with pH between 3 and 11.	8
2.4	Hummel and Dreyer type chromatograms obtained for 40 and 80 ppm ethyl benzoate concentration in the eluent	16
2.5	Various components of the SPME device	20
2.6	Temperature dependence of the adsorption versus time profiles obtained for methamphetamine	22
2.7	Demonstration of myoglobin (M) and cytochrome (C) displacement by lysozyme (L) with time (A) and protein extraction using a polyacrylic- acid-coated fibre (B).	22
2.8	Adsorption curve under different convection conditions	23
2.9	Boundary layer model	24
2.10	SPME configuration	25
3.1	Typical GC response (ethanol + 2-heptanone)	32
3.2	Typical GC response (ethanol + 2-nonanone)	33
4.1	SPME results for different sub-samples of 0.6 mM 2-heptanone standard solution	37
4.2	Effect of extraction time at 25°C for 2-heptanone on the SPME results	39
4.3	Effect of extraction time at 25°C for 2-nonanone on the SPME results	39
4.4	GC response curve of 2-heptanone standard solution	40
4.5	GC response curve of 2-nonanone standard solution (7 μ m PDMS coating fibre, extraction time 5 minutes)	41

List of figures

4.6	GC response curve of 2-nonanone standard solution (30 μm PDMS coating fibre; extraction time: 5 min)	42
4.7	GC response curve of 2-nonanone standard solution (30 μm PDMS coating fibre, extraction time: 30 seconds)	43
5.1	The effect of the addition of ethanol to 0.2 mM 2-nonanone standard solution on the GC response	48
5.2	The effect of ethanol addition to a flavour--protein solution on the GC response	49
5.3	The effect of ethanol addition to a flavour--protein solution on the GC response (0.2 mM 2-nonanone in 2% WPI solution; 30 μm PDMS fibre; extraction time: 5 min)	50
5.4	The interaction between WPI and 2-heptanone with time (0.4 mM 2-heptanone solution; 100 μm PDMS fibre; extraction time: 5 min)	52
5.5	The interaction between WPI and 2-heptanone with time (0.8 mM 2-heptanone solution; 100 μm PDMS fibre; extraction time: 5 min)	52
5.6	The interaction between WPI and 2-nonanone with time (0.2 mM 2-nonanone solution; 30 μm PDMS fibre; extraction time: 5 min)	53
5.7	The interaction between WPI and 2-nonanone with time (0.2 mM 2-nonanone solution; 30 μm PDMS fibre; extraction time: 30 seconds)	53
5.8	SPME results for the binding of 2-heptanone to Na-CN (100 μm PDMS fibre; extraction time: 5 min)	54
5.9	Figure 5.9. SPME result for the binding of 2-nonanone to Na-CN	55
5.10	Klotz plot for 2-nonanone binding to Na-CN	56
5.11	SPME result of the binding of 2-heptanone to WPI (100 μm PDMS fibre; extraction time: 5 min)	57
5.12	SPME result of the binding of 2-nonanone to WPI (30 μm PDMS fibre; extraction time: 5 min)	57
5.13	SPME result of the binding of 2-nonanone to WPI (30 μm PDMS fibre; extraction time: 30 seconds)	58
5.14	Amount of bound 2-nonanone at different WPI concentrations	58
5.15	Klotz plot for 2-heptanone binding to WPI	60
5.16	Klotz plot for the binding of 2-nonanone to WPI at WPI concentrations $\leq 0.5\%$	61

List of figures

5.17	Klotz plot for the binding of 2-nonanone to WPI at WPI concentrations $\geq 1\%$	62
5.18	Klotz plot for 2-nonanone binding to β -lactoglobulin in WPI, assuming that all the binding occurred with β -lactoglobulin alone at WPI concentrations $\leq 0.5\%$	63
5.19	Klotz plot for 2-nonanone binding to β -lactoglobulin in WPI, assuming that all the binding occurred with β -lactoglobulin alone at WPI concentrations $\geq 1\%$.	64
5.20	The proportion of monomer β -lactoglobulin as a function of the protein concentration (simulated data using an association constant $K = 4.88 \times 10^4 \text{ M}^{-1}$)	65
6.1	The effect of pH on the binding of 2-nonannone to WPI (1%)	68
6.2	Klotz plot for 2-nonanone and WPI binding at pH = 3	69
6.3	Klotz plot for 2-nonanone and WPI binding at pH = 4	69
6.4	Klotz plot for 2-nonanone and WPI binding at pH = 6	70
6.5	WPI–2-nonanone binding parameters at different pH values	71
6.6	Klotz plot for binding of 2-nonanone to β -lactoglobulinbinding in WPI solution at pH = 4	72
6.7	Klotz plot for binding of 2-nonanone to β -lactoglobulin in WPI solution at pH = 6	73
6.8	The effect of sodium palmitate on the binding of 2-nonanone to WPI	76
6.9	The effect of added sodium palmitate on β -lactoglobulin–2-nonanone binding	78
6.10	Comparison of the competition model and experimental results (0.5% WPI, 0.2 mM 2-nonanone)	81

LIST OF TABLES

Table	Title	Page
2.1	Comparison of binding constants obtained with different methods	12
3.1	Temperature and conditioning recommendations for fibres	30
3.2	Retention times for the two components in standard solutions	33
3.3	Retention times of individual components	34
4.1	Container and liquid amount used for optimization of SPME conditions	36
4.2	The SPME-GC results obtained using different containers	36
4.3	The SPME results for 2-heptanone standard samples	40
4.4	The SPME results of 2-nonanone standard samples	41
6.1	WPI–2-nonanone binding parameters at different pH values	70
6.2	β -Lactoglobulin–2-nonanone binding parameters at different pH values	72
6.3	Bound amount of 2-nanonone in different sodium palmitate added	80