

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

STUDIES ON THE EFFECTIVENESS OF VARIOUS INSECTICIDES AND
DRYING IN CONTROLLING GRANARY WEEVIL
(Sitophilus granarius L.)
IN STORED WHEAT SEED

A thesis presented in partial fulfilment
of the requirement for the degree of
Master in Agricultural Science (Seed Technology)
at
Massey University, New Zealand

MD. NURUL ISLAM
March 1986

ABSTRACT

This study was designed to examine the effects of various insecticidal materials and of repeated seed drying during storage on the population dynamics and survival of Granary Weevil (Sitophilus granarius L.) in wheat. The study was conducted in three parts. The first experiment examined the immediate and longer term effectiveness of the contact insecticide malathion and the fumigant phosphine along with influence of repeated drying to safe moisture contents on granary weevil infestation. Wheat seed was stored for 165 days at 25 C and 80% R.H. Treatments were examined every 30 days to study the development of Sitophilus granarius populations and associated damage to wheat seed and evaluation made of the effects of drying, malathion and phosphine on the established infestation and on seed quality. Granary weevils increased by about a factor of X 10 every 60 days and did extensive damage to chemically untreated seed. Repeated drying reduced the rate of increase but did not eliminate the insect population. Malathion dust added to infested seed severely checked insect development and when combined with drying destroyed the infestation completely. Malathion also displayed considerable residual effect and had no adverse effect on seed viability. Phosphine was found to be totally effective in eradicating an established population of granary weevils from seed without affecting seed quality.

In a second experiment malathion was sprayed onto jute squares at 2.5% and at one half and one quarter of this rate. Treated squares were stored for 90 days at 20 C, ambient RH of 70 - 90% or 30 C, ambient RH of 60 - 80% and the residual toxicity of the deposit was assayed with live insects at intervals after treatment. Malathion was also applied at 2.5% concentration to the outside of grain filled sacks which were then placed individually into large plastic bags into which adult granary weevils were introduced at 7 day intervals. After 56 days storage, counts were made of live and dead insects inside the sacks to assess protective effect of the malathion treatment. On jute squares malathion was completely effective at all concentrations and at both storage temperatures (20 C and 30 C) immediately after application and also after 7 days. Thereafter, it lost its effectiveness slowly over the next 90 days storage. In whole sack treatment malathion was found to provide only immediate protection at both temperatures and was inadequate after only a few days.

In the third experiment wheat seeds, uninfested and infested with Sitophilus granarius, were mixed with ground neem seed of each of two species of neem (Azadirachta indica and Melia azadirach) at 1 g per 20 g wheat and were stored at 25 C and 80% R.H. Seeds were examined for live and dead insects and germination assessed after 90 days storage. Little or no direct mortality of adults was recorded but there was indirect evidence of suppression of egg laying particularly with Azadirachta indica. Neem seed powder did not affect the viability of the wheat seed.

This study has clearly shown the short term residual effectiveness of malathion, the immediate eradicator action of phosphine and the poor performance of the natural insecticidal chemical in neem seed on granary weevil infestation in wheat. The results also show the maintenance of low seed moisture contents in wheat to be a practical method of reducing insect populations. The role of granary weevil in damaging seed was clearly seen by X-ray photography and by the extent of types of abnormal seedlings found in positional germination tests. In the absence of effective control Sitophilus granarius has the potential to devastate wheat seed quality in terms of both purity and germination in as little as 90 days.

ACKNOWLEDGEMENT

I wish to express my heartfelt gratitude to my supervisors Dr P.G. Fenemore, Dr M.J. Hill and Mr E. Roberts for their invaluable help, encouragement, constructive criticisms and advice in the preparation of this thesis.

I am also greatly indebted to Dr M.J. Hill, Director, Seed Technology Centre for his concern for my personal welfare during my study period in New Zealand.

My sincere appreciation is also due to:

Mrs D.E.M. Meech for her supervision in laboratory work.

Dr I.L. Gordon for his help and suggestions on the computer analysis of the experimental data.

Mr C.R. Johnstone for the preparation of seed samples and arranging storage facilities.

Mrs K.A. Johnstone for her technical assistance in the laboratory work.

Mrs A.M. Davies for her untiring and excellent work in typing the draft and final manuscript of the thesis.

The staff of the central photographic unit, Massey University for their assistance with photography.

All friends for their help and encouragement. Fred, my friend, deserves my heartiest gratitude for his direct help in computing and tidying up of the thesis.

I also wish to thank the Government of Bangladesh and Bangladesh Agricultural Development Corporation for allowing me to study in New Zealand.

Finally my special thanks and appreciation to my parents, wife and children for their understanding, patience and encouragement during my stay in New Zealand.

CONTENTS

	<u>Page</u>
ABSTRACT	ii
ACKNOWLEDGEMENTS	iv
LIST OF TABLES	vii
LIST OF FIGURES	viii
LIST OF PLATES	ix
LIST OF APPENDICES	x
INTRODUCTION	1
CHAPTER 1 - REVIEW OF LITERATURE	4
Introduction	4
Insect Pest of Stored Seed	4
Granary Weevil	5
Control of Insect Pests in Stored Seed	9
Sanitation and Management	9
Physical Measures	9
Chemical Measures	11
Malathion	13
Phosphine	16
Neem Seed	19
Effects of Storage Conditions on Seed Longevity	21
CHAPTER 2 - AN EVALUATION OF MALATHION DUST, FUMIGATION WITH PHOSPHINE AND SEED DRYING IN CONTROLLING AN ESTABLISHED INFESTATION OF <u>Sitophilus granarius</u> AND THEIR EFFECT ON SEED QUALITY.	25
Introduction	25
Materials and Methods	25
Results and Discussion	34
Initial Quality Level of Wheat Seed	34
Insect Population	35
Seed Moisture Content	41
Normal Germination	44
Abnormal Seedlings	53
Types of Abnormal Seedlings	55
Internal Insect Damage to Seed	56

	vi
CHAPTER 3 - EFFECTIVENESS OF MALATHION APPLIED AS SPRAY TO THE OUTSIDE OF STORAGE SACKS AS A PREVENTATIVE MEASURE AGAINST INFESTATION.	71
Introduction	71
Materials and Methods	71
Results and Discussion	75
Effectiveness of Malathion Sprayed onto Jute Squares	75
Effectiveness of Malathion Sprayed onto Jute Sacks	77
CHAPTER 4 - EVALUATION OF GROUND NEEM SEED AS AN INSECTICIDE.	80
Introduction	80
Material and Methods	80
Results and Discussion	83
Effect of Ground Neem Seed on an Established Insect Population	83
Seed Quality	86
CHAPTER 5 - GENERAL DISCUSSION AND CONCLUSION	89
BIBLIOGRAPHY	94
APPENDICES	

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Initial quality of wheat seed	34
2. Numbers of live and (dead) adult granary weevils in the different treatments up to 165 days of storage (mean)	36
2a. Numbers of live and (dead) adult granary weevils in malathion and phosphine treatments upto 165 days storage (mean)	37
3. Percentage of normal germination of seeds from different treatments upto 165 days storage (mean)	46
4. Categories of abnormal seedlings of wheat seed in positional germination test after X-ray photography	70
5. Numbers of granary weevil adults affected out of 20 on jute squares at intervals after treatment	76
6. Numbers of live and (dead) granary weevil adults in untreated seed stored in malathion treated sacks, 56 days after infestation	78
7. Numbers of live and (dead) granary weevil adults in different treatments upto 90 days storage (mean)	84
8. Percentages of normal germination of wheat seeds of all treatments upto 60 days storage.	87

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Insect population development in different treatments with time	38
2. Changes in seed moisture content in undried treatments	42
3. Changes in seed moisture content in dried treatments	43
4. Changes in germination percentage of different treatments during storage	47
5. Changes in abnormal seedling percentage in different treatments during storage	54
6-13. Positional germination results of seeds tested in the same sequence as shown in Plates 13 to 20	59-67

LIST OF PLATES

<u>Plate</u>	<u>Page</u>
1. Vacuum insect counter	27
2. Containers for storing wheat seed	27
3. Large plastic drums used to hold one complete replicate in experiment 1	29
4. Pump for circulating air through large drums	29
5. Aspirated wet and dry bulb thermometers	31
6. Agromatic portable moisture meter	31
7. Physical appearance of uninfested untreated wheat seed after 120 days of storage	50
8. Physical appearance of insect infested and malathion treated wheat seed after 120 days of storage	50
9. Physical appearance of insect infested and phosphine fumigated seeds after 120 days of storage	51
10. Physical appearance of insect infested and untreated seeds after 120 days of storage	51
11. Abnormal types: From Ia to IIIId	57
12. Abnormal types: From IIIe to VIa	58
13-20 X-ray photographs of seeds of different treatments after 60 days of storage	60-67
21. Granary weevil adults confined beneath petridish on jute square	74
22. Jute sack with wheat seed inside polythene bag	74
23. Neem (<u>Azadirachta indica</u>) seed	81
24. Neem (<u>Melia azedarach</u>) seed	81

LIST OF APPENDICESAppendix

1. Percentage of abnormal seedlings in different treatments up to 165 days of storage (mean).
2. Percentage of main types of abnormalities in wheat seeds after 30, 60, 90, 120 and 165 days storage.
3. Analysis of variance of the experimental data on the number of live granary weevil adults of insect infested treatments after 30, 60, 90, 120 and 165 days storage.
- 3a. Analysis of variance of the experimental data on the number of live granary weevil adults in malathion and phosphine treatments after 30, 60, 90, 120 and 165 days storage.
4. Analysis of variance of the experimental data on normal germination percentage of seeds after 30, 60, 90, 120 and 165 days storage.
5. Analysis of variance of the experimental data on the percentage of abnormal seedlings after 30, 60, 90, 120 and 165 days storage.
6. Analysis of variance of the experimental data on the numbers of live granary weevil adults after 0, 7 and 14 days of malathion treatment.
7. Analysis of variance of the experimental data on the numbers of live granary weevil adults in neem seed trial after 0, 30, 40, 60 and 90 days of storage.
8. Analysis of variance of the experimental data on normal germination of seeds in neem seed trial after 60 days storage.

INTRODUCTION

Shortage of food is a major problem in most developing countries. Low crop yields aggravate this problem. Among the factors that increase the yield of agricultural products, the successful production and availability of high quality seed is most important. Even though many countries particularly in the humid tropics and sub-tropics, have developed the necessary technology for the production of high quality seeds they often face great difficulties in maintaining high seed quality in storage, particularly where lengthy storage periods occur between harvest and the next sowing season.

The combination of high humidity and high temperature environments are disastrous in their effects on the viability of stored seeds (Delouche et al., 1973). Some information on how high temperature and humidity affect the seed viability in storage has been provided by Islam (1984). Dehumidified cold storage is the best solution for storing most types of crop seeds. However, financial constraints in most developing countries, such as Bangladesh at present preclude the commercial use of environmentally conditioned seed stores. A common storage practice now being adopted there is to keep the moisture content of stored seed at low levels by repeated drying.

Under high humidity and temperature storage conditions insects can be one of the most important single causes of damage to seeds if infestation is not prevented or controlled (Henderson and Christensen, 1961). Hall (1970) states that at least 10% of harvested food crops are destroyed by pests in storage, and that current losses of 30% are apparently common in large areas of the world especially in some of the tropical and sub-tropical countries where the need for more food is greatest. High temperature and high relative humidity ambient conditions, are most detrimental to seed longevity in storage and at the same time provide very favourable environments for the development of insect populations. This makes safe seed storage in such countries even more difficult. The present study was undertaken to investigate some aspects of chemical control of insects in stored seed.

In Bangladesh wheat and rice seeds are generally stored on a large scale. The storage of wheat seed however, is generally considered to be more difficult. Weevils are the most common insects that infest stored seeds in Bangladesh. The insecticide malathion and

the fumigant phosphine (liberated from Phostoxin tablets) are the chemicals mainly used there to combat insect infestation of seed. However, little research has been done there on the mode of action of these chemicals or on their effects on seed quality and thus practical use depends on the recommendations of manufacturers or suppliers. This study was therefore, undertaken to investigate several aspects of the action of malathion and phosphine when used on wheat seed infested by granary weevil (Sitophilus granarius) under simulated tropical storage conditions.

The increasing cost of some insecticides and also the rapid development of resistance by insects against a particular insecticide necessitates the search for new but relatively cheap insecticides. Recently neem seed has received wide attention as a possible cheap but effective natural pesticide (Ivbijaro, 1983). Neem is readily available in most tropical and subtropical countries including Bangladesh. This study therefore, included an evaluation of the effectiveness of two botanical species of neem seed (Azadirachta indica and Melia Azedarach) as an insecticide against granary weevil.

In summary therefore, the objectives of the present study were to investigate the following aspects of chemical control of granary weevil (Sitophilus granarius L.) in stored wheat seed (Triticum aestivum).

- (1) To study the development of granary weevil populations and associated damage done to wheat seed in storage.
- (2) To evaluate the effect of seed drying on insect population development and on seed quality.

- (3) To evaluate the effectiveness of the contact insecticide, malathion, the fumigant phosphine, and the natural pesticide neem seed, in controlling an established infestation of granary weevil in wheat seed.
- (4) To study the effectiveness of malathion applied as a spray to the outside of storage sacks as a preventative measure against infestation.
- (5) To investigate the effect of the above chemicals on wheat seed quality.