

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Polymer Dynamics Studied by Dynamic Light Scattering

A thesis presented in partial fulfilment of
the requirements for the degree of
Master of Science
in Physics at
Massey University

by

David Thomas CLARK

1998

Abstract

Theoretical treatments of dynamic properties of polymer solutions are reviewed. Particular emphasis is placed on the discussion of diffusion in polymer solutions. The relationship between the slow diffusion coefficient found by Dynamic Light Scattering (DLS) and the Self Diffusion Coefficient is shown.

An introduction to DLS theory is given. The experimental techniques involved in DLS measurements are discussed.

Concentration dependencies of the DLS observed slow diffusion coefficient in ternary polymer solutions of polystyrene-poly vinyl methyl ether-solvent have been measured. Solvents used were toluene, carbon tetrachloride and tetrachloroethylene and the polymer molar masses were $110\,000\text{ gmol}^{-1}$ for both polystyrene (PS) and poly vinyl methyl ether (PVME). Solvents were chosen to be very nearly isorefractive with PVME. Toluene is very nearly an equally good solvent for the polymer pair PS and PVME, while tetrachloroethylene is equally poor for both polymers. Carbon tetrachloride is an unequal quality solvent for this polymer pair. The results of these DLS measurements are reported.

Four sets of experiments are described. The first is the effect on the relationship between D_I (the slow mode diffusion coefficient found by DLS) and D_S (the self diffusion coefficient) of different mass fractions in the polymer solution. It is found the mass fraction, x , has little effect on the observed diffusion coefficient. Secondly the effect of the polymer-polymer interaction parameter, χ , on the relationship between D_I and D_S is investigated. It is found solutions formed with equal quality solvents have D_I nearly equal to D_S . But solutions formed with unequal quality solvents have D_I considerably less than D_S and that these solutions suffer phase separation at lower concentration. Thirdly the effect of polymer molar mass on the relationship between D_I and D_S is investigated. These are found to be in line with those expected from literature.

Fourthly the considerable discrepancy between D_I and D_S which is manifest in 110 000 g mol^{-1} PS/ 110 000 g mol^{-1} PVME/ toluene solutions at polymer volume fractions greater than 0.4 is investigated. The diffusion coefficient found, D_I , does not fit a unique concentration power law. Two unique regions are seen with concentration exponents of -11 ± 4 and -14.9 ± 0.7 for the low and high concentration regions respectively. This is found to agree with results found in literature.

Acknowledgments

I wish to express my sincere gratitude to the following people for their involvement in this work:

Firstly to Jesus, the Author and Perfector of my faith, who gives people a second chance. To you goes the glory.

My supervisor Assoc. Professor Neil Pinder, for providing both theoretical and experimental "know how", and infinite patience. Your enthusiasm was greatly appreciated, as was your advice and proof reading.

The Staff and Students of the Dept. of Physics who have greatly encouraged and enthused myself for the things of physics as well as put up with me for more than a decade.

Dr Hedwig of the Dept. of Chemistry for the use of his warm room and balance.

The Mechanical Workshop Staff for construction of the high concentration transferral apparatus.

GAF for supplying the poly vinyl methyl ether used in this work free of charge. Having worked with this polymer, I now see why, it sticks everywhere.

My Mum and Dad who have provided the example and the enthusiasm to achieve anything. Thanks.

And of course Kerry Wallace, my girlfriend at the time of writing, whose unfailing love and support provides an all-win situation in everything I do. Who patiently proof-read this manuscript, and whose proofed manuscripts I always looked forward to receiving.

Table of Contents

Abstract	ii
Acknowledgements	iv
Table of Contents	v
List of Figures	x
List of Tables	xiii
Chapter 1 Introduction	1
1.1 Polymers and Dynamic Light Scattering	1
1.2 Thesis Organisation	3
Chapter 2 Polymer and DLS Theory	4
2.1 Polymer Outline	4
2.1.1 Polymers-What are they?	4
2.1.2 Polymeric Structure	5
2.2 Properties of Polymers in Solution	7
2.2.1 Compatibility of Polymers	7
2.2.2 Ideal Polymer Chain Model	11
2.2.3 Excluded Volume and the Self Avoiding Walk	12
2.2.4 Theta Solutions	14
2.2.5 Flory-Huggins Mean Field Theory	15

2.2.6	Polymer Motion	20
2.2.6a	The Rouse Model	20
2.2.6b	The Zimm Model	24
2.2.6c	Reptation	29
2.2.6d	The Blob Model	33
2.3	Basic Dynamic Light Scattering Theory	34
2.4	DLS Theory-Ternary Polymer Solutions	40
2.4.1	Benmouna et al Mean Field RPA Theory	40
2.4.2	Extentions to Benmouna Theory	46
2.4.2a	Ternary systems containing polymers of equal molar mass and a solvent of equal quality	51
2.4.2b	Ternary systems containing polymers of unequal molar mass and a solvent of unequal quality	53
2.4.3	Ternary solutions close to phase separation	54
Chapter 3	Experimental Set-up	61
3.1	Dynamic Light Scattering	61
3.1.1	Historical Overview	61
3.1.2	The Dynamic Light Scattering System	61
3.1.3	Laser Operation	63

3.1.4	Spectrometer and Photomultiplier	64
3.1.4a	Refractive Index Matching Bath	64
3.1.4b	Spectrometer Goniometer and Photomultiplier	65
3.1.4c	Optical Alignment of the Spectrometer	65
3.1.5	The Autocorrelator	68
3.1.6	Computer and Software	71
3.1.7	Running Samples	74
3.2	Sample Preparation	76
3.2.1	Glassware cleansing procedures	76
3.2.2	Polymers used in this study	78
3.2.2a	Polymer characteristics	78
3.2.2b	Fractionation of PVME	80
3.2.2c	PVME Handling	81
3.2.3	Solvents used in this study	82
3.2.4	Sample filtration	83
3.2.5	Solution Preparation	84
3.2.6	Sample delivery and storage	85
3.2.7	Weighing	85

Chapter 4	Results and Conclusion	88
4.1	Solutions Studied and Sundry Information	88
4.1.1	Solutions Studied	88
4.1.2	Sundry information about results	93
4.2	Results and Discussion	95
4.2.1	Investigation of the variation of D_I with Mass Fraction	95
4.2.2	Investigation of the variation of D_I with the Flory Polymer-Polymer Interaction Parameter, χ	100
4.2.3	Investigation of the variation of D_I with Molar Mass	105
4.2.4	Investigation into the discrepancy between D_I and D_S at high polymer concentrations	111
Chapter 5	Conclusion	116
5.1	Conclusion	116
5.2	Suggestions for Further Work	120
Appendices: Publications		121
Appendix 1	New Zealand Institute of Physics Conference Poster (1992) Dynamic Light Scattering from Ternary Polymer Solutions	122

Appendix 2	Studies of Ternary Polymer Solutions by Dynamic Light Scattering and Pulsed Field Gradient Nuclear Magnetic Resonance <i>Macromolecular Reports</i> , A31 (SUPPLS, 6 & 7), 1119-1126 (1994)	135
References		144

List of Figures

Chapter 2 Polymer and DLS Theory

Figure 2.2.1	Free Energy of Mixing for Binary mixture versus their composition	8
Figure 2.2.2	Phase diagram of a binary system showing the Upper Critical Point	9
Figure 2.2.3	Phase diagram of a binary system showing the Lower Critical Point	9
Figure 2.2.4	Temperature-concentration diagram for a polymer solution far from LCP according to Daoud and Jannick	10
Figure 2.2.5	A Representation of the Rouse Model beads and spring	20
Figure 2.2.6	A single polymer entangled in the polymeric matrix of other polymers	29
Figure 2.2.7	Reptation-The Rouse Chain in a Tube	31
Figure 2.3.1	Typical scattering geometry in a Dynamic Light Scattering Experiment	34

Chapter 3 Experimental Set-up

Figure 3.1.1	Dynamic Light Scattering Equipment	62
Figure 3.1.2	The Diffusion coefficient of 0.091 μm polystyrene latex solution at various angles of scatter	67

Figure 3.1.3	The first cumulant of a 0.091 μ m polystyrene solution vs q^2	67
Figure 3.2.1	Apparatus for cleaning glassware	77
Figure 3.2.2	The structure of polystyrene	79
Figure 3.2.3	The structure of poly (vinyl methyl ether)	79
Figure 3.2.4	An exploded view of the filtering apparatus	83
Figure 3.2.5	Apparatus for transferring high concentration solutions	85

Chapter 4 Results and Discussion

Figure 4.2.1	The concentration dependence of the diffusion coefficient of PS-PVME-toluene solutions of various mass fractions	96
Figure 4.2.2	The concentration dependence of the diffusion coefficient of PS-PVME-tetrachloroethylene solutions of various mass fractions	98
Figure 4.2.3	The concentration dependence of the diffusion coefficient of PS-PVME-carbon tetrachloride solutions of various mass fractions	99
Figure 4.2.4	The concentration dependence of the diffusion coefficient found by DLS in the PS-PVME system with the solvents toluene, tetrachloroethylene and carbon tetrachloride at a polymer mass fraction of 0.05	103

Figure 4.2.5	The concentration dependence of the diffusion coefficient found by DLS in the PS-PVME system with the solvents toluene and tetrachloroethylene at a polymer mass fraction of 0.12	104
Figure 4.2.6	The concentration dependence of the diffusion coefficient found by DLS in the PS-PVME system with the solvents toluene and carbon tetrachloride at a polymer mass fraction of 0.29	105
Figure 4.2.7	The concentration dependence of the diffusion coefficient obtained by DLS for the PS-PVME system with the solvent toluene at various polymer mass fractions and molar masses	107
Figure 4.2.8	The concentration dependence of the diffusion coefficient obtained by DLS for the PS-PVME system with the solvent carbon tetrachloride at various polymer mass fractions and molar masses	108
Figure 4.2.9	The concentration dependence of the diffusion coefficient obtained by DLS for the PS-PVME system for carbon tetrachloride and toluene at various polymer mass fractions and molar masses	110
Figure 4.2.10	High concentration extension of Daivis's 110 000 gmol^{-1} PS/ 110 000 gmol^{-1} PVME in toluene	113

List of Tables

Chapter 3 Experimental Set-Up

Table 3.2.1 Polymers used in this study 78

Table 3.2.2 Solvents used in sample preparation . . . 82

Chapter 4 Results and Discussion

Table 4.1.1 Sample series investigated 89

Table 4.1.2 Solution series mass and weight fractions . . . 90

Table 4.2.1 Interaction parameter for PS and PVME
in various solvents 101

Chapter 1

Introduction

1.1 Polymers and Dynamic Light Scattering

Since the dawn of the laser, Laser Light Scattering has allowed the investigation of molecular size movements within solutions. Recently ternary polymer solutions containing two polymers and a single solvent have been investigated using Dynamic Light Scattering (DLS). The lack of suitable theory, and apparatus of sufficient quality has limited these investigations to diffusion coefficient dependence on molar mass and concentration. Generally these investigations have utilised special polymers and solvents in order to reduce the complexity of the results found.

Benmouna and Borsali et al [1] [2] [3] [4] have recently produced a theory of diffusion specifically for polymer solutions, which they have extended in ternary polymer solutions. This theory makes several predictions which are easily verifiable by experiment. Daivis [5] has briefly examined various aspects of this theory experimentally, however these results require confirmation and expansion, as well as experimental investigation into recent theoretical advances of this theory.

Daivis's work advanced understanding of diffusion in polymer solutions, however understanding was left incomplete in several areas. In his work Daivis compared diffusion in an equally good solvent for the polymer pair polystyrene (PS) and poly vinyl methyl ether (PVME) with that for the same polymer pair in an unequal quality solvent. The comparison of this diffusion data with data collected with the same polymer pair in an equally poor quality solvent would reveal much about polymer motion in solutions, yet this work was not attempted by Daivis. Daivis noted that the diffusion coefficient measured by DLS deviated from that obtained by Pulse Gradient Spin Echo Nuclear Magnetic Resonance Spectroscopy (PGSE NMR) at high polymer concentrations. This work was not pursued by Daivis. This work investigates aspects of

Daivis's work which have been left incomplete or require further investigation for resolution.

1.2 Thesis Organisation

Chapter 2 summarises existing theories of diffusion in polymer solutions, and DLS theory. This chapter also explains what polymers are and gives a basic understanding of the properties of polymers critical to understanding this thesis.

Chapter 3 summarises experimental procedures for both the DLS equipment and also for solution preparation. This chapter also summarises some of the theoretical background for the DLS equipment, and the sources of polymers and solvents utilised in this work.

Chapter 4 summarises investigations into four aspects of the work previously presented by Daivis et al [5] [6] [7] [8] [9]. The first is the effect on the relationship between D_I and D_S of different mass fractions in the polymer system 110 000 Dalton PS/ 110 000 Dalton PVME/ solvent (where three different solvents were utilised:- toluene, carbon tetrachloride and tetrachloroethylene). Secondly the effect of the polymer-polymer interaction parameter, χ , on the relationship between D_I and D_S is investigated. Thirdly the effect of polymer molar mass on the relationship between D_I and D_S is investigated. Fourthly the considerable discrepancy between D_I and D_S which is manifest in 110 000 Dalton PS/ 110 000 Dalton PVME/ toluene solutions at polymer volume fractions greater than 0.4. The results presented in this chapter are also discussed and compared to results found in literature.

Chapter 5 summarises the main conclusions which can be drawn from this study.