

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

**OVIPOSITION PREFERENCE, LARVAL FEEDING
PREFERENCE AND LARVAL FOOD QUALITY
OF HELIOTHIS ARMIGERA**

A thesis
presented in partial fulfilment of
the requirements for the degree of
Doctor of Philosophy

at
Massey University
Palmerston North
New Zealand

NATHAWUT THANEE

1987

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	i
LIST OF FIGURES	iii
LIST OF TABLES	v
LIST OF PLATES	viii
LIST OF APPENDICES	ix
ABSTRACT	1
CHAPTER 1 INTRODUCTION	3
CHAPTER 2 LITERATURE REVIEW	7
Biology of the genus <u>Heliothis</u>	7
1 Adult	8
2 Egg	10
3 Larva	11
4 Pupa	12
Distribution	15
Economic importance	18
Crop species affected	19
Seasonal development	20
<u>Heliothis armigera</u> (Hubner)	22
1 Distribution and host plants	22
2 Occurrence in Australasia	24
Insect-plant relationships	27
1 Relationships between <u>Heliothis</u> species	

and their food plants	30
2 Oviposition of <u>Heliothis</u> species on plants	32

CHAPTER 3 FIELD OBSERVATIONS OF POPULATION DEVELOPMENT
ON FOUR HOST PLANTS AND SEASONAL DEVELOPMENT

OF HELIOTHIS ARMIGERA IN THE MANAWATU 35

Introduction 35

Field trials 37

Materials and methods 37

1 Field study area 37

2 Plants 40

3 Seasonal occurrence of H.armigera larvae
on host plants in relation to plant phenology 41

4 Damage to tomato fruits by larvae of
H.armigera 42

5 Infestation of aster flowers by larvae of
H.armigera 42

6 Infestation of sweetcorn cobs by larvae of
H.armigera 43

7 Pheromone trapping of H.armigera at
Palmerston North 43

Results 44

1 Seasonal occurrence of H.armigera larvae
on host plants in relation to plant phenology 44

2 Damage to tomato fruits by larvae of
H.armigera 45

3 Infestation of aster flowers by larvae of
H.armigera 45

4 Infestation of sweetcorn cobs by larvae of <u>H.armigera</u>	54
5 Pheromone trapping of <u>H.armigera</u> at Palmerston North	55
Discussion	59
1 Seasonal occurrence of <u>H.armigera</u> larvae on host plants and infestations in relation to plant phenology	59
2 Pheromone trapping of <u>H.armigera</u> at Palmerston North	61
Conclusions	63

CHAPTER 4 OVIPOSITION PREFERENCE OF HELIOTHIS ARMIGERA

ON DIFFERENT PLANTS	64
Introduction	64
Materials and methods	66
1 Oviposition preference of <u>H.armigera</u> for plant parts within plant species (mature flowering plants)	66
2 Oviposition preference of <u>H.armigera</u> between four host plants (immature non-flowering plants)	67
3 Oviposition preference of <u>H.armigera</u> between four host plants (mature flowering plants)	67
4 Effect of plant odour on oviposition preference of <u>H.armigera</u> (mature flowering plants)	68

Results	68
1 Oviposition preference of <u>H.armigera</u> for plant parts within plant species (mature flowering plants)	68
2 Oviposition preference of <u>H.armigera</u> between four host plants (immature non-flowering plants)	78
3 Oviposition preference of <u>H.armigera</u> between four host plants (mature flowering plants)	78
4 Effect of plant odour on oviposition preference of <u>H.armigera</u> (mature flowering plants)	78
Dicussion	81
Conclusions	84

CHAPTER 5 EFFECTS OF DIFFERENT LARVAL FOODS ON

DEVELOPMENT OF <u>HELIOTHIS ARMIGERA</u>	85
Introduction	85
Materials and methods	86
Results	88
1 Effects of different foods on larval development	88
2 Effects of different larval foods on pupal weight, pupal period and sex ratio	92
3 Effects of different larval foods on adult life span and fecundity	95

Discussion	101
1 Larval period	101
2 Percent pupation	102
3 Pupal period	102
4 Pupal weight	103
5 Adult fecundity, life span, oviposition period and egg hatchability	104
6 Ranking and correlation between parameters of larval food quality	105
Conclusions	112

CHAPTER 6 FACTORS DETERMINING FOOD QUALITY FOR

DEVELOPMENT OF <u>HELIOTHIS</u> <u>ARMIGERA</u>	113
Introduction	113
Materials and methods	115
1 Amount of food eaten and larval weight gain	115
2 Dry matter content of food	116
3 Total nitrogen content of food	116
4 Parameters of food utilization	117
Results	118
1 Amount of food eaten and larval weight gain	118
2 Dry matter content of food	120
3 Nitrogen content of food	120
4 Parameters of food utilization	122
Discussion	124
Conclusions	128

CHAPTER 7 FOOD SELECTION AND INDUCTION OF FEEDING

PREFERENCE IN LARVAE OF <u>HELIOTHIS</u> <u>ARMIGERA</u>	129
Introduction	129
Materials and methods	131
1 Preference of newly hatched larvae	
for different foods	133
1.1 Preference tests using leaf discs	133
1.2 Preference for reproductive parts	
of plants	133
2 Induction of feeding preference by	
rearing on a specific food	133
2.1 Rearing on artificial diet and	
plant leaves	133
2.2 Rearing on artificial diet and	
reproductive parts of plants	134
Results	135
1 Preference of newly hatched larvae	
for different foods	135
1.1 Preference tests using leaf discs	135
1.2 Preference for reproductive parts	
of plants	135
2 Induction of feeding preference by	
rearing on a specific food	142
2.1 Rearing on artificial diet and	
plant leaves	142
2.2 Rearing on artificial diet and	
reproductive parts of plants	150

Discussion	155
1 Preference of newly hatched larvae for different foods	155
2 Induction of feeding preference by rearing on a specific food	156
Conclusions	159
CHAPTER 8 GENERAL DISCUSSION	160
1 Seasonal development	160
2 Feeding sites of larvae in the field	164
3 Larval food quality	167
4 Larval feeding preference	170
5 Oviposition preference	173
6 Concluding remarks	176
REFERENCES	177
APPENDICES	201

ACKNOWLEDGEMENTS

The present work was carried out under the supervision of Dr.P.G.Fenemore, to whom I wish to express my gratitude for his encouragement, guidance, continued interest and constructive criticism throughout the course of the study and in the preparation of the manuscript. Grateful acknowledgement is also made to Professor B.P.Springett and Professor B.R. Watkin for their helpful advice, discussion and comment.

I would also like to thank:-

Dr.M.J.Hill for providing plant seeds and allowing the use of his property for light trap.

Dr.P.Singh and Mr.J.S.Dugdale of Entomology Division, DSIR, Auckland for providing information, reference specimens and insect rearing techniques.

Mr.K.Simpson for his advice on Kjeldahl nitrogen content determination.

Dr.I.Gordon and Mr.V.Sereeprasert for their advice on statistical analysis.

Mr.T.Lynch for preparing the experimental sites.

Mrs.L.Mather, Mrs.L.Davis and Mr.H.Neilson, technicians of Plant Health Department for their prompt and efficient help.

Mr.R.Cave for typing this thesis.

The staff of the Massey University Library and Palmerston North MAF Library.

The staff of Massey University Computer Centre.

The staff of Massey University Photographic Unit.

The Meteorology Division, DSIR, Palmerston North for providing meteorological data during experiment.

The SIRATAC Ltd.. New South Wales, Australia for supplying pheromone laminates.

Khon Kaen University, Thailand for giving me this study leave.

New Zealand government for ODA scholarship.

Finally, my family for their encouragement and support.

LIST OF FIGURES

FIGURE	PAGE
2.1 World distribution of <i>Heliothis armigera</i>	23
3.1 Numbers of <i>H.armigera</i> larvae observed every 3 days from December 27, to March 2, 1984-1985 and 1985-1986 on tomato plants.	47
3.2 Numbers of <i>H.armigera</i> larvae observed every 3 days from December 27, to March 2, 1984-1985 and 1985-1986 on aster plants.	48
3.3 Numbers of <i>H.armigera</i> larvae observed every 3 days from December 27, to March 2, 1984-1985 and 1985-1986 on sweetcorn plants.	49
3.4 Mean numbers of <i>H.armigera</i> larvae on different plant parts observed every 3 days from December 27, to March 2, average over the two seasons, 1984-1985 and 1985-1986.	50
3.5 Mean total numbers of <i>H.armigera</i> larvae observed every 3 days between February 4, and March 2, 1985 on different colours of aster flowers.	53
3.6 Relationship between numbers of <i>H.armigera</i> moths caught in pheromone traps and temperature, wind velocity and rainfall at Palmerston North between December 25, 1985 and April 30, 1986.	58
4.1 Mean numbers of eggs laid per female per day by <i>H.armigera</i> at $20 \pm 2^{\circ}\text{C}$ and 16:8(L/D) photoperiod.	70
4.2 Distribution of eggs on different plant parts within plant species.	75

FIGURE	PAGE
4.3 Distribution of eggs between upper halves and lower halves of plants.	76
4.4 Distribution of eggs between upper surfaces and lower surfaces of leaves.	77
4.5 Mean percentages of eggs laid on different plants.	80
7.1 Arrangement for preference tests.	132
7.2 Percentages of newly hatched larvae of <i>H.armigera</i> on leaf discs of four host plants.	139
7.3 Percentages of newly hatched larvae of <i>H.armigera</i> on reproductive parts of plants.	141
7.4 Feeding preference on plant leaves of fourth instar larvae of <i>H.armigera</i> reared on artificial diet.	145
7.5 Feeding preference on plant leaves of fourth instar larvae of <i>H.armigera</i> reared on lucerne and aster leaves.	148
7.6 Feeding preference on plant leaves of fourth instar larvae of <i>H.armigera</i> reared on tomato and sweetcorn leaves.	149
7.7 Feeding preference on reproductive parts of plants of fourth instar larvae of <i>H.armigera</i> reared on artificial diet.	152
7.8 Feeding preference on reproductive parts of plants of fourth instar larvae of <i>H.armigera</i> reared on a specific food.	154

LIST OF TABLES

TABLE	PAGE
2.1 Development of <i>Heliothis armigera conferta</i> on an artificial diet at $25\pm 1^{\circ}\text{C}$ and 16:8(L/D) photoperiod.	14
2.2 Distribution of <i>Heliothis</i> species in the world.	16
3.1 Herbicides, rates and application times to field plots.	41
3.2 Damage to each size category of tomato fruits by <i>H. armigera</i> larvae between February 4, and March 4, 1985.	51
3.3 Mean total numbers of aster flowers and mean numbers of <i>H. armigera</i> larvae per 10 flowers observed between February 4, and March 2, 1985.	52
3.4 Infestation of sweetcorn cobs by larvae of <i>H. armigera</i> between February 4, and March 2, 1985.	54
3.5 Pheromone trap catches of <i>H. armigera</i> moths in experimental plot area between December 1985 and April 1986.	56
3.6 Correlation coefficients(r) and probability(P) between environmental factors and pheromone trap catches of <i>H. armigera</i> .	57
4.1 Percentages of eggs laid of <i>H. armigera</i> on different parts of mature flowering lucerne plants.	71
4.2 Percentages of eggs laid of <i>H. armigera</i> on different parts of mature flowering tomato plants.	72
4.3 Percentages of eggs laid of <i>H. armigera</i> on different parts of mature flowering aster plants.	73

TABLE	PAGE
4.4 Percentages of eggs laid of <i>H.armigera</i> on different parts of mature flowering sweetcorn plants.	74
4.5 Oviposition preference of <i>H.armigera</i> between four host plants.	79
5.1 Cumulative percent mortality of each instar of <i>H.armigera</i> larvae through to pupation when larvae were reared on different plant parts in controlled laboratory conditions.	89
5.2 Effects of different foods on larval development through to pupation.	90
5.3 Effects of shifting larvae from poor to higher quality food after the third moult on larval development and pupation.	91
5.4 Effects of different larval foods on the pupal stage of <i>H.armigera</i> .	93
5.5 Effects of shifting larvae from poor to higher quality food after the third moult on pupal stage of <i>H.armigera</i> .	94
5.6 Effects of different larval foods on adult stage and fecundity of <i>H.armigera</i> .	97
5.7 Effects of shifting larvae from poor to higher quality food on the adult stage and fecundity of <i>H.armigera</i> .	99
5.8 Ranking of different foods according to biological parameters.	108
5.9 Fitness indices for different larval foods.	109

TABLE	PAGE
6.1 Amount of food eaten and total weight gained by <i>H.armigera</i> larvae when fed on 8 different foods.	119
6.2 Dry matter content, water content and nitrogen content of each food.	121
6.3 Parameters of food utilization.	123
7.1 Percentages of newly hatched larvae of <i>H.armigera</i> on leaf discs of four host plants.	137
7.2 Percentages of newly hatched larvae of <i>H.armigera</i> on reproductive parts of host plants.	140
7.3 Feeding preference of fourth instar larvae reared on artificial diet.	144
7.4 Feeding preference of fourth instar larvae reared on specific foods.	146
7.5 Feeding preference of fourth instar larvae raised on artificial diet then offered reproductive parts of plants.	151
7.6 Induction of food preference by fourth instar larvae raised on reproductive parts of plants.	
8.1 Oviposition preference and feeding preference by <i>H.armigera</i> larvae on four host plants.	176

LIST OF PLATES

PLATE	PAGE
3.1 General view of experimental area.	39
4.1 Stems of aster showing soft long hairs and <u>H. armigera</u> eggs.	84(a)

LIST OF APPENDICES

APPENDIX	PAGE
1 Food plants of <i>Heliothis</i> species.	201
2 The composition of the formulated diet for <i>H.armigera</i> .	217
3 Percent of eggs laid by <i>H.armigera</i> on lucerne.	218
4 Percent of eggs laid by <i>H.armigera</i> on tomato.	219
5 Percent of eggs laid by <i>H.armigera</i> on aster.	220
6 Percent of eggs laid by <i>H.armigera</i> on sweetcorn.	221
7 Percent of eggs laid by <i>H.armigera</i> on 4 different plants.	222